



Radiation Measurement Results of 150 items in November 2011



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	ShimohamahaneKawa Akita Pref.	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Mobara, Chiba Pref.	Oct-11	Cs137	4.5 Bq/Kg raw	± 2.7 Bq/Kg raw	4.5	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Nihonmatsu	Oct-11	Cs137	5.6 Bq/Kg raw	± 2.9 Bq/Kg raw	5.6	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Okugawa, Nishiaizu, Yama	Sep-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.9	Cs137	6.0 Bq/Kg raw	
			Cs134	3.9 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice (Koshihikari)	Nishiaizu, Yama	Oct-11	Cs137	4.1 Bq/Kg raw	± 2.8 Bq/Kg raw	4.1	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Furudono, Ishikawa	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.0	Cs137	6.0 Bq/Kg raw	
			Cs134	4.0 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Shioki, Yotsukura, Iwaki	Oct-11	Cs137	5.5 Bq/Kg raw	± 3.0 Bq/Kg raw	9.6	Cs137	6.0 Bq/Kg raw	
			Cs134	4.1 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Kamiogawa, Ogawa, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	6.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Kamiogawa, Ogawa, Iwaki	Oct-11	Cs137	3.7 Bq/Kg raw	± 2.7 Bq/Kg raw	7.0	Cs137	6.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Nakashio, Taira, Iwaki	Oct-11	Cs137	4.9 Bq/Kg raw	± 2.8 Bq/Kg raw	7.3	Cs137	6.0 Bq/Kg raw	
			Cs134	2.4 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Numanouchi, Taira, Iwaki	Oct-10	Cs137	4.0 Bq/Kg raw	± 2.7 Bq/Kg raw	7.0	Cs137	6.0 Bq/Kg raw	
			Cs134	3.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice (Koshihikari)	Tono, Iwaki	Oct-11	Cs137	6.2 Bq/Kg raw	± 3.2 Bq/Kg raw	13.3	Cs137	6.0 Bq/Kg raw	
			Cs134	7.1 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice (MilkyQueen)	Komoda, Kashima, Iwaki	Unknown	Cs137	3.4 Bq/Kg raw	± 2.8 Bq/Kg raw	6.8	Cs137	7.0 Bq/Kg raw	
			Cs134	3.4 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice (Koshihikari)	Shimoyada, Kashima, Iwaki	Oct-11	Cs137	3.5 Bq/Kg raw	± 2.9 Bq/Kg raw	7.2	Cs137	6.0 Bq/Kg raw	
			Cs134	3.7 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Onahamashima, Iwaki	Oct-11	Cs137	3.3 Bq/Kg raw	± 2.8 Bq/Kg raw	5.6	Cs137	6.0 Bq/Kg raw	
			Cs134	2.3 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Soeno, Iwaki	Oct-11	Cs137	4.0 Bq/Kg raw	± 2.9 Bq/Kg raw	8.6	Cs137	6.0 Bq/Kg raw	
			Cs134	4.6 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Watanabe, Iwaki	Sep-11	Cs137	3.5 Bq/Kg raw	± 2.6 Bq/Kg raw	6.7	Cs137	6.0 Bq/Kg raw	
			Cs134	3.2 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice (80% polished)	Oomuro, Taira, Iwaki	Sep-11	Cs137	4.2 Bq/Kg raw	± 2.7 Bq/Kg raw	8.4	Cs137	6.0 Bq/Kg raw	
			Cs134	4.2 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Isohara, Otsuka, Kitaibaraki	Oct-10	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.4	Cs137	6.0 Bq/Kg raw	
			Cs134	2.4 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Isohara, Otsuka, Kitaibaraki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	6.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Kitaibaraki, Ibaraki Pref.	Oct-11	Cs137	2.9 Bq/Kg raw	± 2.8 Bq/Kg raw	5.3	Cs137	6.0 Bq/Kg raw	
			Cs134	2.4 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Hirata, Ishikawa	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.1	Cs137	6.0 Bq/Kg raw	
			Cs134	3.1 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.9	Cs137	6.0 Bq/Kg raw	
			Cs134	3.9 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Akai, Taira, Iwaki	Oct-11	Cs137	2.9 Bq/Kg raw	± 2.8 Bq/Kg raw	6.7	Cs137	6.0 Bq/Kg raw	
			Cs134	3.8 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.8	Cs137	6.0 Bq/Kg raw	
			Cs134	3.8 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Shimoyagyū, Yotsukura, Iwaki	Oct-11	Cs137	3.5 Bq/Kg raw	± 2.7 Bq/Kg raw	6.1	Cs137	6.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Taira, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.7	Cs137	6.0 Bq/Kg raw	
			Cs134	4.7 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Omuro, Taira, Iwaki	Sep-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.3	Cs137	6.0 Bq/Kg raw	
			Cs134	4.3 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Shimokabeya, Taira, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	5.2	Cs137	6.0 Bq/Kg raw	
			Cs134	5.2 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Yoshima, Iwaki	Oct-11	Cs137	6.1 Bq/Kg raw	± 2.9 Bq/Kg raw	6.1	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Kouya, Uchigo, Iwaki	Nov-11	Cs137	3.2 Bq/Kg raw	± 2.4 Bq/Kg raw	3.2	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Matsukusune, Joban, Iwaki	Oct-11	Cs137	3.4 Bq/Kg raw	± 2.5 Bq/Kg raw	3.4	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice (MilkyQueen)	Komoda, Kashima, Iwaki	Unknown	Cs137	3.2 Bq/Kg raw	± 2.7 Bq/Kg raw	6.4	Cs137	7.0 Bq/Kg raw	
			Cs134	3.2 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Rice	Kashima, Iwaki	Sep-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.4	Cs137	6.0 Bq/Kg raw	
			Cs134	2.4 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Tono, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.1	Cs137	6.0 Bq/Kg raw	
			Cs134	4.1 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Yamada, Iwaki	Oct-11	Cs137	2.9 Bq/Kg raw	± 2.8 Bq/Kg raw	5.9	Cs137	6.0 Bq/Kg raw	
			Cs134	3.0 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Watanabe, Iwaki	Aug-11	Cs137	3.8 Bq/Kg raw	± 2.4 Bq/Kg raw	3.8	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Onahamakanari, Iwaki	Oct-11	Cs137	4.7 Bq/Kg raw	± 2.9 Bq/Kg raw	9.3	Cs137	6.0 Bq/Kg raw	
			Cs134	4.6 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Onahamaohara, Iwaki	Oct-11	Cs137	4.6 Bq/Kg raw	± 2.7 Bq/Kg raw	7.5	Cs137	6.0 Bq/Kg raw	
			Cs134	2.9 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Rice	Ueda, Iwaki	Oct-11	Cs137	3.4 Bq/Kg raw	± 2.6 Bq/Kg raw	3.4	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Rice	Soeno, iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.7	Cs137	6.0 Bq/Kg raw	
			Cs134	2.7 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Glutinous rice	Takahagi, Ogawa, Iwaki	Oct-11	Cs137	3.9 Bq/Kg raw	± 2.6 Bq/Kg raw	6.3	Cs137	6.0 Bq/Kg raw	
			Cs134	2.4 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Glutinous rice	Shimoyada, Kashima, Iwaki	Oct-11	Cs137	3.8 Bq/Kg raw	± 3.0 Bq/Kg raw	9.9	Cs137	6.0 Bq/Kg raw	
			Cs134	6.1 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Shimotsuga, Gunma Pref.	Jun-11	Cs137	4.7 Bq/Kg raw	± 2.9 Bq/Kg raw	4.7	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	6.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Kamiogawa, Ogawa, Iwaki	Jul-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Maehara, Ogawa, Iwaki	Jun-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.7	Cs137	6.0 Bq/Kg raw	
			Cs134	3.7 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Akai, Taira, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Shimotakaku, Taira, Iwaki	Aug-11	Cs137	7.0 Bq/Kg raw	± 3.5 Bq/Kg raw	13.2	Cs137	6.0 Bq/Kg raw	
			Cs134	6.2 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Shimoyamaguchi, Taira, Iwaki	Jul-11	Cs137	3.9 Bq/Kg raw	± 2.5 Bq/Kg raw	7.8	Cs137	6.0 Bq/Kg raw	
			Cs134	3.9 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Nishiki, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.0	Cs137	6.0 Bq/Kg raw	
			Cs134	4.0 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Watanabe, Iwaki	Aug-11	Cs137	3.1 Bq/Kg raw	± 2.7 Bq/Kg raw	3.1	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Potato	Onahamasumiyoshi, Iwaki	Aug-11	Cs137	4.8 Bq/Kg raw	± 3.0 Bq/Kg raw	4.8	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Sweet potato	Nisiki, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	6.9	Cs137	6.0 Bq/Kg raw	
			Cs134	6.9 Bq/Kg raw	± 3.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Sweet potato	Tabiuto, Tabito, Iwaki	Nov-11	Cs137	8.1 Bq/Kg raw	± 4.2 Bq/Kg raw	13.7	Cs137	6.0 Bq/Kg raw	
			Cs134	5.6 Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Sweet potato	Ohisa, Iwaki	Nov-11	Cs137	8.0 Bq/Kg raw	± 0.4 Bq/Kg raw	11.6	Cs137	10.0 Bq/Kg raw	
			Cs134	3.6 Bq/Kg raw	± 0.6 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Sweet potato	Onahamaokaona, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.9	Cs137	6.0 Bq/Kg raw	
			Cs134	3.9 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Sweet potato	Hayama, Iwaki	Nov-11	Cs137	3.6 Bq/Kg raw	± 2.4 Bq/Kg raw	3.6	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Taro	Shimokabeya, Taira, Iwaki	Nov-11	Cs137	6.3 Bq/Kg raw	± 3.9 Bq/Kg raw	6.3	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Taro	Tabiuto, Tabito, Iwaki	Nov-11	Cs137	0.5 Bq/Kg raw	± 0.4 Bq/Kg raw	0.5	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Taro	Shimokawa, Izumi, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Yam	Ishikawa, Ishikawa	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	8.2	Cs137	6.0 Bq/Kg raw	
			Cs134	8.2 Bq/Kg raw	± 3.5 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Yam	Ishikawa, Ishikawa	Nov-11	Cs137	7.8 Bq/Kg raw	± 4.3 Bq/Kg raw	15.2	Cs137	6.0 Bq/Kg raw	
			Cs134	7.4 Bq/Kg raw	± 3.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Koriyama	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.7	Cs137	6.0 Bq/Kg raw	
			Cs134	3.7 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Shimoarakawa, Taira Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Shimoarakawa, Taira Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.3	Cs137	6.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Sanuka, Iwaki	Nov-11	Cs137	5.3 Bq/Kg raw	± 3.2 Bq/Kg raw	5.3	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Tairanakayama, Iwaki	Nov-11	Cs137	3.3 Bq/Kg raw	± 2.7 Bq/Kg raw	6.6	Cs137	6.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Furudono, Ishikawa	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Japanese white radish	Takakura, Iwaki	Nov-11	Cs137	5.0 Bq/Kg raw	± 3.5 Bq/Kg raw	10.1	Cs137	6.0 Bq/Kg raw	
			Cs134	5.1 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Japanese white radish	Kashima, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.2	Cs137	6.0 Bq/Kg raw	
			Cs134	4.2 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Carrot	Koriyama	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.8	Cs137	6.0 Bq/Kg raw	
			Cs134	2.8 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Pumpkin	Nihonmatsu	Oct-11	Cs137	6.4 Bq/Kg raw	± 6.4 Bq/Kg raw	10.4	Cs137	10.0 Bq/Kg raw	
			Cs134	4.0 Bq/Kg raw	± 15.5 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Onion	Kaminiida, Yotsukura, Iwaki	Aug-11	Cs137	3.5 Bq/Kg raw	± 2.5 Bq/Kg raw	3.5	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Onion	Nakayama, Taira, Iwaki	Aug-11	Cs137	3.5 Bq/Kg raw	± 2.6 Bq/Kg raw	6.9	Cs137	6.0 Bq/Kg raw	
			Cs134	3.4 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Onion	Shio, Taira, Iwaki	May-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.8	Cs137	6.0 Bq/Kg raw	
			Cs134	2.8 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Onion	Izumi, Iwaki	Jun-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.9	Cs137	6.0 Bq/Kg raw	
			Cs134	2.9 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Onion	Takahagi, Ogawa, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.3	Cs137	6.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Onion	Onahamasumiyoshi, Iwaki	Aug-11	Cs137	5.1 Bq/Kg raw	± 3.6 Bq/Kg raw	9.1	Cs137	6.0 Bq/Kg raw	
			Cs134	4.0 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Onion	Higashidai, Kaneyama, Iwaki	May-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.2	Cs137	6.0 Bq/Kg raw	
			Cs134	3.2 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Green onion	Onahamaokaona, Iwaki	Nov-11	Cs137	6.3 Bq/Kg raw	± 4.5 Bq/Kg raw	6.3	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Green onion	Kashima, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.5	Cs137	6.0 Bq/Kg raw	
			Cs134	3.5 Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Green onion	Shimoarakawa, Taira, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Green onion	Nishigo, Joban, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.7	Cs137	6.0 Bq/Kg raw	
			Cs134	2.7 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Green onion	Shimokawa, Izumi, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.1	Cs137	6.0 Bq/Kg raw	
			Cs134	4.1 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Green onion	Higashidai, Kanayama, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	10.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	10.0	Bq/Kg raw
Chinese cabbage	Furudono, Ishikawa	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	10.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	10.0	Bq/Kg raw
Chinese cabbage	Takakura, Iwaki	Nov-11	Cs137	6.3	Bq/Kg raw	± 4.0	Bq/Kg raw	6.3	Cs137	6.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Chinese cabbage	Ohisaitagizawa, Ohisa, Iwaki	Nov-11	Cs137	8.1	Bq/Kg raw	± 7.1	Bq/Kg raw	8.1	Cs137	6.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Chinese cabbage	Kamiyunagaya, Joban, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.1	Cs137	6.0	Bq/Kg raw
			Cs134	5.1	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Chinese cabbage	Nishigo, Joban, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.2	Cs137	6.0	Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Lettuce	Shimokawa, Izumi, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	6.2	Cs137	6.0	Bq/Kg raw
			Cs134	6.2	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Spinach	Minamishirado, Taira, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	6.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Leaf lettuce	Minamishirado, Taira, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137	6.0	Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Japanese mustard spinach	Sanuka, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.6	Cs137	6.0	Bq/Kg raw
			Cs134	4.6	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Soybean	Tabiuto, Tabito, Iwaki	Nov-11	Cs137	11.9	Bq/Kg raw	± 0.5	Bq/Kg raw	21.8	Cs137	10.0	Bq/Kg raw
			Cs134	9.9	Bq/Kg raw	± 0.7	Bq/Kg raw		Cs134	10.0	Bq/Kg raw
Shitake mushroom	Shiraiwa, Yotsukura, Iwaki	Nov-11	Cs137	304	Bq/Kg raw	± 61.0	Bq/Kg raw	544	Cs137	6.0	Bq/Kg raw
			Cs134	240	Bq/Kg raw	± 48.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Shitake mushroom	Shimokawa, Izumi, Iwaki	Nov-11	Cs137	358	Bq/Kg raw	± 72.0	Bq/Kg raw	667	Cs137	6.0	Bq/Kg raw
			Cs134	309	Bq/Kg raw	± 62.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Shitake mushroom	Negishi, Tono, Iwaki	Nov-11	Cs137	79.7	Bq/Kg raw	± 17.1	Bq/Kg raw	153	Cs137	6.0	Bq/Kg raw
			Cs134	73.1	Bq/Kg raw	± 14.6	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Shitake mushroom	Komagome, Yotsukura, Iwaki	Nov-11	Cs137	8.5	Bq/Kg raw	± 0.4	Bq/Kg raw	12.6	Cs137	10.0	Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 0.6	Bq/Kg raw		Cs134	10.0	Bq/Kg raw
Nameko mushroom	Onahamakamikajiro , Iwaki	Nov-11	Cs137	318	Bq/Kg raw	± 2.9	Bq/Kg raw	652	Cs137	10.0	Bq/Kg raw
			Cs134	334	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134	10.0	Bq/Kg raw
Nameko mushroom	Negishi, Tono, Iwaki	Nov-11	Cs137	85.2	Bq/Kg raw	± 18.5	Bq/Kg raw	161	Cs137	6.0	Bq/Kg raw
			Cs134	75.9	Bq/Kg raw	± 15.4	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Citron	Yotsukura, Iwaki	Nov-11	Cs137	64.9	Bq/Kg raw	± 14.4	Bq/Kg raw	119	Cs137	6.0	Bq/Kg raw
			Cs134	54.3	Bq/Kg raw	± 11.3	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Citron	Minamiodaira, Tabito, Iwaki	Nov-11	Cs137	34.4	Bq/Kg raw	± 8.3	Bq/Kg raw	71.7	Cs137	6.0	Bq/Kg raw
			Cs134	37.3	Bq/Kg raw	± 8.1	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Mandarin orange	Wakabadai, Iwaki	Nov-11	Cs137	44.6	Bq/Kg raw	± 10.3	Bq/Kg raw	81.4	Cs137	6.0	Bq/Kg raw
			Cs134	36.8	Bq/Kg raw	± 8.1	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Mandarin orange	Nakayama, Taira, Iwaki	Nov-11	Cs137	13.8	Bq/Kg raw	± 4.1	Bq/Kg raw	23.7	Cs137	6.0	Bq/Kg raw
			Cs134	9.9	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Mandarin orange	Shimoarakawa, Taira, Iwaki	Oct-11	Cs137	40.9	Bq/Kg raw	± 9.4	Bq/Kg raw	75.5	Cs137	6.0	Bq/Kg raw
			Cs134	34.6	Bq/Kg raw	± 7.5	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Mandarin orange	Onahamasuwa, Iwaki	Nov-11	Cs137	22.8	Bq/Kg raw	± 5.8	Bq/Kg raw	41.1	Cs137	6.0	Bq/Kg raw
			Cs134	18.3	Bq/Kg raw	± 4.5	Bq/Kg raw		Cs134	4.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Mandarin orange	Onahamanakachosakai, Iwaki	Nov-11	Cs137	9.3 Bq/Kg raw	± 3.6 Bq/Kg raw	22.5	Cs137	6.0 Bq/Kg raw	
			Cs134	13.2 Bq/Kg raw	± 3.5 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Mandarin orange	Ena, Iwaki	Nov-11	Cs137	31.7 Bq/Kg raw	± 7.7 Bq/Kg raw	63.4	Cs137	6.0 Bq/Kg raw	
			Cs134	31.7 Bq/Kg raw	± 7.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Mandarin orange	Tamatsuyu, Izumi, Iwaki	Nov-11	Cs137	9.7 Bq/Kg raw	± 3.4 Bq/Kg raw	17.2	Cs137	6.0 Bq/Kg raw	
			Cs134	7.5 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Mandarin orange	Onahamaminatogaoka, Iwaki	Nov-11	Cs137	18.3 Bq/Kg raw	± 5.2 Bq/Kg raw	35.5	Cs137	6.0 Bq/Kg raw	
			Cs134	17.2 Bq/Kg raw	± 4.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Lemon	Onahamaminatogaoka, Iwaki	Nov-11	Cs137	17.5 Bq/Kg raw	± 5.0 Bq/Kg raw	25.4	Cs137	6.0 Bq/Kg raw	
			Cs134	7.9 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Ohisa, Iwaki	Nov-11	Cs137	43.8 Bq/Kg raw	± 10.0 Bq/Kg raw	78.8	Cs137	6.0 Bq/Kg raw	
			Cs134	35.0 Bq/Kg raw	± 7.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Nishiogawa, Ogawa, Iwaki	Nov-11	Cs137	3.3 Bq/Kg raw	± 2.6 Bq/Kg raw	6.3	Cs137	6.0 Bq/Kg raw	
			Cs134	3.0 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	10.7 Bq/Kg raw	± 3.9 Bq/Kg raw	22.8	Cs137	6.0 Bq/Kg raw	
			Cs134	12.1 Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Minamiodaira, Tabito, Iwaki	Nov-11	Cs137	15.4 Bq/Kg raw	± 4.5 Bq/Kg raw	28.4	Cs137	6.0 Bq/Kg raw	
			Cs134	13.0 Bq/Kg raw	± 3.5 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Nakayama, Taira, Iwaki	Nov-11	Cs137	10.9 Bq/Kg raw	± 3.9 Bq/Kg raw	24.7	Cs137	6.0 Bq/Kg raw	
			Cs134	13.8 Bq/Kg raw	± 3.7 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Iino, Chuodai, Iwaki	Nov-11	Cs137	3.5 Bq/Kg raw	± 3.4 Bq/Kg raw	10.5	Cs137	6.0 Bq/Kg raw	
			Cs134	7.0 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Onahamanakachozakai, Iwaki	Nov-11	Cs137	3.8 Bq/Kg raw	± 2.6 Bq/Kg raw	6.9	Cs137	6.0 Bq/Kg raw	
			Cs134	3.1 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Persimmon	Onahamakamikajiro, Iwaki	Nov-11	Cs137	50.9 Bq/Kg raw	± 0.7 Bq/Kg raw	100	Cs137	10.0 Bq/Kg raw	
			Cs134	49.5 Bq/Kg raw	± 0.9 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Persimmon	Ena, Iwaki	Nov-11	Cs137	5.1 Bq/Kg raw	± 3.1 Bq/Kg raw	13.2	Cs137	6.0 Bq/Kg raw	
			Cs134	8.1 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Kiwi fruit	Yotsukura, Iwaki	Nov-11	Cs137	25.1 Bq/Kg raw	± 6.3 Bq/Kg raw	47.5	Cs137	6.0 Bq/Kg raw	
			Cs134	22.4 Bq/Kg raw	± 5.3 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Kiwi fruit	Nakayama, Taira, Iwaki	Nov-11	Cs137	24.7 Bq/Kg raw	± 6.3 Bq/Kg raw	45.2	Cs137	6.0 Bq/Kg raw	
			Cs134	20.5 Bq/Kg raw	± 4.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Kiwi fruit	Nakayama, Taira, Iwaki	Nov-11	Cs137	29.6 Bq/Kg raw	± 7.5 Bq/Kg raw	65.6	Cs137	6.0 Bq/Kg raw	
			Cs134	36.0 Bq/Kg raw	± 7.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Kiwi fruit	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	4.5 Bq/Kg raw	± 3.0 Bq/Kg raw	12.8	Cs137	6.0 Bq/Kg raw	
			Cs134	8.3 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Kiwi fruit	Onahamasuwa, Iwaki	Nov-11	Cs137	13.8 Bq/Kg raw	± 4.5 Bq/Kg raw	32.6	Cs137	6.0 Bq/Kg raw	
			Cs134	18.8 Bq/Kg raw	± 4.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Apple kiwi fruit	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	5.7 Bq/Kg raw	± 2.9 Bq/Kg raw	10.5	Cs137	6.0 Bq/Kg raw	
			Cs134	4.8 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Honey	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	14.8 Bq/Kg raw	± 4.2 Bq/Kg raw	20.7	Cs137	6.0 Bq/Kg raw	
			Cs134	5.9 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Honey	Shimookeuri, Kawamae, Iwaki	Oct-11	Cs137	34.4 Bq/Kg raw	± 8.2 Bq/Kg raw	65.3	Cs137	6.0 Bq/Kg raw	
			Cs134	30.9 Bq/Kg raw	± 6.9 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Honey	Hirakubo, Taira, Iwaki	Nov-11	Cs137	27.3 Bq/Kg raw	± 0.6 Bq/Kg raw	51.7	Cs137	10.0 Bq/Kg raw	
			Cs134	24.4 Bq/Kg raw	± 0.8 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Honey	Kawamae, Iwaki	Nov-11	Cs137	10.4 Bq/Kg raw	± 3.2 Bq/Kg raw	17.5	Cs137	6.0 Bq/Kg raw	
			Cs134	7.1 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Honey	Kawamae, Iwaki	Nov-11	Cs137	32.1 Bq/Kg raw	± 7.9 Bq/Kg raw	55.5	Cs137	6.0 Bq/Kg raw	
			Cs134	23.4 Bq/Kg raw	± 5.7 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Honey (Acacia)	Tomioka, Futaba	May-10	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Propolis	Brazil	Jan-01	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Salmon roe (Soy marinated)	Natsugawa, Taira, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Pickled vegetables	Taira, Iwaki	Nov-11	Cs137	2.7 Bq/Kg raw	± 2.6 Bq/Kg raw	6.9	Cs137	6.0 Bq/Kg raw	
			Cs134	4.2 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Miso	Taira, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
School lunch	Unknown	Nov-11	Cs137	4.8 Bq/Kg raw	± 0.4 Bq/Kg raw	8.3	Cs137	10.0 Bq/Kg raw	
			Cs134	3.5 Bq/Kg raw	± 0.6 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Spring water	Sashio, Miwa, Iwaki	Nov-11	Cs137	12.5 Bq/Kg raw	± 0.5 Bq/Kg raw	41.6	Cs137	10.0 Bq/Kg raw	
			Cs134	29.1 Bq/Kg raw	± 0.8 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Water	Tabiuto, Tabito, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Well water	Ohisa, Iwaki	Nov-11	Cs137	2.4 Bq/Kg raw	± 0.2 Bq/Kg raw	2.4	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Well water	Shimonagai, Miwa, Iwaki	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Well water	Taki, Tono, Iwaki	Nov-11	Cs137	0.7 Bq/Kg raw	± 0.2 Bq/Kg raw	0.7	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Well water	Koya, Uchigo, Iwaki	Nov-11	Cs137	17.8 Bq/Kg raw	± 0.3 Bq/Kg raw	41.5	Cs137	10.0 Bq/Kg raw	
			Cs134	23.7 Bq/Kg raw	± 0.4 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Well water	Unknown	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Drinking water (Tap water)	Miwa, Iwaki	Unknown	Cs137	2.4 Bq/Kg raw	± 8.3 Bq/Kg raw	3.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.6 Bq/Kg raw	± 49.5 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Mineral water	Unknown	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	10.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	10.0 Bq/Kg raw	
Dried rice straw	Oura, Yotsukura, Iwaki	Nov-11	Cs137	90.4 Bq/Kg raw	± 32.5 Bq/Kg raw	174	Cs137	6.0 Bq/Kg raw	
			Cs134	83.1 Bq/Kg raw	± 26.2 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Thatch plant	Uwadaira, Ogawa, Iwaki	Nov-11	Cs137	15.5 Bq/Kg raw	± 11.7 Bq/Kg raw	41.5	Cs137	6.0 Bq/Kg raw	
			Cs134	26.0 Bq/Kg raw	± 10.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	