



# Radiation Measurement Results of 314 items in December 2011



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

**Radioactive contamination level may differ according to sampling points even within the same address.**

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Brown rice (Koshihikari)	Wajima, Ishikawa	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice (Akitakomachi)	Yuzawa, Akita	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Mitsuke, Niigata	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Sekinami, Kitaibaraki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.1	Cs137	6.0 Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Namie, Futaba	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.6	Cs137	7.0 Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Kashima, Minamisoma	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.8	Cs137	7.0 Bq/Kg生
			Cs134	4.8	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Iwase, Sukagawa	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.4	Cs137	7.0 Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice (Hitomebore)	Kawanumagun, Aizubange	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.1	Cs137	6.0 Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice (Sasanishiki)	Aizuwakamatsu	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.6	Cs137	6.0 Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Yamagun, Nishiaizu	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Natsui, Ono, Tamura	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.3	Cs137	7.0 Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Yotsukura, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.5	Bq/Kg生	3.1	Cs137	6.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Yaguki, Yotsukura, Iwaki	Oct-11	Cs137	4.7	Bq/Kg生	± 2.7	Bq/Kg生	8.1	Cs137	7.0 Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Yaguki, Yotsukura, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.7	Bq/Kg生	7.5	Cs137	6.0 Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Yaguki, Yotsukura, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.7	Bq/Kg生	7.5	Cs137	6.0 Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	6.2	Bq/Kg生	± 3.2	Bq/Kg生	12.2	Cs137	6.0 Bq/Kg生
			Cs134	6.0	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Kabeya, Taira, Iwaki	Oct-11	Cs137	4.4	Bq/Kg生	± 2.7	Bq/Kg生	8.5	Cs137	7.0 Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Takaku, Taira, Iwaki	Oct-11	Cs137	4.1	Bq/Kg生	± 2.9	Bq/Kg生	8.2	Cs137	7.0 Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Brown rice	Yamadama, Iwaki	Oct-11	Cs137	5.7	Bq/Kg生	± 3.0	Bq/Kg生	12.1	Cs137	7.0 Bq/Kg生
			Cs134	6.4	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Miwa, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.7	Bq/Kg生	6.4	Cs137	7.0 Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Miwa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.5	Cs137	7.0 Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Shimoogawa, Ogawa, Iwaki	Sep-11	Cs137	4.2	Bq/Kg生	± 2.7	Bq/Kg生	6.8	Cs137	6.0 Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Ogawa, Iwaki	Oct-11	Cs137	4.2	Bq/Kg生	± 2.9	Bq/Kg生	7.2	Cs137	7.0 Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Tono, Iwaki	Oct-11	Cs137	2.6	Bq/Kg生	± 2.6	Bq/Kg生	2.6	Cs137	7.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.3	Cs137	6.0 Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Tono, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.1	Cs137	6.0 Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Yamada, Iwaki	Sep-11	Cs137	9.2	Bq/Kg生	± 3.4	Bq/Kg生	15.2	Cs137	6.0 Bq/Kg生
			Cs134	6.0	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Kawabe, Iwaki	Oct-11	Cs137	5.1	Bq/Kg生	± 3.3	Bq/Kg生	15.3	Cs137	6.0 Bq/Kg生
			Cs134	10.2	Bq/Kg生	± 3.2	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Koya, Uchigo, Iwaki	Sep-11	Cs137	7.1	Bq/Kg生	± 3.1	Bq/Kg生	10.6	Cs137	7.0 Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Kuroda, Tabito, Iwaki	Oct-11	Cs137	6.5	Bq/Kg生	± 3.1	Bq/Kg生	9.9	Cs137	7.0 Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Numabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.0	Cs137	6.0 Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	4.0 Bq/Kg生
Brown rice	Nakoso, Iwaki	Oct-11	Cs137	4.6	Bq/Kg生	± 3.0	Bq/Kg生	10.3	Cs137	7.0 Bq/Kg生
			Cs134	5.7	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Nishigo, Joban, Iwaki	Dec-11	Cs137	4.2	Bq/Kg生	± 2.8	Bq/Kg生	7.7	Cs137	7.0 Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.9	Bq/Kg生	9.6	Cs137	7.0 Bq/Kg生
			Cs134	6.3	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	5.0	Bq/Kg生	± 2.7	Bq/Kg生	8.4	Cs137	7.0 Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Fujiwara, Joban, Iwaki	Sep-11	Cs137	5.3	Bq/Kg生	± 2.9	Bq/Kg生	8.7	Cs137	7.0 Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 3.0	Bq/Kg生	10.8	Cs137	7.0 Bq/Kg生
			Cs134	7.1	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Kuramochi, kasima, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 2.8	Bq/Kg生	7.4	Cs137	7.0 Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Tamatsuyu, Izumi, Iwaki	Oct-11	Cs137	3.4	Bq/Kg生	± 2.8	Bq/Kg生	7.4	Cs137	7.0 Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Kamikamado, Watanabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.0	Cs137	7.0 Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Kamikamado, Watanabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0 Bq/Kg生
Brown rice	Bora, Watanabe, Iwaki	Oct-11	Cs137	15.0	Bq/Kg生	± 4.5	Bq/Kg生	23.8	Cs137	7.0 Bq/Kg生
			Cs134	8.8	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0 Bq/Kg生

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Rice	Hokkaido	2010	Cs137	2.5	Bq/Kg 生	± 2.4	Bq/Kg 生	2.5	Cs137	6.0 Bq/Kg 生
			Cs134	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Nagai, Yamagata	Sep-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	0.0	Cs137	7.0 Bq/Kg 生
			Cs134	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Hanakawa, Kitaiharaki, Ibaraki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	3.5	Cs137	7.0 Bq/Kg 生
			Cs134	3.5	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Sendai,Miyagi	Oct-11	Cs137	3.7	Bq/Kg 生	± 2.5	Bq/Kg 生	3.7	Cs137	6.0 Bq/Kg 生
			Cs134	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Nihonmatsu	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	2.3	Cs137	6.0 Bq/Kg 生
			Cs134	2.3	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Aizumisato, Onuma	Oct-11	Cs137	4.0	Bq/Kg 生	± 2.8	Bq/Kg 生	7.9	Cs137	7.0 Bq/Kg 生
			Cs134	3.9	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Natsui,Ono, Tamura	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	3.0	Cs137	7.0 Bq/Kg 生
			Cs134	3.0	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Ono, Tamura	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	3.2	Cs137	7.0 Bq/Kg 生
			Cs134	3.2	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Yaguki, Yotsukura,Iwaki	Nov-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	2.4	Cs137	6.0 Bq/Kg 生
			Cs134	2.4	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Yaguki, Yotsukura,Iwaki	Nov-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	2.4	Cs137	6.0 Bq/Kg 生
			Cs134	2.4	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Komagome, Yotsukura,Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	3.9	Cs137	6.0 Bq/Kg 生
			Cs134	3.9	Bq/Kg 生	± 2.0	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	4.5	Cs137	7.0 Bq/Kg 生
			Cs134	4.5	Bq/Kg 生	± 2.3	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Akai,Iwaki	Oct-10	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	3.9	Cs137	6.0 Bq/Kg 生
			Cs134	3.9	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Akai,Iwaki	Oct-10	Cs137	2.5	Bq/Kg 生	± 2.5	Bq/Kg 生	2.5	Cs137	6.0 Bq/Kg 生
			Cs134	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Kamiya,Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	4.2	Cs137	6.0 Bq/Kg 生
			Cs134	4.2	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Nakakamiya, Taira,Iwaki	Unknown	Cs137	4.4	Bq/Kg 生	± 2.6	Bq/Kg 生	8.4	Cs137	7.0 Bq/Kg 生
			Cs134	4.0	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Shimokamiya, Taira,Iwaki	Nov-11	Cs137	4.4	Bq/Kg 生	± 2.7	Bq/Kg 生	6.5	Cs137	6.0 Bq/Kg 生
			Cs134	2.1	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Shimokatayose, Taira,Iwaki	Oct-11	Cs137	4.7	Bq/Kg 生	± 2.7	Bq/Kg 生	7.2	Cs137	6.0 Bq/Kg 生
			Cs134	2.5	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Takaku,Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	2.8	Cs137	6.0 Bq/Kg 生
			Cs134	2.8	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Takaku,Taira, Iwaki	Oct-11	Cs137	3.4	Bq/Kg 生	± 2.7	Bq/Kg 生	6.7	Cs137	7.0 Bq/Kg 生
			Cs134	3.3	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Shimotakaku, Taira,Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	3.0	Cs137	7.0 Bq/Kg 生
			Cs134	3.0	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Nakashio, Taira,Iwaki	Oct-10	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	2.3	Cs137	6.0 Bq/Kg 生
			Cs134	2.3	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	4.0 Bq/Kg 生
Rice	Kamihirakubo, Taira,Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	2.8	Cs137	7.0 Bq/Kg 生
			Cs134	2.8	Bq/Kg 生	± 2.2	Bq/Kg 生		Cs134	6.0 Bq/Kg 生
Rice	Oyasaku, Yoshima,Iwaki	Oct-11	Cs137	0.0	Bq/Kg 生	± 0.0	Bq/Kg 生	4.2	Cs137	6.0 Bq/Kg 生
			Cs134	4.2	Bq/Kg 生	± 2.1	Bq/Kg 生		Cs134	4.0 Bq/Kg 生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Rice	Nishiogawa, Ogawa, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.5	Cs137 7.0 Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.2	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Nemoto, Ogawa, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 2.7	Bq/Kg生	3.8	Cs137 7.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Tabito, Iwaki	Oct-11	Cs137	4.4	Bq/Kg生	± 2.7	Bq/Kg生	7.3	Cs137 7.0 Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.2	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Kuroda, Tabito, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.5	Cs137 7.0 Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 6.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Kadono, Tono, Iwaki	Oct-11	Cs137	5.1	Bq/Kg生	± 3.5	Bq/Kg生	10.7	Cs137 7.0 Bq/Kg生
			Cs134	5.6	Bq/Kg生	± 2.8	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Kadono, Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.0	Cs137 7.0 Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Kashima, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.9	Cs137 7.0 Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.9	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Kamiyada, Joban, Iwaki	Unknown	Cs137	3.9	Bq/Kg生	± 2.7	Bq/Kg生	7.3	Cs137 6.0 Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Yamada, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.5	Cs137 6.0 Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Yamada, Iwaki	Oct-11	Cs137	3.2	Bq/Kg生	± 2.5	Bq/Kg生	3.2	Cs137 6.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Matsunoshita, Kawabe, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.7	Cs137 7.0 Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Ohara, Onahama, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.0	Cs137 6.0 Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Honya, Izumi, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.9	Cs137 7.0 Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Soeno, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.9	Cs137 6.0 Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Nakoso, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.6	Bq/Kg生	7.5	Cs137 7.0 Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Nakoso, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 2.7	Bq/Kg生	7.7	Cs137 7.0 Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.2	Bq/Kg生		Cs134 6.0 Bq/Kg生
Rice	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.5	Cs137 6.0 Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 4.0 Bq/Kg生
Rice	Unknown	Dec-11	Cs137	3.3	Bq/Kg生	± 2.7	Bq/Kg生	6.4	Cs137 7.0 Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134 6.0 Bq/Kg生
Glutinous rice (Brown rice)	Tabito, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 2.9	Bq/Kg生	8.8	Cs137 7.0 Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134 6.0 Bq/Kg生
Glutinous rice (Brown rice)	Nakoso, Iwaki	Sep-11	Cs137	11.9	Bq/Kg生	± 4.0	Bq/Kg生	22.1	Cs137 6.0 Bq/Kg生
			Cs134	10.2	Bq/Kg生	± 3.2	Bq/Kg生		Cs134 4.0 Bq/Kg生
Glutinous rice (Brown rice)	Izumida, Watanabe, Iwaki	Oct-11	Cs137	6.3	Bq/Kg生	± 3.1	Bq/Kg生	9.0	Cs137 6.0 Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.3	Bq/Kg生		Cs134 4.0 Bq/Kg生
Glutinous rice	Nanyo, Yamagata	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.5	Cs137 7.0 Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134 6.0 Bq/Kg生
Glutinous rice	Onami, Fukushima	Nov-11	Cs137	10.0	Bq/Kg生	± 3.7	Bq/Kg生	17.8	Cs137 7.0 Bq/Kg生
			Cs134	7.8	Bq/Kg生	± 2.9	Bq/Kg生		Cs134 6.0 Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Glutinous rice	Takakura, Iwaki	Oct-11	Cs137	5.4 Bq/Kg生	± 2.9 Bq/Kg生	11.9	Cs137	6.0 Bq/Kg生	
			Cs134	6.5 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Glutinous rice	Kagono, Tono, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.5	Cs137	7.0 Bq/Kg生	
			Cs134	3.5 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Glutinous rice	Tono, Iwaki	Oct-11	Cs137	4.8 Bq/Kg生	± 2.7 Bq/Kg生	8.8	Cs137	6.0 Bq/Kg生	
			Cs134	4.0 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Glutinous rice	Onahama, Iwaki	Oct-11	Cs137	6.8 Bq/Kg生	± 0.4 Bq/Kg生	11.1	Cs137	8.0 Bq/Kg生	
			Cs134	4.3 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Glutinous rice	Ogawa, Iwaki	Oct-11	Cs137	0.7 Bq/Kg生	± 0.3 Bq/Kg生	1.0	Cs137	8.0 Bq/Kg生	
			Cs134	0.3 Bq/Kg生	± 0.5 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Potato	Godo, Miwa, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Potato	Shimokabeya, Taira, Iwaki	Jul-11	Cs137	1.5 Bq/Kg生	± 0.4 Bq/Kg生	7.3	Cs137	8.0 Bq/Kg生	
			Cs134	5.8 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Potato	Shimokabeya, Taira, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.1	Cs137	6.0 Bq/Kg生	
			Cs134	3.1 Bq/Kg生	± 3.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Potato	Shimotakaku, Taira, Iwaki	Jun-11	Cs137	3.5 Bq/Kg生	± 0.4 Bq/Kg生	4.4	Cs137	8.0 Bq/Kg生	
			Cs134	0.9 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Potato	Shimohirakubo, Taira, Iwaki	Unknown	Cs137	1.9 Bq/Kg生	± 19.8 Bq/Kg生	1.9	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Potato	Nakayama, Taira, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.6	Cs137	6.0 Bq/Kg生	
			Cs134	3.6 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Sweet potato	Midaisakai, Uchigo, Iwaki	Dec-11	Cs137	2.8 Bq/Kg生	± 0.4 Bq/Kg生	3.1	Cs137	10.0 Bq/Kg生	
			Cs134	0.3 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Potato	Shimohata, Kashima, Iwak	Aug-11	Cs137	4.9 Bq/Kg生	± 0.4 Bq/Kg生	6.2	Cs137	8.0 Bq/Kg生	
			Cs134	1.3 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Potato	Tono, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.9	Cs137	6.0 Bq/Kg生	
			Cs134	2.9 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Potato	Honya, Izumi, Iwaki	Jul-11	Cs137	4.1 Bq/Kg生	± 3.3 Bq/Kg生	4.1	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Potato	Kaidomari, Tabito, Iwak	Sep-11	Cs137	5.9 Bq/Kg生	± 3.2 Bq/Kg生	5.9	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Potato	Nishigo, Joban, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.5	Cs137	7.0 Bq/Kg生	
			Cs134	3.5 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Potato	Nagasaki, Iwaki	Jul-11	Cs137	1.2 Bq/Kg生	± 0.4 Bq/Kg生	4.0	Cs137	10.0 Bq/Kg生	
			Cs134	2.8 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Potato	Onahama, Iwaki	Jun-11	Cs137	1.4 Bq/Kg生	± 0.4 Bq/Kg生	2.7	Cs137	8.0 Bq/Kg生	
			Cs134	1.3 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Potato	Onahama, Iwaki	Jun-11	Cs137	4.8 Bq/Kg生	± 0.4 Bq/Kg生	10.1	Cs137	8.0 Bq/Kg生	
			Cs134	5.3 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Potato	Ohara, Onahama, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	5.3	Cs137	7.0 Bq/Kg生	
			Cs134	5.3 Bq/Kg生	± 2.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Taro	Kurobe, Toyama Pref.	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	5.1	Cs137	7.0 Bq/Kg生	
			Cs134	5.1 Bq/Kg生	± 3.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Taro	Hosoya, Yotsukura, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Taro	Shimoyamaguchi , Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Taro	Minamiotaira, Tabito, Iwaki	Nov-11	Cs137	4.6 Bq/Kg生	± 2.6 Bq/Kg生	4.6	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Taro	Kamiyunagaya, Joban, Iwaki	Nov-11	Cs137	0.8 Bq/Kg生	± 0.3 Bq/Kg生	0.8	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Taro	Shimoyamaguchi , Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Taro	Tamatsuyu, Izumi, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Taro	Tamatsuyu, Izumi, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Taro	Okaona, Onahama, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Yam	Kamikajiro, Onahama, Iwaki	Dec-11	Cs137	13.0 Bq/Kg生	± 3.6 Bq/Kg生	25.0	Cs137	8.0 Bq/Kg生	
			Cs134	12.0 Bq/Kg生	± 5.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Yam	Kamikajiro, Onahama, Iwaki	Dec-11	Cs137	3.2 Bq/Kg生	± 0.4 Bq/Kg生	5.0	Cs137	8.0 Bq/Kg生	
			Cs134	1.8 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Yam	Kuroda, Tabito, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	5.8	Cs137	7.0 Bq/Kg生	
			Cs134	5.8 Bq/Kg生	± 3.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Yam	Takakura, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.7	Cs137	6.0 Bq/Kg生	
			Cs134	3.7 Bq/Kg生	± 3.3 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Yam	Tamatsuyu, Izumi, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生	
			Cs134	3.7 Bq/Kg生	± 2.7 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Yam	Oyasaku, Yoshima, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Japanese white radish (with leaf)	Nakakabeya, Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.6	Cs137	7.0 Bq/Kg生	
			Cs134	2.6 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish	Ohisa, Ohisa, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.7	Cs137	6.0 Bq/Kg生	
			Cs134	2.7 Bq/Kg生	± 2.7 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Japanese white radish	Shimoyoshima, Yoshima, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Japanese white radish	Kitayoshima, Yoshima, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish	Kamihirakubo, Taira, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.3	Cs137	6.0 Bq/Kg生	
			Cs134	2.3 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Japanese white radish	Akai, Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生	
			Cs134	3.1 Bq/Kg生	± 2.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish	Shimotakaku, Taira, Iwaki	Dec-11	Cs137	0.2 Bq/Kg生	± 0.4 Bq/Kg生	0.2	Cs137	8.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Japanese white radish	Shimoyamaguchi , Taira, Iwaki	Dec-11	Cs137	5.6 Bq/Kg生	± 3.4 Bq/Kg生	10.2	Cs137	6.0 Bq/Kg生	
			Cs134	4.6 Bq/Kg生	± 2.8 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Japanese white radish	Shimoyamaguchi , Taira, Iwaki	Dec-11	Cs137	5.6 Bq/Kg生	± 3.4 Bq/Kg生	10.2	Cs137	6.0 Bq/Kg生	
			Cs134	4.6 Bq/Kg生	± 2.8 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Japanese white radish	Shibahara, Ogawa, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.7	Cs137	7.0 Bq/Kg生	
			Cs134	2.7 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish	Fujiwara, Joban, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish	Nishigo, Joban, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Japanese white radish	Tsuzura, Uchigo, Iwaki	Dec-11	Cs137	4.3	Bq/Kg生	± 3.3 Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shiramizu, Uchigo, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	10.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0	Bq/Kg生
Japanese white radish	Shimokawa, Izumi, Iwaki	Dec-11	Cs137	2.7	Bq/Kg生	± 2.4 Bq/Kg生	4.9	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 2.1 Bq/Kg生		Cs134	4.0	Bq/Kg生
Japanese white radish	Kamikajiro, Onahama, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0	Bq/Kg生
Japanese white radish	Kaidomari, Tabito, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	3.4	Cs137	6.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.9 Bq/Kg生		Cs134	4.0	Bq/Kg生
Japanese white radish	Baba, Nishiki, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	10.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0	Bq/Kg生
Japanese white radish	Watanabe, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	3.6	Cs137	6.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 3.4 Bq/Kg生		Cs134	4.0	Bq/Kg生
Round Japanese white radish	Nakayama, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	2.4	Cs137	6.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.1 Bq/Kg生		Cs134	4.0	Bq/Kg生
Japanese white radish(leaf)	Shiramizu, Uchigo, Iwaki	Dec-11	Cs137	8.9	Bq/Kg生	± 8.9 Bq/Kg生	22.3	Cs137	6.0	Bq/Kg生
			Cs134	13.4	Bq/Kg生	± 7.3 Bq/Kg生		Cs134	4.0	Bq/Kg生
Carrot	Akai, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	3.5	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 3.1 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Ohisa, Ohisa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	4.7	Cs137	6.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 3.3 Bq/Kg生		Cs134	4.0	Bq/Kg生
Chinese cabbage	Yashikimae, Yotsukura, Iwaki	Dec-11	Cs137	38.6	Bq/Kg生	± 9.0 Bq/Kg生	72.2	Cs137	7.0	Bq/Kg生
			Cs134	33.6	Bq/Kg生	± 7.4 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Tamayama, Yotsukura, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	5.9	Bq/Kg生	± 3.8 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kitayoshima, Yoshima, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 3.6 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kamiya, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	7.3	Bq/Kg生	± 4.5 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kamihirakubo, Taira, Iwaki	Dec-11	Cs137	1.4	Bq/Kg生	± 0.4 Bq/Kg生	1.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Akai, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	4.9	Cs137	7.0	Bq/Kg生
			Cs134	4.9	Bq/Kg生	± 3.2 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimoarakawa, Taira, Iwaki	Dec-11	Cs137	8.5	Bq/Kg生	± 6.6 Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimoyamaguchi, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0	Bq/Kg生
Chinese cabbage	Shiramizu, Uchigo, Iwaki	Dec-11	Cs137	7.1	Bq/Kg生	± 5.8 Bq/Kg生	15.1	Cs137	6.0	Bq/Kg生
			Cs134	8.0	Bq/Kg生	± 4.7 Bq/Kg生		Cs134	4.0	Bq/Kg生
Chinese cabbage	Tsuzura, Uchigo, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	5.5	Cs137	7.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 3.9 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Nishigo, Joban, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 3.2 Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Nishigo, Joban, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0	Bq/Kg生
Chinese cabbage	Mizunoya, Joban, Iwaki	Dec-11	Cs137	5.7	Bq/Kg生	± 5.3 Bq/Kg生	12.4	Cs137	6.0	Bq/Kg生
			Cs134	6.7	Bq/Kg生	± 4.3 Bq/Kg生		Cs134	4.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Chinese cabbage	Nigishi, Tono, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Chinese cabbage	Miyamada, Tono, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Chinese cabbage	Tamatsuyu, Izumi, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.6
			Cs134	6.6	Bq/Kg生	± 4.5	Bq/Kg生	
Chinese cabbage	Misawa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.4
			Cs134	4.4	Bq/Kg生	± 4.3	Bq/Kg生	
Chinese cabbage	Minamiotaira, Tabito, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.1
			Cs134	4.1	Bq/Kg生	± 3.2	Bq/Kg生	
Chinese cabbage	Nagasaki, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Chinese cabbage	Kamikajiro, Onahama, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Chinese cabbage	Soeno, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.2
			Cs134	6.2	Bq/Kg生	± 4.0	Bq/Kg生	
Chinese cabbage	Watanabe, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.5
			Cs134	6.5	Bq/Kg生	± 5.8	Bq/Kg生	
Green onion	Kurobe, Toyama Pref.	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.7
			Cs134	5.7	Bq/Kg生	± 3.0	Bq/Kg生	
Green onion	Hosoya, Yotsukura, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3
			Cs134	3.3	Bq/Kg生	± 2.8	Bq/Kg生	
Green onion	Tamayama, Yotsukura, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Green onion	Shimoyamaguchi, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.9
			Cs134	4.9	Bq/Kg生	± 3.1	Bq/Kg生	
Green onion	Akai, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.7
			Cs134	3.7	Bq/Kg生	± 3.3	Bq/Kg生	
Green onion	Shimohirakubo, Taira, Iwaki	Dec-11	Cs137	0.8	Bq/Kg生	± 0.4	Bq/Kg生	0.8
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Green onion	Wakabadai, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Green onion	Mizunoya, Joban, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.6
			Cs134	5.6	Bq/Kg生	± 5.3	Bq/Kg生	
Green onion	Tamatsuyu, Izumi, Iwaki	Dec-11	Cs137	4.4	Bq/Kg生	± 4.5	Bq/Kg生	4.4
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Green onion	Minamiotaira, Tabito, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Green onion	Kamikajiro, Onahama, Iwaki	Dec-11	Cs137	4.1	Bq/Kg生	± 0.4	Bq/Kg生	4.1
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Green onion	Teramawari, Onahama, Iwaki	Dec-11	Cs137	4.7	Bq/Kg生	± 4.2	Bq/Kg生	8.3
			Cs134	3.6	Bq/Kg生	± 3.5	Bq/Kg生	
Green onion	Teramawari, Onahama, Iwaki	Dec-11	Cs137	6.1	Bq/Kg生	± 4.2	Bq/Kg生	6.1
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	
Onion	Izumizaki, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.2
			Cs134	6.2	Bq/Kg生	± 3.1	Bq/Kg生	
Onion	Shimohirakubo, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生	

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Onion	Ohara, Onahama, Iwaki	Jul-11	Cs137	8.8 Bq/Kg生	± 4.9 Bq/Kg生	8.8	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Onion	Ohara, Onahama, Iwaki	Sep-11	Cs137	4.8 Bq/Kg生	± 3.0 Bq/Kg生	7.6	Cs137	6.0 Bq/Kg生	
			Cs134	2.8 Bq/Kg生	± 2.5 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Lettuce	Takano, Uchigo, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	8.1	Cs137	7.0 Bq/Kg生	
			Cs134	8.1 Bq/Kg生	± 4.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Cabbage	Chibe Pref.	Dec-11	Cs137	7.7 Bq/Kg生	± 5.8 Bq/Kg生	17.2	Cs137	7.0 Bq/Kg生	
			Cs134	9.5 Bq/Kg生	± 4.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Cabbage	Akai, Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	6.6	Cs137	7.0 Bq/Kg生	
			Cs134	6.6 Bq/Kg生	± 4.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Cabbage	Akai, Taira, Iwaki	Dec-11	Cs137	7.8 Bq/Kg生	± 6.7 Bq/Kg生	16.9	Cs137	7.0 Bq/Kg生	
			Cs134	9.1 Bq/Kg生	± 5.5 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Cabbage	Fujiwara, Joban, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	5.7	Cs137	7.0 Bq/Kg生	
			Cs134	5.7 Bq/Kg生	± 3.7 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Spinach	Nishigo, Joban, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	8.7	Cs137	7.0 Bq/Kg生	
			Cs134	8.7 Bq/Kg生	± 5.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese mustard spinach	Taihaku, Sendai, Miyagi Pref.	Nov-11	Cs137	0.5 Bq/Kg生	± 71.4 Bq/Kg生	0.5	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Japanese mustard spinach	Akai, Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	6.1	Cs137	7.0 Bq/Kg生	
			Cs134	6.1 Bq/Kg生	± 3.5 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese mustard spinach	Shimoyoshima, Yoshima, Iwaki	Dec-11	Cs137	11.2 Bq/Kg生	± 7.1 Bq/Kg生	19.0	Cs137	7.0 Bq/Kg生	
			Cs134	7.8 Bq/Kg生	± 5.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Potherb mustard	Nishigo, Joban, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	6.5	Cs137	7.0 Bq/Kg生	
			Cs134	6.5 Bq/Kg生	± 4.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Yukina green	Taihaku, Sendai, Miyagi Pref.	Nov-11	Cs137	0.1 Bq/Kg生	± 270 Bq/Kg生	0.1	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Garland chrysanthemum	Okaona, Onahama, Iwaki	Dec-11	Cs137	7.2 Bq/Kg生	± 6.6 Bq/Kg生	7.2	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Garland chrysanthemum	Shiramizu, Uchigo, Iwaki	Dec-11	Cs137	5.3 Bq/Kg生	± 0.4 Bq/Kg生	11.2	Cs137	10.0 Bq/Kg生	
			Cs134	5.9 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Herana green	Yashikimae, Yotsukura, Iwaki	Dec-11	Cs137	12.8 Bq/Kg生	± 4.1 Bq/Kg生	20.5	Cs137	7.0 Bq/Kg生	
			Cs134	7.7 Bq/Kg生	± 2.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Petit Vert	Kashima, Chuodai, Iwaki	Dec-11	Cs137	36.4 Bq/Kg生	± 0.5 Bq/Kg生	72.9	Cs137	10.0 Bq/Kg生	
			Cs134	36.5 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Broccoli	Kamiya, Taira, Iwaki	Dec-11	Cs137	8.4 Bq/Kg生	± 5.7 Bq/Kg生	17.2	Cs137	7.0 Bq/Kg生	
			Cs134	8.8 Bq/Kg生	± 4.6 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Broccoli	Nishigo, Joban, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Broccoli	Baba, Nishiki, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Yacon(dried)	Miyamada, Tonol, Iwaki	Oct-11	Cs137	175 Bq/Kg生	± 1.4 Bq/Kg生	365	Cs137	10.0 Bq/Kg生	
			Cs134	190 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Shiitake mushroom	Yumototenjin, Joban, Iwaki	Dec-11	Cs137	253 Bq/Kg生	± 2.0 Bq/Kg生	477	Cs137	10.0 Bq/Kg生	
			Cs134	224 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Shiitake mushroom	Taki, Tono, Iwaki	Nov-11	Cs137	144 Bq/Kg生	± 1.2 Bq/Kg生	275	Cs137	8.0 Bq/Kg生	
			Cs134	131 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Shiitake mushroom	Minamiodaira, Tabito, Iwaki	Dec-11	Cs137	121 Bq/Kg生	± 1.0 Bq/Kg生	234	Cs137	10.0 Bq/Kg生	
			Cs134	113 Bq/Kg生	± 1.1 Bq/Kg生		Cs134	10.0 Bq/Kg生	

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Shiitake mushroom	Kamikajiro, Onahama, Iwaki	Dec-11	Cs137	88.0 Bq/Kg生	± 0.6 Bq/Kg生	157	Cs137	10.0 Bq/Kg生
			Cs134	68.8 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	10.0 Bq/Kg生
Shiitake mushroom	Tamatsuyu, Izumi, Iwaki	Nov-11	Cs137	155 Bq/Kg生	± 1.2 Bq/Kg生	305	Cs137	8.0 Bq/Kg生
			Cs134	150 Bq/Kg生	± 1.5 Bq/Kg生		Cs134	8.0 Bq/Kg生
Dried shiitake mushroom	Shimohunao, Joban, Iwaki	Nov-11	Cs137	112 Bq/Kg生	± 29.0 Bq/Kg生	211	Cs137	6.0 Bq/Kg生
			Cs134	98.5 Bq/Kg生	± 24.0 Bq/Kg生		Cs134	4.0 Bq/Kg生
Dried shiitake mushroom	Kawauchi, Futaba	Nov-11	Cs137	1557 Bq/Kg生	± 10.9 Bq/Kg生	2,872	Cs137	8.0 Bq/Kg生
			Cs134	1315 Bq/Kg生	± 13.2 Bq/Kg生		Cs134	8.0 Bq/Kg生
Nameko mushroom	Umegaoka, Yotsukura, Iwaki	Nov-11	Cs137	548 Bq/Kg生	± 4.9 Bq/Kg生	1,091	Cs137	8.0 Bq/Kg生
			Cs134	543 Bq/Kg生	± 5.4 Bq/Kg生		Cs134	8.0 Bq/Kg生
Matsutake mushroom	Tamayama, Yotsukura, Iwaki	Unknown	Cs137	55.9 Bq/Kg生	± 0.6 Bq/Kg生	96.4	Cs137	8.0 Bq/Kg生
			Cs134	40.5 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	8.0 Bq/Kg生
Citron (with peel, seed)	Soeno, Iwaki	Dec-11	Cs137	74.7 Bq/Kg生	± 17.2 Bq/Kg生	141	Cs137	7.0 Bq/Kg生
			Cs134	66.0 Bq/Kg生	± 14.3 Bq/Kg生		Cs134	6.0 Bq/Kg生
Citron	Sekinami, Kitaibaraki	Nov-11	Cs137	58.2 Bq/Kg生	± 0.8 Bq/Kg生	120	Cs137	10.0 Bq/Kg生
			Cs134	61.4 Bq/Kg生	± 1.0 Bq/Kg生		Cs134	10.0 Bq/Kg生
Citron	Misawa, Iwaki	Dec-11	Cs137	40.0 Bq/Kg生	± 9.4 Bq/Kg生	72.3	Cs137	6.0 Bq/Kg生
			Cs134	32.3 Bq/Kg生	± 7.3 Bq/Kg生		Cs134	4.0 Bq/Kg生
Citron	Shimoyoshima, Yoshima, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0 Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0 Bq/Kg生
Citron	Taira, Iwaki	Dec-11	Cs137	72.5 Bq/Kg生	± 0.9 Bq/Kg生	136	Cs137	10.0 Bq/Kg生
			Cs134	63.7 Bq/Kg生	± 1.1 Bq/Kg生		Cs134	10.0 Bq/Kg生
Citron	Shimoarakawa, Taira, Iwaki	Dec-11	Cs137	67.8 Bq/Kg生	± 0.8 Bq/Kg生	127	Cs137	10.0 Bq/Kg生
			Cs134	58.8 Bq/Kg生	± 0.9 Bq/Kg生		Cs134	10.0 Bq/Kg生
Citron	Wakabadai, Taira, Iwaki	Dec-11	Cs137	64.0 Bq/Kg生	± 0.9 Bq/Kg生	124	Cs137	8.0 Bq/Kg生
			Cs134	60.4 Bq/Kg生	± 1.0 Bq/Kg生		Cs134	8.0 Bq/Kg生
Citron	Midaisakai, Uchigo, Iwaki	Dec-11	Cs137	45.6 Bq/Kg生	± 0.7 Bq/Kg生	87.4	Cs137	10.0 Bq/Kg生
			Cs134	41.8 Bq/Kg生	± 0.9 Bq/Kg生		Cs134	10.0 Bq/Kg生
Citron	Kubota, Nakoso, Iwaki	Dec-11	Cs137	19.8 Bq/Kg生	± 0.5 Bq/Kg生	33.7	Cs137	10.0 Bq/Kg生
			Cs134	13.9 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	10.0 Bq/Kg生
Citron	Odaira, Tono, Iwaki	Dec-11	Cs137	42.9 Bq/Kg生	± 1.6 Bq/Kg生	80.3	Cs137	10.0 Bq/Kg生
			Cs134	37.4 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	10.0 Bq/Kg生
Citron	Kamiyunagawa, Joban, Iwaki	Dec-11	Cs137	43.3 Bq/Kg生	± 10.3 Bq/Kg生	77.0	Cs137	7.0 Bq/Kg生
			Cs134	33.7 Bq/Kg生	± 7.8 Bq/Kg生		Cs134	6.0 Bq/Kg生
Citron	Yokodai, Iwaki	Dec-11	Cs137	212 Bq/Kg生	± 1.7 Bq/Kg生	403	Cs137	8.0 Bq/Kg生
			Cs134	191 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	8.0 Bq/Kg生
Citron	Tamagawa, Onahama, Iwaki	Dec-11	Cs137	94.4 Bq/Kg生	± 0.9 Bq/Kg生	177	Cs137	8.0 Bq/Kg生
			Cs134	82.8 Bq/Kg生	± 1.1 Bq/Kg生		Cs134	8.0 Bq/Kg生
Citron	Okaona, Onahama, Iwaki	Dec-11	Cs137	82.4 Bq/Kg生	± 0.9 Bq/Kg生	152	Cs137	10.0 Bq/Kg生
			Cs134	69.9 Bq/Kg生	± 1.0 Bq/Kg生		Cs134	10.0 Bq/Kg生
Mandarin orange	Shimokabeya, Taira, Iwaki	Dec-11	Cs137	47.1 Bq/Kg生	± 0.7 Bq/Kg生	89.5	Cs137	8.0 Bq/Kg生
			Cs134	42.4 Bq/Kg生	± 0.9 Bq/Kg生		Cs134	8.0 Bq/Kg生
Mandarin orange	Takaku, Chuodai, Iwaki	Dec-11	Cs137	33.9 Bq/Kg生	± 8.1 Bq/Kg生	65.5	Cs137	6.0 Bq/Kg生
			Cs134	31.6 Bq/Kg生	± 7.0 Bq/Kg生		Cs134	4.0 Bq/Kg生
Mandarin orange	Kashima, Chuodai, Iwaki	Dec-11	Cs137	127 Bq/Kg生	± 1.1 Bq/Kg生	248	Cs137	10.0 Bq/Kg生
			Cs134	121 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	10.0 Bq/Kg生
Mandarin orange	Noda, Onahama, Iwaki	Dec-11	Cs137	61.4 Bq/Kg生	± 0.8 Bq/Kg生	117	Cs137	8.0 Bq/Kg生
			Cs134	55.2 Bq/Kg生	± 0.9 Bq/Kg生		Cs134	8.0 Bq/Kg生

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Mandarin orange	Sumiyoshi, Onahama, Iwaki	Dec-11	Cs137	25.9 Bq/Kg生	± 0.6 Bq/Kg生	53.9	Cs137	8.0 Bq/Kg生	
			Cs134	28.0 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Mandarin orange	Kubota, Nakoso, Iwaki	Dec-11	Cs137	24.9 Bq/Kg生	± 0.6 Bq/Kg生	44.2	Cs137	10.0 Bq/Kg生	
			Cs134	19.3 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Mandarin orange	Izumigaoka, Iwaki	Nov-11	Cs137	14.8 Bq/Kg生	± 4.4 Bq/Kg生	30.2	Cs137	7.0 Bq/Kg生	
			Cs134	15.4 Bq/Kg生	± 3.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Lemon	Izumigaoka, Iwaki	Dec-11	Cs137	18.0 Bq/Kg生	± 5.2 Bq/Kg生	37.8	Cs137	7.0 Bq/Kg生	
			Cs134	19.8 Bq/Kg生	± 4.8 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese citron	Shimoarakawa, Taira, Iwaki	Dec-11	Cs137	55.7 Bq/Kg生	± 12.9 Bq/Kg生	99.0	Cs137	6.0 Bq/Kg生	
			Cs134	43.3 Bq/Kg生	± 9.7 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Kiwi fruit	Kawauchi	Dec-11	Cs137	12.0 Bq/Kg生	± 0.5 Bq/Kg生	22.6	Cs137	8.0 Bq/Kg生	
			Cs134	10.6 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Kiwi fruit	Sakuragaoka, Iwaki	Nov-11	Cs137	33.2 Bq/Kg生	± 0.6 Bq/Kg生	62.6	Cs137	8.0 Bq/Kg生	
			Cs134	29.4 Bq/Kg生	± 0.9 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Kiwi fruit	Kamiyunagaya, Joban, Iwaki	Nov-11	Cs137	36.4 Bq/Kg生	± 1.8 Bq/Kg生	68.9	Cs137	10.0 Bq/Kg生	
			Cs134	32.5 Bq/Kg生	± 2.5 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Kiwi fruit	Kamiyunagaya, Joban, Iwaki	Nov-11	Cs137	36.4 Bq/Kg生	± 0.7 Bq/Kg生	68.9	Cs137	10.0 Bq/Kg生	
			Cs134	32.5 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Kiwi fruit	Shimohata, Kashima, Iwaki	Nov-11	Cs137	25.5 Bq/Kg生	± 0.6 Bq/Kg生	48.7	Cs137	8.0 Bq/Kg生	
			Cs134	23.2 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Kiwi fruit	Onahama, Iwaki	Dec-11	Cs137	14.0 Bq/Kg生	± 0.5 Bq/Kg生	23.4	Cs137	8.0 Bq/Kg生	
			Cs134	9.4 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Kiwi fruit	Okaona, Onahama, Iwaki	Nov-11	Cs137	15.9 Bq/Kg生	± 0.5 Bq/Kg生	24.9	Cs137	10.0 Bq/Kg生	
			Cs134	9.0 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	10.0 Bq/Kg生	
Kiwi fruit	Soeno, Iwaki	Nov-11	Cs137	5.0 Bq/Kg生	± 3.3 Bq/Kg生	13.2	Cs137	7.0 Bq/Kg生	
			Cs134	8.2 Bq/Kg生	± 3.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Apple	Ozaso, Fukushima	Dec-11	Cs137	16.3 Bq/Kg生	± 4.7 Bq/Kg生	30.9	Cs137	6.0 Bq/Kg生	
			Cs134	14.6 Bq/Kg生	± 3.8 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Persimmon	Kawauchi, Futaba	Nov-11	Cs137	8.9 Bq/Kg生	± 0.5 Bq/Kg生	18.6	Cs137	8.0 Bq/Kg生	
			Cs134	9.7 Bq/Kg生	± 0.7 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Persimmon	Nakachozakai, Onahama, Iwaki	Dec-11	Cs137	10.5 Bq/Kg生	± 0.5 Bq/Kg生	16.5	Cs137	8.0 Bq/Kg生	
			Cs134	6.0 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Persimmon	Noda, Onahama, Iwaki	Nov-11	Cs137	4.8 Bq/Kg生	± 0.4 Bq/Kg生	9.1	Cs137	8.0 Bq/Kg生	
			Cs134	4.3 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Dried persimmon	Ueda, Iwaki	Nov-11	Cs137	18.2 Bq/Kg生	± 5.0 Bq/Kg生	37.4	Cs137	6.0 Bq/Kg生	
			Cs134	19.2 Bq/Kg生	± 4.6 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Dried persimmon	Nakahirakubo, Taira, Iwaki	Dec-11	Cs137	23.9 Bq/Kg生	± 0.6 Bq/Kg生	46.4	Cs137	8.0 Bq/Kg生	
			Cs134	22.5 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Dried persimmon	Komagome, Yotsukura, Iwaki	Nov-11	Cs137	19.8 Bq/Kg生	± 0.6 Bq/Kg生	41.8	Cs137	8.0 Bq/Kg生	
			Cs134	22.0 Bq/Kg生	± 0.8 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Wild boar meat	Saiso, Miwa, Iwaki	Dec-11	Cs137	104 Bq/Kg生	± 21.0 Bq/Kg生	185	Cs137	6.0 Bq/Kg生	
			Cs134	81.0 Bq/Kg生	± 16.2 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Wild boar meat	Miwa, Iwaki	Unknown	Cs137	69.3 Bq/Kg生	± 15.1 Bq/Kg生	131	Cs137	7.0 Bq/Kg生	
			Cs134	62.1 Bq/Kg生	± 12.6 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Wild boar meat	Miwa, Iwaki	Dec-11	Cs137	90.1 Bq/Kg生	± 1.0 Bq/Kg生	167	Cs137	8.0 Bq/Kg生	
			Cs134	77.0 Bq/Kg生	± 1.2 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Wild boar meat	Fujiwara, Joban, Iwaki	Dec-11	Cs137	0.6 Bq/Kg生	± 0.4 Bq/Kg生	0.6	Cs137	10.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0 Bq/Kg生	

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Wild boar meat	Fujiwara, Joban, Iwaki	Nov-11	Cs137	119 Bq/Kg生	± 1.2 Bq/Kg生	221	Cs137	10.0	Bq/Kg生
			Cs134	102 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	10.0	Bq/Kg生
Honey	Yotsukura, Iwaki	Oct-11	Cs137	70.2 Bq/Kg生	± 15.1 Bq/Kg生	135	Cs137	6.0	Bq/Kg生
			Cs134	64.4 Bq/Kg生	± 12.9 Bq/Kg生		Cs134	4.0	Bq/Kg生
Milk	Honmiya	Unknown	Cs137	1.3 Bq/Kg生	± 0.2 Bq/Kg生	1.3	Cs137	5.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	5.0	Bq/Kg生
Milk (School lunch)	Honmiya	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	10.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0	Bq/Kg生
Strawberry yogurt	Unknown	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0	Bq/Kg生
Milk powder	Unknown	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0	Bq/Kg生
Milk powder	Unknown	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0	Bq/Kg生
Ramen (driednoodles)	Wheat(Foreign) Seaweed(Soma)	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	5.0	Cs137	6.0	Bq/Kg生
			Cs134	5.0 Bq/Kg生	± 2.7 Bq/Kg生		Cs134	4.0	Bq/Kg生
Ramen noodles	Unknown	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0	Bq/Kg生
Salted plum	Okaona, Onahama, Iwaki	Jun-11	Cs137	24.8 Bq/Kg生	± 6.3 Bq/Kg生	49.1	Cs137	6.0	Bq/Kg生
			Cs134	24.3 Bq/Kg生	± 5.6 Bq/Kg生		Cs134	4.0	Bq/Kg生
Pickled vegetables	Hirakubo, Iwaki	Nov-11	Cs137	2.4 Bq/Kg生	± 2.3 Bq/Kg生	5.2	Cs137	6.0	Bq/Kg生
			Cs134	2.8 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	4.0	Bq/Kg生
Tofu	Hokkaido	Oct-10	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.2	Cs137	6.0	Bq/Kg生
			Cs134	2.2 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	4.0	Bq/Kg生
Natto	Unknown	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	8.0	Bq/Kg生
Mixed vegetables	Shimoarakawa, Taira, Iwaki	Dec-11	Cs137	5.7 Bq/Kg生	± 0.4 Bq/Kg生	13.3	Cs137	10.0	Bq/Kg生
			Cs134	7.6 Bq/Kg生	± 0.6 Bq/Kg生		Cs134	10.0	Bq/Kg生
School Lunch	Uchigo, Iwaki	Dec-11	Cs137	3.9 Bq/Kg生	± 2.2 Bq/Kg生	3.9	Cs137	6.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0	Bq/Kg生
School Lunch	Uchigo, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.3	Cs137	6.0	Bq/Kg生
			Cs134	3.3 Bq/Kg生	± 2.0 Bq/Kg生		Cs134	4.0	Bq/Kg生
School Lunch	Kamiyunagaya, Joban, Iwaki	Dec-11	Cs137	3.3 Bq/Kg生	± 2.5 Bq/Kg生	3.3	Cs137	6.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0	Bq/Kg生
A Meal	Unknown	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	4.1 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0	Bq/Kg生
3 Meals	Unknown	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.9	Cs137	7.0	Bq/Kg生
			Cs134	3.9 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0	Bq/Kg生
Sawdust	Gunma	May-11	Cs137	23.8 Bq/Kg生	± 37.0 Bq/Kg生	52.1	Cs137	6.0	Bq/Kg生
			Cs134	28.3 Bq/Kg生	± 7.8 Bq/Kg生		Cs134	4.0	Bq/Kg生
Tap water	Kamiogawa, Ogawa, Iwaki	Dec-11	Cs137	1.8 Bq/Kg生	± 0.2 Bq/Kg生	1.8	Cs137	10.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0	Bq/Kg生
Tap water	Shimohirakubo, Taira, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	5.0	Bq/Kg生
Tap water	Kashimma, Chuodai, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	10.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	10.0	Bq/Kg生
Tap water	Okaona, Onahama, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	5.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Well water (spring water)	Ronden, Furudono, Ishikawa	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water (spring water)	Nakadera, Miwa, Iwaki	Dec-11	Cs137	16.3	Bq/Kg生	± 1.6	Bq/Kg生	35.5	Cs137 10.0 Bq/Kg生
			Cs134	19.2	Bq/Kg生	± 2.0	Bq/Kg生		Cs134 10.0 Bq/Kg生
Well water	Nakadera, Miwa, Iwaki	Dec-11	Cs137	1.0	Bq/Kg生	± 0.2	Bq/Kg生	1.0	Cs137 8.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 8.0 Bq/Kg生
Well water	Shimonagayama, Miwa, Iwaki	Dec-11	Cs137	22.2	Bq/Kg生	± 0.3	Bq/Kg生	49.4	Cs137 5.0 Bq/Kg生
			Cs134	27.2	Bq/Kg生	± 0.4	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Shimonagayama, Miwa, Iwaki	Dec-11	Cs137	2.3	Bq/Kg生	± 1.5	Bq/Kg生	2.3	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 3.0 Bq/Kg生
Well water	Shimonagayama, Miwa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.0	Cs137 5.0 Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.4	Bq/Kg生		Cs134 3.0 Bq/Kg生
Well water	Shimonagayama, Miwa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.8	Cs137 7.0 Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.3	Bq/Kg生		Cs134 6.0 Bq/Kg生
Well water	Godo, Miwa, Iwaki	Dec-11	Cs137	14.7	Bq/Kg生	± 0.3	Bq/Kg生	33.9	Cs137 5.0 Bq/Kg生
			Cs134	19.2	Bq/Kg生	± 0.4	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Miwa, Iwaki	Dec-11	Cs137	2.6	Bq/Kg生	± 1.5	Bq/Kg生	4.6	Cs137 7.0 Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.3	Bq/Kg生		Cs134 6.0 Bq/Kg生
Well water	Shimonagayama, Miwa, Iwaki	Dec-11	Cs137	12.0	Bq/Kg生	± 0.3	Bq/Kg生	28.2	Cs137 10.0 Bq/Kg生
			Cs134	16.2	Bq/Kg生	± 0.4	Bq/Kg生		Cs134 10.0 Bq/Kg生
Well water	Shimoichigaya, Miwa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 8.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 8.0 Bq/Kg生
Well water (domestic water)	Miwa, Iwaki	Dec-11	Cs137	16.3	Bq/Kg生	± 0.3	Bq/Kg生	35.5	Cs137 10.0 Bq/Kg生
			Cs134	19.2	Bq/Kg生	± 0.4	Bq/Kg生		Cs134 10.0 Bq/Kg生
Well water	Koya, Uchigo, Iwaki	Dec-11	Cs137	10.4	Bq/Kg生	± 0.2	Bq/Kg生	21.2	Cs137 5.0 Bq/Kg生
			Cs134	10.8	Bq/Kg生	± 0.4	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Hosoya, Yotsukura, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water (above)	Shibahara, Ogawa, Iwaki	Dec-11	Cs137	9.1	Bq/Kg生	± 0.2	Bq/Kg生	16.4	Cs137 10.0 Bq/Kg生
			Cs134	7.3	Bq/Kg生	± 0.4	Bq/Kg生		Cs134 10.0 Bq/Kg生
Well water (below)	Shibahara, Ogawa, Iwaki	Dec-11	Cs137	5.1	Bq/Kg生	± 0.2	Bq/Kg生	13.3	Cs137 10.0 Bq/Kg生
			Cs134	8.2	Bq/Kg生	± 0.3	Bq/Kg生		Cs134 10.0 Bq/Kg生
Well water	Iritono, Tono, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Oodaira, Tono, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Hosoya, Yotsukura, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Tamayama, Yotsukura, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Shibahara, Ogawa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water	Nakahirakubo, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 5.0 Bq/Kg生
Well water (domestic water)	Hisanohama, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 10.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 10.0 Bq/Kg生
Well water (domestic water)	Tanoami, Hisanohama, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137 10.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134 10.0 Bq/Kg生

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Well water (spring water)	Tabito, iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	10.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	10.0 Bq/Kg生
Well water	Koya, Uchigo, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	10.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	10.0 Bq/Kg生
Groundwater (domestic water)	Godo, Miwa, Iwaki	Dec-11	Cs137	35.2	Bq/Kg生	± 0.4	Bq/Kg生	87.7	Cs137	5.0 Bq/Kg生
			Cs134	52.5	Bq/Kg生	± 0.5	Bq/Kg生		Cs134	5.0 Bq/Kg生
Spring water (drinking water)	Kaminagai, Miwa, Iwaki	Dec-11	Cs137	3.3	Bq/Kg生	± 6.3	Bq/Kg生	4.7	Cs137	10.0 Bq/Kg生
			Cs134	1.4	Bq/Kg生	± 22.2	Bq/Kg生		Cs134	10.0 Bq/Kg生
Spring water (drinking water)	Saiso, Miwa, Iwaki	Dec-11	Cs137	0.9	Bq/Kg生	± 0.2	Bq/Kg生	0.9	Cs137	10.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	10.0 Bq/Kg生
Spring water	Minaiodaira, Tabito, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	5.0 Bq/Kg生
Spring water	Shimomisaka, Miwa, Iwaki	Dec-11	Cs137	1.3	Bq/Kg生	± 0.2	Bq/Kg生	1.4	Cs137	5.0 Bq/Kg生
			Cs134	0.1	Bq/Kg生	± 0.5	Bq/Kg生		Cs134	5.0 Bq/Kg生
Spring water	Godo, Miwa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	5.0 Bq/Kg生

