



Radiation Measurement Results of 328 Items in January 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Brown rice	Aomori	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.4	Cs137	6.0 Bq/Kg生		
			Cs134	2.4 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Brown rice	Yamagata	Oct-11	Cs137	3.7 Bq/Kg生	± 3.0 Bq/Kg生	8.5	Cs137	7.0 Bq/Kg生		
			Cs134	4.8 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Yamagata	Unknown	Cs137	4.4 Bq/Kg生	± 2.8 Bq/Kg生	8.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.7 Bq/Kg生	± 2.5 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Niigata	Oct-11	Cs137	3.2 Bq/Kg生	± 2.5 Bq/Kg生	3.2	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Niigata, Niigata	Oct-11	Cs137	4.5 Bq/Kg生	± 2.5 Bq/Kg生	4.5	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Ibaraki	Oct-11	Cs137	4.1 Bq/Kg生	± 2.7 Bq/Kg生	6.4	Cs137	7.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Kumamoto	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.1 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Isohara, Ibaraki	Oct-11	Cs137	2.4 Bq/Kg生	± 1.9 Bq/Kg生	5.4	Cs137	6.0 Bq/Kg生		
			Cs134	3.0 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Brown rice	Hanakawa, Kitaibaraki, Ibaraki	Oct-11	Cs137	3.4 Bq/Kg生	± 2.6 Bq/Kg生	7.5	Cs137	7.0 Bq/Kg生		
			Cs134	4.1 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Futaba	Sep-11	Cs137	2.9 Bq/Kg生	± 2.7 Bq/Kg生	2.9	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Naraha, Futaba	Oct-10	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.6	Cs137	7.0 Bq/Kg生		
			Cs134	3.6 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Namie, Futaba	Nov-10	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.9	Cs137	7.0 Bq/Kg生		
			Cs134	3.9 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Aizu	Nov-11	Cs137	4.2 Bq/Kg生	± 2.5 Bq/Kg生	7.0	Cs137	7.0 Bq/Kg生		
			Cs134	2.8 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Tanagura, Higashishirakawa	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.8	Cs137	6.0 Bq/Kg生		
			Cs134	1.8 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Brown rice	Yabuki, Nishishirakawa	Oct-11	Cs137	2.9 Bq/Kg生	± 2.0 Bq/Kg生	6.3	Cs137	6.0 Bq/Kg生		
			Cs134	3.4 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Brown rice	Ishikawa, Ishikawa	Oct-11	Cs137	2.8 Bq/Kg生	± 2.4 Bq/Kg生	6.3	Cs137	7.0 Bq/Kg生		
			Cs134	3.5 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Takasato, Kitakata	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.6	Cs137	7.0 Bq/Kg生		
			Cs134	2.6 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Brown rice	Miwa, Iwaki	Nov-11	Cs137	2.2 Bq/Kg生	± 2.1 Bq/Kg生	4.2	Cs137	7.0 Bq/Kg生		
			Cs134	2.0 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	6.0 Bq/Kg生		

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Hosoya, Yotsukura, Iwaki	Nov-11	Cs137	5.3	Bq/Kg生	± 3.0	Bq/Kg生	10.6	Cs137	7.0	Bq/Kg生
			Cs134	5.3	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	6.1	Bq/Kg生	± 3.2	Bq/Kg生	10.2	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimokabeya, Taira, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 3.1	Bq/Kg生	8.2	Cs137	7.0	Bq/Kg生
			Cs134	5.1	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kabeya, Taira, Iwaki	Nov-11	Cs137	3.2	Bq/Kg生	± 2.2	Bq/Kg生	9.5	Cs137	5.0	Bq/Kg生
			Cs134	6.3	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	3.0	Bq/Kg生
Brown rice	Takaku, Taira, Iwaki	Oct-11	Cs137	6.6	Bq/Kg生	± 3.0	Bq/Kg生	10.6	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Takaku, Taira, Iwaki	Oct-11	Cs137	6.9	Bq/Kg生	± 2.5	Bq/Kg生	13.8	Cs137	6.0	Bq/Kg生
			Cs134	6.9	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Kamiktakaku, Taira, Iwaki	Oct-11	Cs137	7.4	Bq/Kg生	± 3.2	Bq/Kg生	10.0	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Akai, Taira, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Akai, Taira, Iwaki	Oct-11	Cs137	4.0	Bq/Kg生	± 2.7	Bq/Kg生	6.4	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Akai, Taira, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.5	Bq/Kg生	8.3	Cs137	7.0	Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nakahirakubo, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.6	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Hirakubo, Taira, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.5	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	4.2	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Hirakubo, Taira, Iwaki	Nov-11	Cs137	4.8	Bq/Kg生	± 3.1	Bq/Kg生	8.9	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ogawa, Iwaki	Nov-11	Cs137	6.4	Bq/Kg生	± 3.0	Bq/Kg生	10.4	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ogawa, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.8	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Ogawa, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Kuwabara, Ogawa, Iwaki	Oct-11	Cs137	8.6	Bq/Kg生	± 3.4	Bq/Kg生	15.0	Cs137	7.0	Bq/Kg生
			Cs134	6.4	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kamiogawa, Ogawa, Iwaki	Oct-11	Cs137	2.3	Bq/Kg生	± 1.9	Bq/Kg生	5.8	Cs137	6.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Shimoogawa, Ogawa, Iwaki	Oct-11	Cs137	5.5	Bq/Kg生	± 2.8	Bq/Kg生	9.3	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimoogawa, Ogawa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimoogawa, Ogawa, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.8	Bq/Kg生	6.8	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Yoshima, Iwaki	Sep-11	Cs137	2.2	Bq/Kg生	± 2.1	Bq/Kg生	5.4	Cs137	6.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Sakakigoya, Yoshima, Iwaki	Nov-11	Cs137	5.1	Bq/Kg生	± 2.8	Bq/Kg生	8.9	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kamiyunagaya, Joban, Iwaki	Sep-11	Cs137	3.1	Bq/Kg生	± 2.7	Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	4.2	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Misawa, Joban, Iwaki	Oct-11	Cs137	2.8 Bq/Kg生	± 2.5 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生	
			Cs134	4.1 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Iritono, Tono, Iwaki	Oct-11	Cs137	2.9 Bq/Kg生	± 2.0 Bq/Kg生	2.9	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.2	Cs137	7.0 Bq/Kg生	
			Cs134	4.2 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Tono, Iwaki	Oct-11	Cs137	4.7 Bq/Kg生	± 2.7 Bq/Kg生	4.7	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Iritono, Tono, Iwaki	Oct-11	Cs137	3.6 Bq/Kg生	± 2.5 Bq/Kg生	7.2	Cs137	7.0 Bq/Kg生	
			Cs134	3.6 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Negishi, Tono, Iwaki	Oct-11	Cs137	4.6 Bq/Kg生	± 2.8 Bq/Kg生	7.6	Cs137	7.0 Bq/Kg生	
			Cs134	3.0 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Negishi, Tono, Iwaki	Oct-11	Cs137	5.2 Bq/Kg生	± 2.7 Bq/Kg生	8.2	Cs137	7.0 Bq/Kg生	
			Cs134	3.0 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Taki, Tono, Iwaki	Nov-11	Cs137	6.0 Bq/Kg生	± 2.6 Bq/Kg生	10.0	Cs137	6.0 Bq/Kg生	
			Cs134	4.0 Bq/Kg生	± 2.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Taki, Tono, Iwaki	Nov-11	Cs137	3.4 Bq/Kg生	± 2.3 Bq/Kg生	7.1	Cs137	6.0 Bq/Kg生	
			Cs134	3.7 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Soeno, Iwaki	Nov-11	Cs137	3.1 Bq/Kg生	± 2.0 Bq/Kg生	5.8	Cs137	6.0 Bq/Kg生	
			Cs134	2.7 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Watanabe, Iwaki	Oct-11	Cs137	4.3 Bq/Kg生	± 2.8 Bq/Kg生	8.9	Cs137	7.0 Bq/Kg生	
			Cs134	4.6 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Kawabe, Iwaki	Sep-11	Cs137	7.5 Bq/Kg生	± 3.3 Bq/Kg生	14.8	Cs137	7.0 Bq/Kg生	
			Cs134	7.3 Bq/Kg生	± 2.8 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Tabito, Iwaki	Unknown	Cs137	4.2 Bq/Kg生	± 2.7 Bq/Kg生	7.0	Cs137	7.0 Bq/Kg生	
			Cs134	2.8 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Misawa, Iwaki	Oct-11	Cs137	7.2 Bq/Kg生	± 3.5 Bq/Kg生	17.2	Cs137	7.0 Bq/Kg生	
			Cs134	10.0 Bq/Kg生	± 3.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Nakoso, Iwaki	Oct-11	Cs137	2.9 Bq/Kg生	± 2.8 Bq/Kg生	5.6	Cs137	7.0 Bq/Kg生	
			Cs134	2.7 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Nakoso, Iwaki	Oct-11	Cs137	5.2 Bq/Kg生	± 2.3 Bq/Kg生	8.2	Cs137	6.0 Bq/Kg生	
			Cs134	3.0 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Nakaoka, Iwaki	Sep-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.4	Cs137	7.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Aomori Pref.	Oct-11	Cs137	2.6 Bq/Kg生	± 2.4 Bq/Kg生	2.6	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Aomori Pref.	Nov-11	Cs137	2.4 Bq/Kg生	± 2.3 Bq/Kg生	4.8	Cs137	7.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Yamagata Pref.	Unknown	Cs137	3.9 Bq/Kg生	± 2.6 Bq/Kg生	7.4	Cs137	7.0 Bq/Kg生	
			Cs134	3.5 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Obanazawa, Yamagata Pref.	Oct-11	Cs137	3.7 Bq/Kg生	± 2.4 Bq/Kg生	6.1	Cs137	7.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Misato, Toda, Miyagi Pref.	Oct-11	Cs137	5.1 Bq/Kg生	± 2.6 Bq/Kg生	7.5	Cs137	7.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Fukui Pref.	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.9	Cs137	6.0 Bq/Kg生	
			Cs134	1.9 Bq/Kg生	± 1.6 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Ibaraki Pref.	Oct-11	Cs137	2.3 Bq/Kg生	± 2.3 Bq/Kg生	2.3	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Kawajiri, Kitaibaraki	Oct-11	Cs137	2.8	Bq/Kg生	± 1.9	Bq/Kg生	6.8	Cs137	6.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Naraha, Futaba	Oct-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ono, Tamura	Oct-11	Cs137	3.4	Bq/Kg生	± 2.5	Bq/Kg生	3.4	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Date	Oct-11	Cs137	2.6	Bq/Kg生	± 2.0	Bq/Kg生	5.3	Cs137	6.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Nihonmatsu	Unknown	Cs137	2.3	Bq/Kg生	± 1.9	Bq/Kg生	4.2	Cs137	6.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Aizu	Nov-11	Cs137	4.4	Bq/Kg生	± 2.4	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Aizumisato, Onuma	Unknown	Cs137	2.0	Bq/Kg生	± 2.0	Bq/Kg生	2.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Miwa, Iwaki	Unknown	Cs137	2.6	Bq/Kg生	± 2.1	Bq/Kg生	5.0	Cs137	6.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Saiso, Miwa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.1	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Saiso, Miwa, Iwaki	Oct-11	Cs137	4.7	Bq/Kg生	± 2.4	Bq/Kg生	7.5	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Unknown	Cs137	4.2	Bq/Kg生	± 2.8	Bq/Kg生	7.0	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Nov-11	Cs137	5.1	Bq/Kg生	± 2.7	Bq/Kg生	8.2	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.6	Bq/Kg生	6.5	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimoyagyu, Yotsukura, Iwaki	Oct-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.0	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimoyagyu, Yotsukura, Iwaki	May-11	Cs137	3.5	Bq/Kg生	± 2.7	Bq/Kg生	6.1	Cs137	6.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Nagatomo, Yotsukura, Iwaki	Nov-11	Cs137	5.1	Bq/Kg生	± 2.3	Bq/Kg生	5.1	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kawamae, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.6	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Haragoya, Taira, Iwaki	Sep-11	Cs137	3.1	Bq/Kg生	± 2.7	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Oomuro, Taira, Iwaki	Oct-11	Cs137	3.2	Bq/Kg生	± 2.5	Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimokabeya, Taira, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.5	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Akai, Taira, Iwaki	Oct-11	Cs137	4.2	Bq/Kg生	± 2.3	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Akai, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.0	Bq/Kg生	5.1	Cs137	6.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.7	Bq/Kg生	8.3	Cs137		Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134		Bq/Kg生
Rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.7	Bq/Kg生	8.3	Cs137		Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134		Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Rice	Fujima, Taira, Iwaki	Oct-11	Cs137	2.0 Bq/Kg生	± 2.0 Bq/Kg生	5.1	Cs137	6.0 Bq/Kg生		
			Cs134	3.1 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Shimoyamaguchi, Taira, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.2	Cs137	7.0 Bq/Kg生		
			Cs134	4.2 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.8	Cs137	7.0 Bq/Kg生		
			Cs134	4.8 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kabeya, Taira, Iwaki	Nov-11	Cs137	3.4 Bq/Kg生	± 2.7 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生		
			Cs134	3.5 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Suginami, Taira, Iwaki	Oct-11	Cs137	2.7 Bq/Kg生	± 2.0 Bq/Kg生	5.4	Cs137	6.0 Bq/Kg生		
			Cs134	2.7 Bq/Kg生	± 1.6 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Suginami, Taira, Iwaki	Oct-11	Cs137	3.3 Bq/Kg生	± 2.0 Bq/Kg生	5.5	Cs137	6.0 Bq/Kg生		
			Cs134	2.2 Bq/Kg生	± 1.6 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Nakashio, Taira, Iwaki	Oct-11	Cs137	2.8 Bq/Kg生	± 2.3 Bq/Kg生	5.1	Cs137	7.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Katayose, Taira, Iwaki	Oct-11	Cs137	3.8 Bq/Kg生	± 2.5 Bq/Kg生	7.8	Cs137	7.0 Bq/Kg生		
			Cs134	4.0 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Ogawa, Iwaki	Nov-11	Cs137	2.8 Bq/Kg生	± 2.5 Bq/Kg生	6.7	Cs137	7.0 Bq/Kg生		
			Cs134	3.9 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Ogawa, Iwaki	Oct-11	Cs137	4.5 Bq/Kg生	± 2.4 Bq/Kg生	4.5	Cs137	6.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Ogawa, Iwaki	Oct-11	Cs137	3.7 Bq/Kg生	± 2.3 Bq/Kg生	5.8	Cs137	7.0 Bq/Kg生		
			Cs134	2.1 Bq/Kg生	± 2.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kamiogawa, Ogawa, Iwaki	Oct-11	Cs137	2.7 Bq/Kg生	± 2.4 Bq/Kg生	5.8	Cs137	7.0 Bq/Kg生		
			Cs134	3.1 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Takasaki, Ogawa, Iwaki	Oct-11	Cs137	2.6 Bq/Kg生	± 2.0 Bq/Kg生	2.6	Cs137	6.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Takasaki, Ogawa, Iwaki	Oct-11	Cs137	3.1 Bq/Kg生	± 2.0 Bq/Kg生	5.6	Cs137	6.0 Bq/Kg生		
			Cs134	2.5 Bq/Kg生	± 1.6 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Yoshima, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.4	Cs137	7.0 Bq/Kg生		
			Cs134	3.4 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kamiyoshima, Yoshima, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.0	Cs137	7.0 Bq/Kg生		
			Cs134	4.0 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kouya, Uchigo, Iwaki	Oct-11	Cs137	3.0 Bq/Kg生	± 2.4 Bq/Kg生	5.6	Cs137	7.0 Bq/Kg生		
			Cs134	2.6 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.3	Cs137	7.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	4.5 Bq/Kg生	± 5.4 Bq/Kg生	8.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.6 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	3.5 Bq/Kg生	± 2.4 Bq/Kg生	6.1	Cs137	7.0 Bq/Kg生		
			Cs134	2.6 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kamiyunagaya, Joban, Iwaki	Jan-12	Cs137	4.0 Bq/Kg生	± 2.5 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生		
			Cs134	2.9 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kamiyada, Joban, Iwaki	Oct-11	Cs137	3.7 Bq/Kg生	± 2.7 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kuramochi, Kasima, Iwaki	Unknown	Cs137	3.4 Bq/Kg生	± 2.5 Bq/Kg生	6.3	Cs137	7.0 Bq/Kg生		
			Cs134	2.9 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Watanabe, Iwaki	Nov-11	Cs137	4.0 Bq/Kg生	± 2.6 Bq/Kg生	7.9	Cs137	7.0 Bq/Kg生		
			Cs134	3.9 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Rice	Watanabe, Iwaki	Nov-11	Cs137	3.1 Bq/Kg生	± 2.4 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生	
			Cs134	3.8 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Watanabe, Iwaki	Oct-11	Cs137	2.1 Bq/Kg生	± 2.1 Bq/Kg生	4.9	Cs137	6.0 Bq/Kg生	
			Cs134	2.8 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Tanabe, Watanabe, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.1	Cs137	7.0 Bq/Kg生	
			Cs134	4.1 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kamado, Watanabe, Iwaki	Unknown	Cs137	3.9 Bq/Kg生	± 2.6 Bq/Kg生	6.4	Cs137	7.0 Bq/Kg生	
			Cs134	2.5 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Tabito, Iwaki	Sep-11	Cs137	3.1 Bq/Kg生	± 2.4 Bq/Kg生	5.4	Cs137	7.0 Bq/Kg生	
			Cs134	2.3 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Tabito, Iwaki	Nov-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.0	Cs137	7.0 Bq/Kg生	
			Cs134	3.0 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kuroda, Tabito, Iwaki	Oct-11	Cs137	4.9 Bq/Kg生	± 2.6 Bq/Kg生	7.5	Cs137	7.0 Bq/Kg生	
			Cs134	2.6 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kuroda, Tabito, Iwaki	Oct-11	Cs137	3.3 Bq/Kg生	± 2.6 Bq/Kg生	6.8	Cs137	7.0 Bq/Kg生	
			Cs134	3.5 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Misawa, Iwaki	Oct-11	Cs137	4.9 Bq/Kg生	± 3.0 Bq/Kg生	9.2	Cs137	7.0 Bq/Kg生	
			Cs134	4.3 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Nakoso, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.9	Cs137	7.0 Bq/Kg生	
			Cs134	2.9 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kawahara, Kawabe, Iwaki	Sep-11	Cs137	4.8 Bq/Kg生	± 2.2 Bq/Kg生	8.6	Cs137	6.0 Bq/Kg生	
			Cs134	3.8 Bq/Kg生	± 1.8 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	2.5 Bq/Kg生	± 2.4 Bq/Kg生	2.5	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Taki, Tono, Iwaki	Oct-11	Cs137	2.9 Bq/Kg生	± 1.9 Bq/Kg生	2.9	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Tono, Iwaki	Oct-11	Cs137	2.3 Bq/Kg生	± 2.0 Bq/Kg生	4.0	Cs137	6.0 Bq/Kg生	
			Cs134	1.7 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生	
			Cs134	3.7 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Iritono, Tono, Iwaki	Dec-11	Cs137	2.8 Bq/Kg生	± 2.7 Bq/Kg生	6.8	Cs137	7.0 Bq/Kg生	
			Cs134	4.0 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Tono, Iwaki	Sep-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Tono, Iwaki	Nov-11	Cs137	2.8 Bq/Kg生	± 2.4 Bq/Kg生	5.2	Cs137	7.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Takijiri, Izumi, Iwaki	Sep-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.5	Cs137	7.0 Bq/Kg生	
			Cs134	3.5 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kawabe, Iwaki	Sep-11	Cs137	6.1 Bq/Kg生	± 3.0 Bq/Kg生	10.8	Cs137	7.0 Bq/Kg生	
			Cs134	4.7 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Shirokajiro, Onahama, Iwaki	Nov-11	Cs137	2.8 Bq/Kg生	± 2.4 Bq/Kg生	2.8	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kanari, Onahama, Iwaki	Sep-11	Cs137	4.7 Bq/Kg生	± 2.7 Bq/Kg生	11.1	Cs137	7.0 Bq/Kg生	
			Cs134	6.4 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Okaona, Onahama, Iwaki	Oct-11	Cs137	3.3 Bq/Kg生	± 2.9 Bq/Kg生	10.5	Cs137	7.0 Bq/Kg生	
			Cs134	7.2 Bq/Kg生	± 2.8 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Ohara, Onahama, Iwaki	Oct-11	Cs137	3.6 Bq/Kg生	± 2.4 Bq/Kg生	3.6	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Sanuka, Iwaki	Dec-11	Cs137	4.8	Bq/Kg生	± 2.8	Bq/Kg生	12.1	Cs137	7.0	Bq/Kg生
			Cs134	7.3	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nakaoka, Iwaki	Sep-11	Cs137	3.0	Bq/Kg生	± 2.3	Bq/Kg生	6.2	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yamada, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.0	Cs137	5.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	3.0	Bq/Kg生
Rice	Unknown	Unknown	Cs137	4.4	Bq/Kg生	± 2.5	Bq/Kg生	7.5	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 7.0	Bq/Kg生	2.2	Cs137	7.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 6.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (Brown rice)	Soeno, Iwaki	Nov-11	Cs137	3.4	Bq/Kg生	± 2.0	Bq/Kg生	3.4	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice (Brown rice)	Tono, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.8	Bq/Kg生	7.4	Cs137	7.0	Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice	Nakata, Koriyama	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.6	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice	Katahira, Koriyama	Nov-11	Cs137	4.4	Bq/Kg生	± 2.6	Bq/Kg生	7.7	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice	Yotsukura, Iwaki	Nov-11	Cs137	4.7	Bq/Kg生	± 2.7	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice	Kitamuki, Yotsukura, Iwaki	Nov-11	Cs137	11.2	Bq/Kg生	± 3.7	Bq/Kg生	20.4	Cs137	7.0	Bq/Kg生
			Cs134	9.2	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	2.6	Bq/Kg生	± 1.9	Bq/Kg生	5.1	Cs137	6.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice	Makunouchi, Taira, Iwaki	Oct-11	Cs137	10.2	Bq/Kg生	± 3.1	Bq/Kg生	17.8	Cs137	6.0	Bq/Kg生
			Cs134	7.6	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice	Kashima, Iwaki	Oct-11	Cs137	7.0	Bq/Kg生	± 3.2	Bq/Kg生	13.9	Cs137	7.0	Bq/Kg生
			Cs134	6.9	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Flour	Nagasaki, Iwaki	Jun-11	Cs137	50.7	Bq/Kg生	± 11.6	Bq/Kg生	98.7	Cs137	7.0	Bq/Kg生
			Cs134	48.0	Bq/Kg生	± 10.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Aizu	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Hirata, Ishikawa	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.5	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kamiyada, Joban, Iwaki	Aug-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.0	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Fujiwara, Joban, Iwaki	Jul-11	Cs137	5.8	Bq/Kg生	± 3.2	Bq/Kg生	10.9	Cs137	7.0	Bq/Kg生
			Cs134	5.1	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Ohara, Onahama, Iwaki	Jul-11	Cs137	4.5	Bq/Kg生	± 2.4	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Okaona, Onahama, Iwaki	Jul-11	Cs137	4.3	Bq/Kg生	± 2.7	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Ohara, Onahama, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.4	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kashima, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.5	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Hashirikuma, Kashima, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Potato	Shimoogawa, Iwaki	Jul-11	Cs137	3.5 Bq/Kg生	± 2.6 Bq/Kg生	6.2	Cs137	7.0 Bq/Kg生		
			Cs134	2.7 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Shioda, Ogawa, Iwaki	Jun-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	5.8	Cs137	7.0 Bq/Kg生		
			Cs134	5.8 Bq/Kg生	± 3.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Iritono, Tono, Iwaki	Jul-11	Cs137	4.8 Bq/Kg生	± 2.8 Bq/Kg生	7.5	Cs137	7.0 Bq/Kg生		
			Cs134	2.7 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Moegidai, Izumi, Iwaki	Jul-11	Cs137	5.5 Bq/Kg生	± 3.1 Bq/Kg生	5.5	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Takijiri, Izumi, Iwaki	Aug-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.0	Cs137	7.0 Bq/Kg生		
			Cs134	4.0 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Yamada, Iwaki	Jun-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Kawabe, Iwaki	Jun-11	Cs137	3.4 Bq/Kg生	± 2.9 Bq/Kg生	7.5	Cs137	7.0 Bq/Kg生		
			Cs134	4.1 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Komagome, Yotsukura, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.1 Bq/Kg生	± 3.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Kamiyoshima, Yoshima, Iwaki	Nov-11	Cs137	2.8 Bq/Kg生	± 2.2 Bq/Kg生	2.8	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Takaku, Chuodai, Iwaki	Jan-12	Cs137	7.6 Bq/Kg生	± 7.6 Bq/Kg生	7.6	Cs137	6.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Potato	Miyamachi, Uchigo, Iwaki	Jan-12	Cs137	4.7 Bq/Kg生	± 3.2 Bq/Kg生	4.7	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Akai, Taira, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	20.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	20.0 Bq/Kg生		
Potato	Numanouchi, Taira, Iwaki	Jul-11	Cs137	3.5 Bq/Kg生	± 2.8 Bq/Kg生	7.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.6 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Nakahirakubo, Taira, Iwaki	Jun-11	Cs137	1.2 Bq/Kg生	± 6.7 Bq/Kg生	5.9	Cs137	8.0 Bq/Kg生		
			Cs134	4.7 Bq/Kg生	± 5.3 Bq/Kg生		Cs134	8.0 Bq/Kg生		
Potato	Kitashirado, Taira, Iwaki	Jul-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	6.6	Cs137	7.0 Bq/Kg生		
			Cs134	6.6 Bq/Kg生	± 3.4 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Potato	Shimoyamaguchi, Taiara, Iwaki	Jun-11	Cs137	5.0 Bq/Kg生	± 3.0 Bq/Kg生	8.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.1 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Taro	Hirakubo, Taira, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.1	Cs137	7.0 Bq/Kg生		
			Cs134	4.1 Bq/Kg生	± 3.9 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Taro	Ohara, Onahama, Iwaki	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Sweet Potato	Moegidai, Izumi, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生		
			Cs134	3.7 Bq/Kg生	± 2.7 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Sweet Potato	Shimoyunagaya, Joban, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.9	Cs137	7.0 Bq/Kg生		
			Cs134	3.9 Bq/Kg生	± 2.7 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Sweet Potato	Takasaka, Uchigo, Iwaki	Sep-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Sweet Potato	Kamiyoshima, Yoshima, Iwaki	Oct-11	Cs137	4.8 Bq/Kg生	± 2.5 Bq/Kg生	7.4	Cs137	7.0 Bq/Kg生		
			Cs134	2.6 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Yam	Kamitakaku, Taira, Iwaki	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Yam	Shimogawa, Izumi, Iwaki	Jan-12	Cs137	4.4 Bq/Kg生	± 3.1 Bq/Kg生	4.4	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Yam	Kanari, Onahama, Iwaki	Dec-11	Cs137	2.8	Bq/Kg生	± 2.4	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried stems of taro	Moegidai, Izumi, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Pumpkin	Kamioogoe, Taira, Iwaki	Unknown	Cs137	6.1	Bq/Kg生	± 3.9	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Pumpkin	Shimoyunagaya, Joban, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 3.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Sekimoto, Kitaibaraki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.5	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimogawa, Izumi, Iwaki	Jan-12	Cs137	8.5	Bq/Kg生	± 5.0	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kuroda, Tabito, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.7	Cs137	7.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 4.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Yamada, Iwaki	Jan-12	Cs137	6.5	Bq/Kg生	± 5.2	Bq/Kg生	15.2	Cs137	7.0	Bq/Kg生
			Cs134	8.7	Bq/Kg生	± 4.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kamitakaku, Taira, Iwaki	Jan-12	Cs137	5.6	Bq/Kg生	± 4.8	Bq/Kg生	10.8	Cs137	7.0	Bq/Kg生
			Cs134	5.2	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Suginami, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Takaku, Taira, Iwaki	Jan-12	Cs137	5.2	Bq/Kg生	± 5.0	Bq/Kg生	10.9	Cs137	7.0	Bq/Kg生
			Cs134	5.7	Bq/Kg生	± 4.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shimogawa, Izumi, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.6	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shimogawa, Izumi, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.6	Cs137	7.0	Bq/Kg生
			Cs134	5.6	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Takijiri, Izumi, Iwaki	Jan-12	Cs137	6.1	Bq/Kg生	± 3.2	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Tamatsuyu, Izumi, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	5.0	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Moegidai, Izumi, Iwaki	Jan-12	Cs137	3.5	Bq/Kg生	± 3.3	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Hashirikuma, Kashima, Iwaki	Jan-12	Cs137	4.5	Bq/Kg生	± 2.7	Bq/Kg生	4.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Miyamada, Tono, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Iritono, Tono, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Takaku, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.1	Cs137	7.0	Bq/Kg生
			Cs134	5.1	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Hosoya, Yotsukura, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.7	Cs137	7.0	Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Kawabe, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Miya, Uchigo, Iwaki	Jan-12	Cs137	6.9	Bq/Kg生	± 3.3	Bq/Kg生	10.3	Cs137	7.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 3.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Gohoden, Nishiki, Iwaki	Jan-11	Cs137	2.9	Bq/Kg生	± 2.7	Bq/Kg生	5.8	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Japanese white radish	Kattsuo, Toyoma, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.7	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Cabbage and radish mix	Shimokajiro, Onahama, Iwaki	Jan-12	Cs137	4.6	Bq/Kg生	± 4.3	Bq/Kg生	4.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried radish	Moegidai, Izumi, Iwaki	Dec-11	Cs137	122	Bq/Kg生	± 35.0	Bq/Kg生	250	Cs137	7.0	Bq/Kg生
			Cs134	128	Bq/Kg生	± 35.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Carrot	Nakayama, Taira, Iwaki	Jan-12	Cs137	4.8	Bq/Kg生	± 6.6	Bq/Kg生	4.8	Cs137	8.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Carrot	Shimokabeya, Taira, Iwaki	Jan-12	Cs137	7.7	Bq/Kg生	± 4.0	Bq/Kg生	7.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Carrot	Shimohirakubo, Taira, Iwaki	Jan-12	Cs137	4.1	Bq/Kg生	± 3.4	Bq/Kg生	9.7	Cs137	7.0	Bq/Kg生
			Cs134	5.6	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Nakashio, Taira, Iwaki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.6	Cs137	7.0	Bq/Kg生
			Cs134	4.6	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Shioda, Ogawa, Iwaki	Jun-11	Cs137	5.8	Bq/Kg生	± 3.8	Bq/Kg生	9.8	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 3.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Tamagawa, Onahama, Iwaki	Jan-12	Cs137	3.3	Bq/Kg生	± 3.1	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Moegidai, Izumi, Iwaki	Jun-11	Cs137	5.0	Bq/Kg生	± 3.4	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Gohoden, Nishiki, Iwaki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.5	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Sekita, Nakoso, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Shimokabeya, Taira, Iwaki	Jan-12	Cs137	6.0	Bq/Kg生	± 4.8	Bq/Kg生	11.9	Cs137	7.0	Bq/Kg生
			Cs134	5.9	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Kamitakaku, Taira, Iwaki	Oct-12	Cs137	4.2	Bq/Kg生	± 3.8	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Fujiwara, Joban, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	6.3	Bq/Kg生	± 4.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Okaona, Onahama, Iwaki	Jan-12	Cs137	4.7	Bq/Kg生	± 4.5	Bq/Kg生	11.0	Cs137	7.0	Bq/Kg生
			Cs134	6.3	Bq/Kg生	± 3.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Ohara, Onahama, Iwaki	Jan-12	Cs137	3.8	Bq/Kg生	± 2.5	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Burdock	Moegidai, Izumi, Iwaki	Jan-12	Cs137	5.5	Bq/Kg生	± 4.0	Bq/Kg生	10.2	Cs137	7.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 3.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yacon	Shimoogoe, Taira, Iwaki	Dec-11	Cs137	3.8	Bq/Kg生	± 2.2	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Cabbage	Nishigo, Joban, Iwaki	Dec-11	Cs137	5.5	Bq/Kg生	± 5.4	Bq/Kg生	11.5	Cs137	7.0	Bq/Kg生
			Cs134	6.0	Bq/Kg生	± 4.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Cabbage	Sugawara, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Cabbage	Kuroda, Tabito, Iwaki	Jan-12	Cs137	8.3	Bq/Kg生	± 5.1	Bq/Kg生	13.9	Cs137	7.0	Bq/Kg生
			Cs134	5.6	Bq/Kg生	± 4.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Cabbage	Kattsuo, Toyoma, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Broccoli	Shimogawa, Izumi, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.8	Cs137	7.0	Bq/Kg生
			Cs134	6.8	Bq/Kg生	± 4.1	Bq/Kg生		Cs134	6.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rape plant	Nakayama, Taira, Iwaki	Jan-11	Cs137	2.2	Bq/Kg生	± 4.8	Bq/Kg生	4.4	Cs137	8.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Spinach	Akai, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	20.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	20.0	Bq/Kg生
Greens	Maebara, Ogawa, Iwaki	Jan-12	Cs137	7.8	Bq/Kg生	± 10.4	Bq/Kg生	9.5	Cs137	8.0	Bq/Kg生
			Cs134	1.7	Bq/Kg生	± 11.4	Bq/Kg生		Cs134	8.0	Bq/Kg生
Qing-geng-cai	Tamagawa, Onahama, Iwaki	Jan-12	Cs137	3.7	Bq/Kg生	± 3.6	Bq/Kg生	3.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese mustard spinach	Akai, Taira, Iwaki	Jan-12	Cs137	10.1	Bq/Kg生	± 8.1	Bq/Kg生	17.7	Cs137	7.0	Bq/Kg生
			Cs134	7.6	Bq/Kg生	± 7.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Ohara, Onahama, Iwaki	Jan-12	Cs137	3.8	Bq/Kg生	± 2.5	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rape	Takaku, Chuodai, Iwaki	Jan-12	Cs137	5.8	Bq/Kg生	± 4.3	Bq/Kg生	11.5	Cs137	7.0	Bq/Kg生
			Cs134	5.7	Bq/Kg生	± 3.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Turnip rape	Nagasaki, Iwaki	Jan-12	Cs137	10.3	Bq/Kg生	± 8.3	Bq/Kg生	23.2	Cs137	7.0	Bq/Kg生
			Cs134	12.9	Bq/Kg生	± 6.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Purple stem mustard	Shimoogoe, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	6.7	Bq/Kg生	± 3.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mandarin orange	Iayuzaka, Yotsukura, Iwaki	Jan-12	Cs137	77.9	Bq/Kg生	± 16.7	Bq/Kg生	142	Cs137	7.0	Bq/Kg生
			Cs134	64.3	Bq/Kg生	± 13.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mandarin orange	Izumigaoka, Iwaki	Jan-12	Cs137	24.3	Bq/Kg生	± 8.1	Bq/Kg生	37.0	Cs137	8.0	Bq/Kg生
			Cs134	12.7	Bq/Kg生	± 5.6	Bq/Kg生		Cs134	8.0	Bq/Kg生
Mandarin orange	Sanuka, Iwaki	Dec-11	Cs137	4.8	Bq/Kg生	± 2.8	Bq/Kg生	12.1	Cs137	7.0	Bq/Kg生
			Cs134	7.3	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mandarin orange	Onahama, Iwaki	Jan-12	Cs137	10.9	Bq/Kg生	± 7.0	Bq/Kg生	20.1	Cs137	8.0	Bq/Kg生
			Cs134	9.2	Bq/Kg生	± 4.7	Bq/Kg生		Cs134	8.0	Bq/Kg生
Mandarin orange	Shirokajiro, Onahama, Iwaki	Jan-11	Cs137	24.3	Bq/Kg生	± 6.1	Bq/Kg生	38.6	Cs137	7.0	Bq/Kg生
			Cs134	14.3	Bq/Kg生	± 3.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mandarin orange	Kitanosaku, Obama, Iwaki	Dec-11	Cs137	7.5	Bq/Kg生	± 6.3	Bq/Kg生	20.9	Cs137	8.0	Bq/Kg生
			Cs134	13.4	Bq/Kg生	± 4.9	Bq/Kg生		Cs134	8.0	Bq/Kg生
Mandarin orange	Takasaka, Uchigo, Iwaki	Nov-11	Cs137	19.3	Bq/Kg生	± 4.9	Bq/Kg生	32.5	Cs137	7.0	Bq/Kg生
			Cs134	13.2	Bq/Kg生	± 3.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mandarin orange	Nishiogawa, Ogawa, Iwaki	Jan-12	Cs137	11.7	Bq/Kg生	± 5.8	Bq/Kg生	14.5	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Apple	Senoue, Fukushima	Dec-11	Cs137	12.7	Bq/Kg生	± 5.2	Bq/Kg生	24.1	Cs137	7.0	Bq/Kg生
			Cs134	11.4	Bq/Kg生	± 4.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Citron	Iayuzaka, Yotsukura, Iwaki	Jan-12	Cs137	206	Bq/Kg生	± 41.0	Bq/Kg生	382	Cs137	7.0	Bq/Kg生
			Cs134	176	Bq/Kg生	± 35.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Citron	Yotsukura, Iwaki	Jan-12	Cs137	139	Bq/Kg生	± 28.0	Bq/Kg生	245	Cs137	7.0	Bq/Kg生
			Cs134	106	Bq/Kg生	± 21.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Citron	Kitanosaku, Obama, Iwaki	Jan-12	Cs137	30.4	Bq/Kg生	± 8.9	Bq/Kg生	49.2	Cs137	8.0	Bq/Kg生
			Cs134	18.8	Bq/Kg生	± 5.6	Bq/Kg生		Cs134	8.0	Bq/Kg生
Citron	Miya, Uchigo	Jan-12	Cs137	91.2	Bq/Kg生	± 17.2	Bq/Kg生	133	Cs137	8.0	Bq/Kg生
			Cs134	41.5	Bq/Kg生	± 9.9	Bq/Kg生		Cs134	8.0	Bq/Kg生
Citron	Izumigaoka, Iwaki	Jan-12	Cs137	59.7	Bq/Kg生	± 12.4	Bq/Kg生	86.8	Cs137	8.0	Bq/Kg生
			Cs134	27.1	Bq/Kg生	± 6.7	Bq/Kg生		Cs134	8.0	Bq/Kg生
Citron	Ususaku, Watanabe, Iwaki	Jan-12	Cs137	54.2	Bq/Kg生	± 12.0	Bq/Kg生	105	Cs137	7.0	Bq/Kg生
			Cs134	50.5	Bq/Kg生	± 10.5	Bq/Kg生		Cs134	6.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
kabosu (citrus fruit)	Tsuzura, Uchigo, Iwaki	Jan-12	Cs137	191	Bq/Kg生	± 38.0	Bq/Kg生	347	Cs137	7.0	Bq/Kg生
			Cs134	156	Bq/Kg生	± 31.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Blueberry	Kamiyoshima, Yoshima, Iwaki	Oct-11	Cs137	28.6	Bq/Kg生	± 6.9	Bq/Kg生	58.2	Cs137	7.0	Bq/Kg生
			Cs134	29.6	Bq/Kg生	± 6.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Blueberry	Fujiwara, Joban, Iwaki	Aug-11	Cs137	49.4	Bq/Kg生	± 11.1	Bq/Kg生	97.0	Cs137	7.0	Bq/Kg生
			Cs134	47.6	Bq/Kg生	± 9.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Kiwi fruit	Shimoogawa, Iwaki	Dec-11	Cs137	14.3	Bq/Kg生	± 4.1	Bq/Kg生	20.9	Cs137	7.0	Bq/Kg生
			Cs134	6.6	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Kiwi fruit	Ogawa, Iwaki	Jan-12	Cs137	20.8	Bq/Kg生	± 5.3	Bq/Kg生	34.7	Cs137	7.0	Bq/Kg生
			Cs134	13.9	Bq/Kg生	± 3.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Kiwi fruit	Nishiogawa, Ogawa, Iwaki	Nov-11	Cs137	6.6	Bq/Kg生	± 5.9	Bq/Kg生	13.1	Cs137	8.0	Bq/Kg生
			Cs134	6.5	Bq/Kg生	± 4.7	Bq/Kg生		Cs134	8.0	Bq/Kg生
Kiwi fruit	Shimoarakawa, Taira, Iwaki	Nov-11	Cs137	11.3	Bq/Kg生	± 4.0	Bq/Kg生	23.9	Cs137	7.0	Bq/Kg生
			Cs134	12.6	Bq/Kg生	± 3.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Kiwi fruit	Miya, Uchigo, Iwaki	Jan-12	Cs137	16.7	Bq/Kg生	± 7.9	Bq/Kg生	29.7	Cs137	8.0	Bq/Kg生
			Cs134	13.0	Bq/Kg生	± 5.7	Bq/Kg生		Cs134	8.0	Bq/Kg生
Chinese citron	Negoya, kawabe, Iwaki	Jan-12	Cs137	52.0	Bq/Kg生	± 11.9	Bq/Kg生	96.3	Cs137	7.0	Bq/Kg生
			Cs134	44.3	Bq/Kg生	± 9.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mirliton	Kamiyoshima, Yoshima, Iwaki	Nov-11	Cs137	4.2	Bq/Kg生	± 2.4	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Matsutake mushroom	Yotsukura, Iwaki	Oct-11	Cs137	128	Bq/Kg生	± 26.0	Bq/Kg生	231	Cs137	7.0	Bq/Kg生
			Cs134	103	Bq/Kg生	± 21.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Maitake mushroom	Sekita, Nakoso, Iwaki	Dec-11	Cs137	3.3	Bq/Kg生	± 2.9	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Tabito, Iwaki	Oct-11	Cs137	198	Bq/Kg生	± 40.0	Bq/Kg生	374	Cs137	7.0	Bq/Kg生
			Cs134	176	Bq/Kg生	± 35.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Shimogawa, Izumi, Iwaki	Nov-11	Cs137	116	Bq/Kg生	± 24.0	Bq/Kg生	217	Cs137	7.0	Bq/Kg生
			Cs134	101	Bq/Kg生	± 20.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried shiitake mushroom	Yokoyama, Izumi, Iwaki	Oct-11	Cs137	1844	Bq/Kg生	± 211.8	Bq/Kg生	3,116	Cs137	8.0	Bq/Kg生
			Cs134	1272	Bq/Kg生	± 137.8	Bq/Kg生		Cs134	8.0	Bq/Kg生
Dried shiitake mushroom	Nagasaki, Iwaki	Dec-11	Cs137	2850	Bq/Kg生	± 570	Bq/Kg生	5,190	Cs137	7.0	Bq/Kg生
			Cs134	2340	Bq/Kg生	± 470	Bq/Kg生		Cs134	6.0	Bq/Kg生
Butterbur sprout	Kitashirado, Taira, Iwaki	Jan-12	Cs137	18.0	Bq/Kg生	± 9.7	Bq/Kg生	46.2	Cs137	7.0	Bq/Kg生
			Cs134	28.2	Bq/Kg生	± 9.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Butterbur	Tabito, Iwaki	May-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.4	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Butterbur	Ogawa, Iwaki	Unknown	Cs137	12.4	Bq/Kg生	± 5.5	Bq/Kg生	17.4	Cs137	7.0	Bq/Kg生
			Cs134	5.0	Bq/Kg生	± 4.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Udo	Tabito, Iwaki	May-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.9	Cs137	7.0	Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Red beans	Tabito, Iwaki	Nov-11	Cs137	5.6	Bq/Kg生	± 5.2	Bq/Kg生	5.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried Panda beans	Yokoyama, Izumi, Iwaki	Oct-11	Cs137	7.2	Bq/Kg生	± 7.9	Bq/Kg生	13.4	Cs137	8.0	Bq/Kg生
			Cs134	6.2	Bq/Kg生	± 6.4	Bq/Kg生		Cs134	8.0	Bq/Kg生
Wild boa meat	Ibaraki Pref.	Nov-11	Cs137	79.5	Bq/Kg生	± 12.0	Bq/Kg生	193	Cs137	8.0	Bq/Kg生
			Cs134	113	Bq/Kg生	± 19.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Wild boa meat (female)	Iwaki	Nov-11	Cs137	226	Bq/Kg生	± 45.0	Bq/Kg生	410	Cs137	7.0	Bq/Kg生
			Cs134	184	Bq/Kg生	± 37.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Wild boa meat (male)	Iwaki	Nov-11	Cs137	171	Bq/Kg生	± 34.0	Bq/Kg生	299	Cs137	7.0	Bq/Kg生
			Cs134	128	Bq/Kg生	± 26.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Wild boa meat	Kamiarakawa, Taira, Iwaki	Nov-11	Cs137	110	Bq/Kg生	± 22.0	Bq/Kg生	207	Cs137	7.0	Bq/Kg生
			Cs134	96.7	Bq/Kg生	± 19.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Wild boa meat	Iwaki	Nov-11	Cs137	190	Bq/Kg生	± 38.0	Bq/Kg生	366	Cs137	7.0	Bq/Kg生
			Cs134	176	Bq/Kg生	± 35.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Egg	Matsukusune, Joban, Iwaki	Jan-12	Cs137	3.1	Bq/Kg生	± 2.9	Bq/Kg生	7.8	Cs137	7.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried noodles	Foreign(product)	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.3	Cs137	6.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Dried seaweed noodles	Foreign(product) noodles/Soma seaweed	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.6	Cs137	6.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生
Buckwheat	Kouzashi, Aizuwakamatsu	Nov-11	Cs137	4.4	Bq/Kg生	± 3.4	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Honey	Takijiri, Izumi, Iwaki	Sep-11	Cs137	23.2	Bq/Kg生	± 6.5	Bq/Kg生	42.3	Cs137	8.0	Bq/Kg生
			Cs134	19.1	Bq/Kg生	± 4.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Honey	Kabeya, Taira, Iwaki	Oct-11	Cs137	32.1	Bq/Kg生	± 7.5	Bq/Kg生	62.5	Cs137	7.0	Bq/Kg生
			Cs134	30.4	Bq/Kg生	± 6.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Konjac	Tabito, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Salted plum	Kawabe, Iwaki	Jun-11	Cs137	122	Bq/Kg生	± 16.4	Bq/Kg生	200	Cs137	8.0	Bq/Kg生
			Cs134	78.4	Bq/Kg生	± 10.1	Bq/Kg生		Cs134	8.0	Bq/Kg生
Salted butterbur	Sekimoto, Kitaibaraki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Soy sauce	Unknown	Unknown	Cs137	9.3	Bq/Kg生	± 2.5	Bq/Kg生	15.1	Cs137	5.0	Bq/Kg生
			Cs134	5.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	3.0	Bq/Kg生
Salt	Misaki Park offshore, Iwaki	Jul-11	Cs137	12.4	Bq/Kg生	± 8.8	Bq/Kg生	30.9	Cs137	5.0	Bq/Kg生
			Cs134	18.5	Bq/Kg生	± 8.5	Bq/Kg生		Cs134	3.0	Bq/Kg生
School lunch	Joban, Iwaki	Unknown	Cs137	1.8	Bq/Kg生	± 4.7	Bq/Kg生	4.5	Cs137	8.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
School lunch	Uchigo, Iwaki	Unknown	Cs137	4.4	Bq/Kg生	± 2.3	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
School lunch	Uchigo, Iwaki	Unknown	Cs137	4.8	Bq/Kg生	± 2.5	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yogurt	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Milk	Ootsuki, Koriyama	Jan-12	Cs137	1.7	Bq/Kg生	± 1.4	Bq/Kg生	1.7	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Milk	Unknown	Unknown	Cs137	2.0	Bq/Kg生	± 1.3	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Breast milk	Onahama, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.5	Cs137	20.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 3.7	Bq/Kg生		Cs134	20.0	Bq/Kg生
Yuzu juice	Shirokajiro, Onahama, Iwaki	Jan-12	Cs137	18.2	Bq/Kg生	± 4.1	Bq/Kg生	33.9	Cs137	5.0	Bq/Kg生
			Cs134	15.7	Bq/Kg生	± 3.5	Bq/Kg生		Cs134	3.0	Bq/Kg生
Plum wine	Shimoarakawa, Taira, Iwaki	Jul-11	Cs137	22.9	Bq/Kg生	± 5.0	Bq/Kg生	41.5	Cs137	5.0	Bq/Kg生
			Cs134	18.6	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Shimoarakawa, Taira, Iwaki	Jan-12	Cs137	1.7	Bq/Kg生	± 1.5	Bq/Kg生	3.8	Cs137	5.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.4	Bq/Kg生		Cs134	3.0	Bq/Kg生

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Tap water	Minamishirado, Taira, Iwaki	Dec-11	Cs137	1.6 Bq/Kg生	± 1.4 Bq/Kg生	3.0	Cs137	5.0 Bq/Kg生		
			Cs134	1.4 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Yotsukura, Iwaki	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.5	Cs137	5.0 Bq/Kg生		
			Cs134	2.5 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Eguri, Nishiki, Iwaki	Jan-12	Cs137	1.8 Bq/Kg生	± 1.5 Bq/Kg生	3.2	Cs137	5.0 Bq/Kg生		
			Cs134	1.4 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Izumigaoka, Iwaki	Jan-12	Cs137	2.3 Bq/Kg生	± 1.5 Bq/Kg生	4.0	Cs137	5.0 Bq/Kg生		
			Cs134	1.7 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Shimoyoshima, Yoshima, Iwaki	Jan-12	Cs137	1.9 Bq/Kg生	± 1.5 Bq/Kg生	1.9	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Onahama, Iwaki	Jan-12	Cs137	2.9 Bq/Kg生	± 1.5 Bq/Kg生	5.2	Cs137	7.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 1.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Water	Miwa, Iwaki	Dec-11	Cs137	13.7 Bq/Kg生	± 3.4 Bq/Kg生	25.7	Cs137	7.0 Bq/Kg生		
			Cs134	12.0 Bq/Kg生	± 2.9 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Well water	Nihonmatsu	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.0	Cs137	5.0 Bq/Kg生		
			Cs134	2.0 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Kaminagai, Miwa, Iwaki	Jan-12	Cs137	2.2 Bq/Kg生	± 1.4 Bq/Kg生	2.2	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Yotsukura, Iwaki	Jan-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.4	Cs137	5.0 Bq/Kg生		
			Cs134	1.4 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Tono, Iwaki	Jan-12	Cs137	2.6 Bq/Kg生	± 1.5 Bq/Kg生	4.5	Cs137	5.0 Bq/Kg生		
			Cs134	1.9 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Kamioyoe, Taira, Iwaki	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.8	Cs137	5.0 Bq/Kg生		
			Cs134	1.8 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Iritono, Tono, Iwaki	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.5	Cs137	5.0 Bq/Kg生		
			Cs134	2.5 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Godo, Miwa, Iwaki	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.1	Cs137	5.0 Bq/Kg生		
			Cs134	2.1 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Shibahara, Ogawa, Iwaki	Dec-11	Cs137	2.5 Bq/Kg生	± 1.6 Bq/Kg生	3.9	Cs137	5.0 Bq/Kg生		
			Cs134	1.4 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Ground water	Miwa, Iwaki	Jan-12	Cs137	1.8 Bq/Kg生	± 1.4 Bq/Kg生	1.8	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Ground water	Osokawa, Miwa, Iwaki	Jan-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Kashima, Iwaki	Dec-11	Cs137	1.4 Bq/Kg生	± 1.4 Bq/Kg生	1.4	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Rice straw	Tamatsuyu, Izumi, Iwaki	Oct-11	Cs137	838 Bq/Kg生	± 186.0 Bq/Kg生	1,585	Cs137	7.0 Bq/Kg生		
			Cs134	747 Bq/Kg生	± 157.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Bamboo	Joban, Iwaki	Dec-11	Cs137	346 Bq/Kg生	± 71.0 Bq/Kg生	643	Cs137	7.0 Bq/Kg生		
			Cs134	297 Bq/Kg生	± 59.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Pine leaves	Chuodai, Iwaki	Nov-11	Cs137	81.6 Bq/Kg生	± 22.4 Bq/Kg生	144	Cs137	7.0 Bq/Kg生		
			Cs134	62.4 Bq/Kg生	± 17.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Pine leaves	Chuodai, Iwaki	Nov-11	Cs137	59.3 Bq/Kg生	± 18.9 Bq/Kg生	121	Cs137	7.0 Bq/Kg生		
			Cs134	62.0 Bq/Kg生	± 17.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		

