



# Radiation Measurement Results of 401 Items in February 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Kakuda, Miyagi	Oct-11	Cs137	2.7	Bq/Kg生	± 2.1	Bq/Kg生	6.2	Cs137	6.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Minamiuonuma, Nigata	Unknown	Cs137	2.6	Bq/Kg生	± 2.6	Bq/Kg生	2.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Saku, Nagano	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Chiba	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ibaraki	Oct-11	Cs137	6.5	Bq/Kg生	± 2.9	Bq/Kg生	10.9	Cs137	7.0	Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kamikawasaki, Nihonmatsu	Nov-11	Cs137	6.0	Bq/Kg生	± 2.9	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kunimi, Date	Oct-11	Cs137	69.8	Bq/Kg生	± 14.2	Bq/Kg生	123	Cs137	6.0	Bq/Kg生
			Cs134	53.9	Bq/Kg生	± 10.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Tamura, Koriyama	Nov-11	Cs137	3.3	Bq/Kg生	± 2.5	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Fukushima	Oct-11	Cs137	10.3	Bq/Kg生	± 3.0	Bq/Kg生	16.1	Cs137	6.0	Bq/Kg生
			Cs134	5.8	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Aizu	Oct-11	Cs137	4.1	Bq/Kg生	± 2.4	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shirakawa	Oct-11	Cs137	3.0	Bq/Kg生	± 2.7	Bq/Kg生	7.7	Cs137	7.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ide, Naraha, Futaba	Oct-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Naraha, Futaba	Nov-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.1	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Naraha, Futaba	Oct-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Ishikawa	Oct-11	Cs137	5.0	Bq/Kg生	± 2.7	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kawamae, Iwaki	Unknown	Cs137	3.8	Bq/Kg生	± 1.9	Bq/Kg生	3.8	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Shimookouri, Kawamae, Iwaki	Sep-11	Cs137	6.6	Bq/Kg生	± 3.1	Bq/Kg生	6.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Tajima, Misawa, Iwaki	Sep-11	Cs137	6.7	Bq/Kg生	± 3.6	Bq/Kg生	15.0	Cs137	7.0	Bq/Kg生
			Cs134	8.3	Bq/Kg生	± 3.2	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Yotsukura, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 2.9	Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Yotsukura, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.0	Bq/Kg生	5.6	Cs137	6.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Oono, Yotsukura, Iwaki	Nov-11	Cs137	4.4	Bq/Kg生	± 2.5	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ooura, Yotsukura, Iwaki	Oct-11	Cs137	4.8	Bq/Kg生	± 2.3	Bq/Kg生	7.9	Cs137	6.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Tamayama, Yotsukura, Iwaki	Oct-11	Cs137	3.6	Bq/Kg生	± 3.0	Bq/Kg生	10.8	Cs137	7.0	Bq/Kg生
			Cs134	7.2	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Tamayama, Yotsukura, Iwaki	Oct-11	Cs137	3.5	Bq/Kg生	± 2.8	Bq/Kg生	8.0	Cs137	7.0	Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kitsunozuka, Yotsukura, Iwaki	Oct-11	Cs137	4.3	Bq/Kg生	± 2.4	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kitsunozuka, Yotsukura, Iwaki	Nov-11	Cs137	9.9	Bq/Kg生	± 3.6	Bq/Kg生	18.5	Cs137	7.0	Bq/Kg生
			Cs134	8.6	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Komagome, Yotsukura, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.5	Bq/Kg生	7.0	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Takaku, Taira, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.1	Bq/Kg生	8.6	Cs137	6.0	Bq/Kg生
			Cs134	4.9	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Takaku, Taira, Iwaki	Oct-11	Cs137	2.6	Bq/Kg生	± 2.5	Bq/Kg生	5.8	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimotakaku, Taira, Iwaki	Unknown	Cs137	3.5	Bq/Kg生	± 2.8	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimotakaku, Taira, Iwaki	Unknown	Cs137	2.9	Bq/Kg生	± 2.9	Bq/Kg生	8.4	Cs137	7.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	4.2	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Hirakubo, Taira, Iwaki	Oct-11	Cs137	5.1	Bq/Kg生	± 2.3	Bq/Kg生	7.7	Cs137	6.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Hirakubo, Taira, Iwaki	Nov-11	Cs137	4.2	Bq/Kg生	± 2.7	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nakahirakubo, Taira, Iwaki	Oct-11	Cs137	3.4	Bq/Kg生	± 2.4	Bq/Kg生	3.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Fujima, Taira, Iwaki	Oct-11	Cs137	5.9	Bq/Kg生	± 3.1	Bq/Kg生	10.9	Cs137	7.0	Bq/Kg生
			Cs134	5.0	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Fujima, Taira, Iwaki	Nov-11	Cs137	9.7	Bq/Kg生	± 3.6	Bq/Kg生	14.0	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Yamaguchi, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.2	Bq/Kg生	6.9	Cs137	6.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Kamikatayose, Taira, Iwaki	Oct-11	Cs137	6.9	Bq/Kg生	± 3.0	Bq/Kg生	10.3	Cs137	7.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	7.2	Bq/Kg生	± 2.8	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimoogoe, Taira, Iwaki	Oct-11	Cs137	4.6	Bq/Kg生	± 2.7	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimoogoe, Taira, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 2.7	Bq/Kg生	8.1	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Byakko, Kouya, Uchigo	Oct-11	Cs137	15.5	Bq/Kg生	± 4.5	Bq/Kg生	26.0	Cs137	7.0	Bq/Kg生
			Cs134	10.5	Bq/Kg生	± 3.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Oyasaku, Yoshima, Iwaki	Nov-11	Cs137	4.0	Bq/Kg生	± 2.1	Bq/Kg生	4.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Izumi, Iwaki	Oct-11	Cs137	5.8	Bq/Kg生	± 2.3	Bq/Kg生	5.8	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Izumi, Iwaki	Oct-12	Cs137	3.5	Bq/Kg生	± 2.0	Bq/Kg生	6.3	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Horinouchi, Izumi, Iwaki	Oct-11	Cs137	3.4	Bq/Kg生	± 2.8	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Watanabe, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 1.9	Bq/Kg生	3.3	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Watanabe, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.0	Bq/Kg生	3.9	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Tabito, Iwaki	Oct-11	Cs137	6.5	Bq/Kg生	± 3.1	Bq/Kg生	9.4	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Tabito, Iwaki	Nov-11	Cs137	4.6	Bq/Kg生	± 2.4	Bq/Kg生	4.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nichibu, Tabito, Iwaki	Nov-11	Cs137	3.6	Bq/Kg生	± 2.5	Bq/Kg生	3.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Tono, Iwaki	Oct-11	Cs137	4.8	Bq/Kg生	± 2.4	Bq/Kg生	6.6	Cs137	6.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Tono, Iwaki	Oct-11	Cs137	3.6	Bq/Kg生	± 2.7	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Tono, Iwaki	Sep-11	Cs137	4.1	Bq/Kg生	± 2.5	Bq/Kg生	7.7	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Numabe, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.4	Bq/Kg生	3.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Numabe, Iwaki	Oct-11	Cs137	3.6	Bq/Kg生	± 2.4	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Yamada, Iwaki	Unknown	Cs137	4.5	Bq/Kg生	± 2.2	Bq/Kg生	10.2	Cs137	6.0	Bq/Kg生
			Cs134	5.7	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Kamiyada, Joban, Iwaki	Nov-11	Cs137	3.1	Bq/Kg生	± 2.4	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kamiyada, Joban, Iwaki	Nov-11	Cs137	3.1	Bq/Kg生	± 2.4	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Osamago, Joban, Iwaki	Oct-11	Cs137	4.2	Bq/Kg生	± 2.7	Bq/Kg生	7.4	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Osamago, Joban, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 2.6	Bq/Kg生	4.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Iwagaoka, Joban, Iwaki	Oct-11	Cs137	4.0	Bq/Kg生	± 2.7	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ueda, Iwaki	Sep-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.1	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nishiki, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 2.8	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Onahama, Iwaki	Unknown	Cs137	3.0	Bq/Kg生	± 2.0	Bq/Kg生	4.7	Cs137	6.0	Bq/Kg生
			Cs134	1.7	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Soeno, Iwaki	Oct-11	Cs137	5.0	Bq/Kg生	± 2.7	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Unknown	Unknown	Cs137	5.9	Bq/Kg生	± 2.8	Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Unknown	Oct-11	Cs137	4.1	Bq/Kg生	± 2.1	Bq/Kg生	6.6	Cs137	6.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Unknown	Unknown	Cs137	3.1	Bq/Kg生	± 1.9	Bq/Kg生	3.1	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Hokkaido	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Akita	Oct-11	Cs137	2.7	Bq/Kg生	± 1.8	Bq/Kg生	2.7	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Akita	Sep-11	Cs137	2.8	Bq/Kg生	± 2.4	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yamagata	Oct-11	Cs137	3.8	Bq/Kg生	± 2.4	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nanyo, Yamagata	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miyagi	Oct-11	Cs137	4.1	Bq/Kg生	± 2.7	Bq/Kg生	8.8	Cs137	7.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miyagi	Nov-11	Cs137	3.7	Bq/Kg生	± 2.0	Bq/Kg生	5.4	Cs137	6.0	Bq/Kg生
			Cs134	1.7	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Miyagi	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Furukawa, Miyagi	Unknown	Cs137	3.9	Bq/Kg生	± 2.0	Bq/Kg生	5.8	Cs137	6.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kakuda, Miyagi Pref.	Oct-11	Cs137	2.8	Bq/Kg生	± 2.4	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Watari, Watari, Miyagi	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.8	Cs137	6.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Oyama, Oushu, Iwate	Sep-11	Cs137	5.0	Bq/Kg生	± 2.7	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Sekimoto, Kitaibaraki	Nov-11	Cs137	2.8	Bq/Kg生	± 2.4	Bq/Kg生	6.2	Cs137	7.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kitaibaraki	Dec-11	Cs137	3.1	Bq/Kg生	± 2.0	Bq/Kg生	5.9	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Sekimoto, Kitaibaraki	Nov-11	Cs137	2.4	Bq/Kg生	± 2.0	Bq/Kg生	5.5	Cs137	6.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tsukuba, Kitaibaraki	Oct-11	Cs137	2.5	Bq/Kg生	± 1.9	Bq/Kg生	2.5	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Ibaraki	Mar-11	Cs137	2.8	Bq/Kg生	± 2.6	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tochigi	Unknown	Cs137	2.5	Bq/Kg生	± 2.5	Bq/Kg生	5.6	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tochigi	Oct-11	Cs137	3.9	Bq/Kg生	± 2.0	Bq/Kg生	5.8	Cs137	6.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Fukushima	Unknown	Cs137	2.6	Bq/Kg生	± 2.2	Bq/Kg生	2.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
			Cs137	Cs134	±	±		Cs137	Cs134		
Rice	Fukushima	Unknown	Cs137	3.2	Bq/Kg生	± 2.6	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Iino, Fukushima	Unknown	Cs137	16.1	Bq/Kg生	± 4.5	Bq/Kg生	25.0	Cs137	7.0	Bq/Kg生
			Cs134	8.9	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yamada, Fukushima	Nov-11	Cs137	27.1	Bq/Kg生	± 6.7	Bq/Kg生	50.4	Cs137	7.0	Bq/Kg生
			Cs134	23.3	Bq/Kg生	± 5.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tamura, Koriyama	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.0	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Obama, Nihonmatsu	Oct-11	Cs137	4.2	Bq/Kg生	± 2.7	Bq/Kg生	9.7	Cs137	7.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shibukawa, Nihonmatsu	Oct-11	Cs137	2.8	Bq/Kg生	± 2.0	Bq/Kg生	5.8	Cs137	6.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Okugawa, Nishiaizu, Yama	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Inawashiro, Yama	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.3	Cs137	6.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Toyokawa, Kitakata	Oct-11	Cs137	3.2	Bq/Kg生	± 2.4	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Aizuwakamatsu	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Aizu	Oct-11	Cs137	2.7	Bq/Kg生	± 2.4	Bq/Kg生	2.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Aizu	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Aizubange	Oct-11	Cs137	2.1	Bq/Kg生	± 1.9	Bq/Kg生	2.1	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Shimoyamada, Sukagawa	Oct-11	Cs137	2.9	Bq/Kg生	± 2.4	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Naraha, Futaba	Oct-10	Cs137	2.7	Bq/Kg生	± 1.9	Bq/Kg生	2.7	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tomioka, Futaba	Oct-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tomioka, Futaba	Oct-10	Cs137	2.4	Bq/Kg生	± 1.9	Bq/Kg生	4.9	Cs137	6.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Ono, Tamura	Oct-11	Cs137	3.5	Bq/Kg生	± 2.6	Bq/Kg生	3.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ono, Tamura	Nov-11	Cs137	3.3	Bq/Kg生	± 2.0	Bq/Kg生	5.7	Cs137	6.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tamanoi, Ootama, Adachi	Nov-11	Cs137	3.3	Bq/Kg生	± 2.4	Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Iwase	Oct-11	Cs137	3.0	Bq/Kg生	± 2.0	Bq/Kg生	5.1	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Iwaki	Unknown	Cs137	3.3	Bq/Kg生	± 1.9	Bq/Kg生	5.2	Cs137	6.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Iwaki	Oct-11	Cs137	2.1	Bq/Kg生	± 2.0	Bq/Kg生	2.1	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Takabe, Kawamae, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 1.9	Bq/Kg生	6.3	Cs137	6.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Miwa, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.5	Bq/Kg生	5.4	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miwa, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.0	Bq/Kg生	5.1	Cs137	6.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Miwa, Iwaki	Oct-11	Cs137	5.4	Bq/Kg生	± 2.6	Bq/Kg生	7.6	Cs137	7.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miwa, Iwaki	Nov-11	Cs137	3.2	Bq/Kg生	± 2.4	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miwa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Saiso, Miwa, Iwaki	Oct-11	Cs137	2.2	Bq/Kg生	± 1.8	Bq/Kg生	2.2	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kaminagai, Miwa, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.4	Bq/Kg生	3.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.1	Bq/Kg生	6.6	Cs137	6.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.1	Bq/Kg生	7.5	Cs137	6.0	Bq/Kg生
			Cs134	4.2	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.2	Cs137	6.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	3.5	Bq/Kg生	± 2.4	Bq/Kg生	7.4	Cs137	7.0	Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Unknown	Cs137	2.6	Bq/Kg生	± 1.9	Bq/Kg生	5.4	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tamayama, Yotsukura, Iwaki	Nov-11	Cs137	4.1	Bq/Kg生	± 2.0	Bq/Kg生	6.1	Cs137	6.0	Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tamayama, Yotsukura, Iwaki	Nov-11	Cs137	4.1	Bq/Kg生	± 2.0	Bq/Kg生	6.1	Cs137	6.0	Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yaguki, Yotsukura, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 2.6	Bq/Kg生	6.9	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Oomori, Yotsukura, Iwaki	Oct-11	Cs137	5.7	Bq/Kg生	± 2.7	Bq/Kg生	5.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nagatomo, Yotsukura, Iwaki	Oct-11	Cs137	2.4	Bq/Kg生	± 2.1	Bq/Kg生	4.5	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Nishiogawa, Ogawa, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.4	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shioda, Ogawa, Iwaki	Unknown	Cs137	4.7	Bq/Kg生	± 2.4	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ogawa, Iwaki	Nov-11	Cs137	4.8	Bq/Kg生	± 2.5	Bq/Kg生	4.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ogawa, Iwaki	Oct-11	Cs137	4.7	Bq/Kg生	± 2.4	Bq/Kg生	4.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Taira, Iwaki	Nov-11	Cs137	3.2	Bq/Kg生	± 2.0	Bq/Kg生	5.8	Cs137	6.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Taira, Iwaki	2011	Cs137	2.4	Bq/Kg生	± 2.0	Bq/Kg生	4.5	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kabeya, Taira, Iwaki	2011	Cs137	2.6	Bq/Kg生	± 2.0	Bq/Kg生	4.7	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Rice	Natsui, Taira, Iwaki	Sep-11	Cs137	6.0 Bq/Kg生	± 2.8 Bq/Kg生	6.0	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kusano, Taira, Iwaki	Sep-11	Cs137	2.5 Bq/Kg生	± 2.0 Bq/Kg生	4.7	Cs137	6.0 Bq/Kg生		
			Cs134	2.2 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	5.4 Bq/Kg生	± 2.8 Bq/Kg生	10.1	Cs137	7.0 Bq/Kg生		
			Cs134	4.7 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Mamikabeya, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.8	Cs137	6.0 Bq/Kg生		
			Cs134	2.8 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Shimokabeya, Taira, Iwaki	Nov-11	Cs137	3.5 Bq/Kg生	± 2.7 Bq/Kg生	6.1	Cs137	7.0 Bq/Kg生		
			Cs134	2.6 Bq/Kg生	± 2.4 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Nakakabeya, Taira, Iwaki	Oct-10	Cs137	2.7 Bq/Kg生	± 1.9 Bq/Kg生	4.4	Cs137	6.0 Bq/Kg生		
			Cs134	1.7 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Nakakabeya, Taira, Iwaki	Oct-11	Cs137	2.5 Bq/Kg生	± 2.1 Bq/Kg生	4.5	Cs137	6.0 Bq/Kg生		
			Cs134	2.0 Bq/Kg生	± 1.8 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Akai, Taira, Iwaki	Oct-11	Cs137	3.0 Bq/Kg生	± 2.4 Bq/Kg生	5.3	Cs137	7.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Akai, Taira, Iwaki	Nov-11	Cs137	3.8 Bq/Kg生	± 2.4 Bq/Kg生	7.0	Cs137	7.0 Bq/Kg生		
			Cs134	3.2 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Akai, Taira, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生		
			Cs134	3.1 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Akai, Taira, Iwaki	Oct-11	Cs137	3.1 Bq/Kg生	± 2.3 Bq/Kg生	5.2	Cs137	7.0 Bq/Kg生		
			Cs134	2.1 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Akai, Taira, Iwaki	Dec-11	Cs137	4.1 Bq/Kg生	± 2.1 Bq/Kg生	6.3	Cs137	6.0 Bq/Kg生		
			Cs134	2.2 Bq/Kg生	± 1.8 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Kamiarakawa, Taira, Iwaki	Oct-11	Cs137	5.0 Bq/Kg生	± 2.7 Bq/Kg生	8.0	Cs137	7.0 Bq/Kg生		
			Cs134	3.0 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Kamiarakawa, Taira, Iwaki	Oct-10	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.8	Cs137	7.0 Bq/Kg生		
			Cs134	2.8 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Takakuyamaguchi, Taira, Iwaki	Nov-11	Cs137	3.4 Bq/Kg生	± 2.0 Bq/Kg生	5.7	Cs137	6.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 1.7 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Takaku, Taira, Iwaki	Oct-11	Cs137	3.6 Bq/Kg生	± 2.0 Bq/Kg生	7.3	Cs137	6.0 Bq/Kg生		
			Cs134	3.7 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	3.1 Bq/Kg生	± 2.6 Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	3.7 Bq/Kg生	± 2.7 Bq/Kg生	6.7	Cs137	7.0 Bq/Kg生		
			Cs134	3.0 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	3.1 Bq/Kg生	± 2.6 Bq/Kg生	6.4	Cs137	7.0 Bq/Kg生		
			Cs134	3.3 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Shimotakaku, Taira, Iwaki	Nov-11	Cs137	2.7 Bq/Kg生	± 2.5 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生		
			Cs134	4.2 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	4.8 Bq/Kg生	± 2.5 Bq/Kg生	8.9	Cs137	7.0 Bq/Kg生		
			Cs134	4.1 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	4.8 Bq/Kg生	± 2.5 Bq/Kg生	7.1	Cs137	7.0 Bq/Kg生		
			Cs134	2.3 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Fujima, Taira, Iwaki	Oct-11	Cs137	3.6 Bq/Kg生	± 2.5 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生		
			Cs134	3.3 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生		
Rice	Fujima, Taira, Iwaki	Oct-11	Cs137	6.1 Bq/Kg生	± 3.0 Bq/Kg生	6.1	Cs137	7.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生		

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Yamaguchi, Taira, Iwaki	Oct-11	Cs137	2.0	Bq/Kg生	± 2.0	Bq/Kg生	4.9	Cs137	6.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yamaguchi, Taira, Iwaki	Oct-11	Cs137	2.8	Bq/Kg生	± 1.9	Bq/Kg生	4.7	Cs137	6.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yamaguchi, Taira, Iwaki	Oct-11	Cs137	2.0	Bq/Kg生	± 2.0	Bq/Kg生	4.9	Cs137	6.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Manome, Taira, Iwaki	Oct-11	Cs137	5.9	Bq/Kg生	± 2.3	Bq/Kg生	7.9	Cs137	6.0	Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Numanouchi, Taira, Iwaki	Nov-12	Cs137	3.1	Bq/Kg生	± 2.8	Bq/Kg生	7.4	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	8.8	Bq/Kg生	± 3.7	Bq/Kg生	12.6	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.4	Bq/Kg生	3.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.3	Bq/Kg生	5.4	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Haragoya, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.8	Cs137	6.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yoshima, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.5	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kamiyoshima, Yoshima, Iwaki	Oct-11	Cs137	2.2	Bq/Kg生	± 2.0	Bq/Kg生	4.4	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kubo, Yoshima, Iwaki	Oct-11	Cs137	6.7	Bq/Kg生	± 2.7	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Imaniida, Yoshima, Iwaki	Oct-11	Cs137	3.4	Bq/Kg生	± 2.5	Bq/Kg生	3.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yoshima, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.3	Bq/Kg生	3.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Takakura, Iwaki	Oct-11	Cs137	6.4	Bq/Kg生	± 3.0	Bq/Kg生	12.8	Cs137	7.0	Bq/Kg生
			Cs134	6.4	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Soeno, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.1	Bq/Kg生	5.3	Cs137	6.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Numabe, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 1.8	Bq/Kg生	3.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Numabe, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 1.8	Bq/Kg生	3.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kawabe, Iwaki	Oct-11	Cs137	5.1	Bq/Kg生	± 2.6	Bq/Kg生	8.1	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ookubo, Kawabe, Iwaki	Oct-11	Cs137	4.1	Bq/Kg生	± 2.1	Bq/Kg生	7.4	Cs137	6.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Byakko, Koya, Iwaki	Oct-11	Cs137	12.2	Bq/Kg生	± 3.5	Bq/Kg生	12.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tabito, Iwaki	Sep-11	Cs137	3.2	Bq/Kg生	± 2.5	Bq/Kg生	5.8	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kuroda, Tabito, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 2.4	Bq/Kg生	4.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tabito, Iwaki	Nov-11	Cs137	2.8	Bq/Kg生	± 2.7	Bq/Kg生	6.0	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Kashima, Iwaki	Nov-11	Cs137	3.9	Bq/Kg生	± 2.4	Bq/Kg生	6.4	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Komoda, Kashima, Iwaki	Oct-11	Cs137	4.3	Bq/Kg生	± 2.6	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Funado, Kashima, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.6	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	2.7	Bq/Kg生	± 2.0	Bq/Kg生	5.6	Cs137	6.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.1	Bq/Kg生	5.1	Cs137	6.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Ryugasawa, Joban, Iwaki	Oct-11	Cs137	3.4	Bq/Kg生	± 1.9	Bq/Kg生	3.4	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kamiyada, Joban, Iwaki	Nov-11	Cs137	4.5	Bq/Kg生	± 2.2	Bq/Kg生	7.4	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Iwagaoka, Joban, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.7	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Onahama, Iwaki	Oct-11	Cs137	2.8	Bq/Kg生	± 2.1	Bq/Kg生	6.0	Cs137	6.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Izumi, Iwaki	Oct-11	Cs137	2.5	Bq/Kg生	± 2.0	Bq/Kg生	2.5	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Izumi, Iwaki	Oct-11	Cs137	3.6	Bq/Kg生	± 2.3	Bq/Kg生	3.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tamatsuyu, Izumi, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.0	Cs137	6.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Honya, Izumi, Iwaki	Nov-11	Cs137	9.6	Bq/Kg生	± 3.0	Bq/Kg生	19.4	Cs137	6.0	Bq/Kg生
			Cs134	9.8	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Honya, Izumi, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.2	Cs137	6.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tono, Iwaki	Oct-11	Cs137	4.3	Bq/Kg生	± 2.2	Bq/Kg生	4.3	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tono, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.6	Bq/Kg生	3.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tono, Iwaki	Unknown	Cs137	5.3	Bq/Kg生	± 2.7	Bq/Kg生	5.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tono, Iwaki	Oct-11	Cs137	5.0	Bq/Kg生	± 2.5	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tono, Iwaki	Sep-11	Cs137	2.9	Bq/Kg生	± 2.3	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tono, Iwaki	Oct-11	Cs137	2.5	Bq/Kg生	± 2.0	Bq/Kg生	4.3	Cs137	6.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	3.5	Bq/Kg生	± 2.0	Bq/Kg生	5.7	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Watanabe, Iwaki	Oct-11	Cs137	4.8	Bq/Kg生	± 2.4	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Watanabe, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.2	Bq/Kg生	6.5	Cs137	6.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Watanabe, Iwaki	Oct-11	Cs137	2.0	Bq/Kg生	± 1.8	Bq/Kg生	4.0	Cs137	6.0	Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Watanabe, Iwaki	Oct-11	Cs137	9.6	Bq/Kg生	± 2.9	Bq/Kg生	14.3	Cs137	6.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Kamikamado, Watanabe, Iwaki	Sep-11	Cs137	3.2	Bq/Kg生	± 2.0	Bq/Kg生	3.2	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yamada, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 1.7	Bq/Kg生	3.1	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yamada, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nishiki, Iwaki	Nov-11	Cs137	3.0	Bq/Kg生	± 2.5	Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nishiki, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3	Cs137	6.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tomitsu, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Unknown	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.4	Cs137	6.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.6	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Unknown	Unknown	Cs137	3.3	Bq/Kg生	± 2.3	Bq/Kg生	5.4	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Unknown	Oct-11	Cs137	3.6	Bq/Kg生	± 2.6	Bq/Kg生	6.2	Cs137	6.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Unknown	Oct-11	Cs137	2.9	Bq/Kg生	± 1.8	Bq/Kg生	4.6	Cs137	6.0	Bq/Kg生
			Cs134	1.7	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Unknown	Unknown	Cs137	3.5	Bq/Kg生	± 2.6	Bq/Kg生	5.7	Cs137	7.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Washed water of rice	Miyagi	Unknown	Cs137	2.3	Bq/Kg生	± 1.4	Bq/Kg生	2.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (brown rice)	Toyama	Unknown	Cs137	3.1	Bq/Kg生	± 2.5	Bq/Kg生	3.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (brown rice)	Nishigo, Joban, Iwaki	Oct-11	Cs137	7.7	Bq/Kg生	± 3.2	Bq/Kg生	11.0	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (brown rice)	Kunimi, Date	Oct-11	Cs137	143.0	Bq/Kg生	± 29.0	Bq/Kg生	254	Cs137	6.0	Bq/Kg生
			Cs134	111.0	Bq/Kg生	± 22.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice (brown rice)	Yaguki, Yotsukura, Iwaki	Oct-11	Cs137	9.9	Bq/Kg生	± 3.2	Bq/Kg生	12.9	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Nagano	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Iwase	Oct-11	Cs137	2.8	Bq/Kg生	± 1.9	Bq/Kg生	2.8	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice (polished rice)	Miwa, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.5	Bq/Kg生	3.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Oomori, Yotsukura, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.5	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Shimoogoe, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.0	Bq/Kg生	6.4	Cs137	6.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Glutinous rice (polished rice)	Tabito, Iwaki	Sep-11	Cs137	4.2	Bq/Kg生	± 2.7	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Nakakamado, Watanabe, Iwaki	Sep-11	Cs137	5.6	Bq/Kg生	± 2.9	Bq/Kg生	8.1	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Nishiki, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 1.9	Bq/Kg生	5.3	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice (polished rice)	Ogawa, Iwaki	Oct-11	Cs137	2.8	Bq/Kg生	± 2.6	Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Akai, taira, Iwaki	Oct-11	Cs137	7.7	Bq/Kg生	± 3.2	Bq/Kg生	15.7	Cs137	7.0	Bq/Kg生
			Cs134	8.0	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (polished rice)	Nishigo, Joban, Iwaki	Oct-11	Cs137	5.0	Bq/Kg生	± 2.8	Bq/Kg生	9.5	Cs137	7.0	Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Onahama, Iwaki	Feb-12	Cs137	7.4	Bq/Kg生	± 7.0	Bq/Kg生	15.1	Cs137	7.0	Bq/Kg生
			Cs134	7.7	Bq/Kg生	± 5.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Hanabatake, Onahama, Iwaki	Feb-12	Cs137	10.4	Bq/Kg生	± 5.5	Bq/Kg生	10.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Onahama, Iwaki	Sep-11	Cs137	56.0	Bq/Kg生	± 13.7	Bq/Kg生	97.8	Cs137	7.0	Bq/Kg生
			Cs134	41.8	Bq/Kg生	± 10.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Whole rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.7	Bq/Kg生	5.3	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Tamai, Ootama, Adachi	Aug-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kamikawasaki, Nihonmatsu	Jun-11	Cs137	7.7	Bq/Kg生	± 3.1	Bq/Kg生	7.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Yabuki, Nishishirakawa	Aug-11	Cs137	4.4	Bq/Kg生	± 2.5	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimotakaku, Taira, Iwaki	Jul-11	Cs137	15.3	Bq/Kg生	± 4.3	Bq/Kg生	21.5	Cs137	6.0	Bq/Kg生
			Cs134	6.2	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	4.0	Bq/Kg生
Potato	Shimotakaku, Taira, Iwaki	Jun-11	Cs137	7.8	Bq/Kg生	± 3.3	Bq/Kg生	11.9	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kamioyama, Ogawa, Iwaki	Jun-11	Cs137	4.7	Bq/Kg生	± 3.2	Bq/Kg生	4.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kaminagai, Miwa, Iwaki	Aug-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Hosoya, Yotsukura, Iwaki	Jun-11	Cs137	8.6	Bq/Kg生	± 3.4	Bq/Kg生	15.7	Cs137	7.0	Bq/Kg生
			Cs134	7.1	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kabeya, Taira, Iwaki	Aug-11	Cs137	4.6	Bq/Kg生	± 3.3	Bq/Kg生	9.8	Cs137	7.0	Bq/Kg生
			Cs134	5.2	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kamikabeya, Taira, Iwaki	Jul-11	Cs137	6.7	Bq/Kg生	± 3.0	Bq/Kg生	10.2	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimokabeya, Taira, Iwaki	Oct-11	Cs137	4.0	Bq/Kg生	± 2.3	Bq/Kg生	6.4	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimokabeya, Taira, Iwaki	Aug-11	Cs137	3.2	Bq/Kg生	± 2.5	Bq/Kg生	7.6	Cs137	7.0	Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimoyamaguchi, Taira, Iwaki	Jun-11	Cs137	3.2	Bq/Kg生	± 3.0	Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Potato	Fujima, Taira, Iwaki	Jul-11	Cs137	28.8	Bq/Kg生	± 7.2	Bq/Kg生	47.3	Cs137	7.0	Bq/Kg生
			Cs134	18.5	Bq/Kg生	± 4.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Fujima, Taira, Iwaki	Jul-11	Cs137	5.6	Bq/Kg生	± 3.4	Bq/Kg生	10.7	Cs137	7.0	Bq/Kg生
			Cs134	5.1	Bq/Kg生	± 3.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Manome, Taira, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.9	Cs137	7.0	Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Yoshima, Iwaki	Jun-11	Cs137	3.5	Bq/Kg生	± 3.2	Bq/Kg生	3.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimoyoshima, Yoshima, Iwaki	Jul-11	Cs137	4.5	Bq/Kg生	± 2.5	Bq/Kg生	4.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimoyoshima, Yoshima, Iwaki	Jul-11	Cs137	3.7	Bq/Kg生	± 3.3	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimoyoshima, Yoshima, Iwaki	Jul-11	Cs137	9.5	Bq/Kg生	± 4.0	Bq/Kg生	9.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Nichibu, Tabito, Iwaki	Aug-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Sadanishi, Onahama, Iwaki	Jul-11	Cs137	5.8	Bq/Kg生	± 3.4	Bq/Kg生	5.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimokajiro, Onahama, Iwaki	Jun-11	Cs137	3.6	Bq/Kg生	± 3.2	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Takijiri, Izumi, Iwaki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.0	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kubota, Nakoso, Iwaki	Sep-11	Cs137	4.1	Bq/Kg生	± 2.6	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Hosoya, Yotsukura, Iwaki	Jul-11	Cs137	12.0	Bq/Kg生	± 5.7	Bq/Kg生	18.3	Cs137	7.0	Bq/Kg生
			Cs134	6.3	Bq/Kg生	± 4.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Soeno, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.4	Cs137	7.0	Bq/Kg生
			Cs134	5.4	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Ogawa, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yam	Toyoma, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.4	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yam	Takakura, Yumoto, Iwaki	Dec-11	Cs137	16.0	Bq/Kg生	± 7.5	Bq/Kg生	22.1	Cs137	8.0	Bq/Kg生
			Cs134	6.1	Bq/Kg生	± 5.1	Bq/Kg生		Cs134	8.0	Bq/Kg生
Yam	Minamishirado, Taira, Iwaki	Nov-11	Cs137	4.6	Bq/Kg生	± 6.3	Bq/Kg生	10.5	Cs137	8.0	Bq/Kg生
			Cs134	5.9	Bq/Kg生	± 4.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Sweet potato	Kitsunozuka, Yotsukura, Iwaki	Nov-11	Cs137	8.0	Bq/Kg生	± 4.1	Bq/Kg生	12.5	Cs137	7.0	Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Taro	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Taro	Yoshima, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Taro (with peel)	Fujima, Taira, Iwaki	Nov-11	Cs137	7.3	Bq/Kg生	± 4.1	Bq/Kg生	11.9	Cs137	7.0	Bq/Kg生
			Cs134	4.6	Bq/Kg生	± 3.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shioda, Ogawa, Iwaki	Unknown	Cs137	4.2	Bq/Kg生	± 3.0	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shimokabeya, Taira, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Japanese white radish	Shimotakaku, Taira, Iwaki	Feb-12	Cs137	4.4	Bq/Kg生	± 4.2	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Manome, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.1	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Akai, Taira, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.8	Cs137	7.0	Bq/Kg生
			Cs134	4.8	Bq/Kg生	± 3.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Sadanishi, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Tamagawa, Onahama, Iwaki	Feb-12	Cs137	4.0	Bq/Kg生	± 3.3	Bq/Kg生	4.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Oohara, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Tsurumaki, Takakura, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.9	Cs137	7.0	Bq/Kg生
			Cs134	4.9	Bq/Kg生	± 3.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried radish	Ogawa, Iwaki	Jan-12	Cs137	83.1	Bq/Kg生	± 27.9	Bq/Kg生	175	Cs137	8.0	Bq/Kg生
			Cs134	92.0	Bq/Kg生	± 21.1	Bq/Kg生		Cs134	8.0	Bq/Kg生
Dried radish	Moegidai, Iwaki	Feb-12	Cs137	226.0	Bq/Kg生	± 55.0	Bq/Kg生	400	Cs137	7.0	Bq/Kg生
			Cs134	174.0	Bq/Kg生	± 42.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Freeze-dried Japanese white radish	Hosoya, Yotsukura, Iwaki	Jan-12	Cs137	26.4	Bq/Kg生	± 8.9	Bq/Kg生	41.8	Cs137	7.0	Bq/Kg生
			Cs134	15.4	Bq/Kg生	± 6.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Reconstituted water of dried radish	Hosoya, Yotsukura, Iwaki	Jan-12	Cs137	2.5	Bq/Kg生	± 1.5	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	1.6	Bq/Kg生	± 1.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Carrot	Watanabe, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Pumpkin	Akai, Taira, Iwaki	Oct-11	Cs137	6.6	Bq/Kg生	± 3.3	Bq/Kg生	11.0	Cs137	7.0	Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Yotsukura, Iwaki	Aug-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Fujima, Taira, Iwaki	Jun-11	Cs137	5.1	Bq/Kg生	± 2.6	Bq/Kg生	8.9	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Kamikabeya, Taira, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Toyoma, Taira, Iwaki	Jun-11	Cs137	5.4	Bq/Kg生	± 3.2	Bq/Kg生	9.4	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Akai, Taira, Iwaki	Jul-11	Cs137	3.4	Bq/Kg生	± 2.5	Bq/Kg生	7.5	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Kubota, Nakoso, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Yoshima, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Kitsunozuka, Yotsukura, Iwaki	Feb-12	Cs137	2.7	Bq/Kg生	± 2.6	Bq/Kg生	6.4	Cs137	7.0	Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Youkoudai, Iwaki	Jan-11	Cs137	4.3	Bq/Kg生	± 3.1	Bq/Kg生	8.0	Cs137	7.0	Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green onion	Kamiyada, Joban, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 3.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Ishikawa, Ishikawa	Dec-11	Cs137	3.5	Bq/Kg生	± 2.4	Bq/Kg生	3.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Chinese cabbage	Kamiogawa, Ogawa, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	9.3	Cs137	7.0	Bq/Kg生
			Cs134	9.3	Bq/Kg生	± 5.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Akai, Taira, Iwaki	Feb-12	Cs137	5.6	Bq/Kg生	± 5.3	Bq/Kg生	5.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Minamishirado, Taira, Iwaki	Feb-12	Cs137	5.2	Bq/Kg生	± 2.9	Bq/Kg生	5.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kabeya, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimokabeya, Taira, Iwaki	Jan-12	Cs137	8.6	Bq/Kg生	± 4.7	Bq/Kg生	8.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimokabeya, Taira, Iwaki	Feb-12	Cs137	6.0	Bq/Kg生	± 4.3	Bq/Kg生	6.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimokabeya, Taira, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Shimotakaku, Taira, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Fujima, Taira, Iwaki	Feb-12	Cs137	7.5	Bq/Kg生	± 4.9	Bq/Kg生	15.5	Cs137	7.0	Bq/Kg生
			Cs134	8.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Tsurumaki, Takakura, Iwaki	Feb-12	Cs137	10.7	Bq/Kg生	± 5.9	Bq/Kg生	18.5	Cs137	7.0	Bq/Kg生
			Cs134	7.8	Bq/Kg生	± 4.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Tamagawa, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Watanabe, Iwaki	Feb-12	Cs137	6.4	Bq/Kg生	± 4.6	Bq/Kg生	7.0	Cs137	7.0	Bq/Kg生
			Cs134	5.1	Bq/Kg生	± 4.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Tabito, Iwaki	Feb-12	Cs137	6.3	Bq/Kg生	± 4.8	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kamiyada, Joban, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	7.5	Cs137	7.0	Bq/Kg生
			Cs134	7.5	Bq/Kg生	± 4.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Kimigatsuka, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.4	Cs137	7.0	Bq/Kg生
			Cs134	6.4	Bq/Kg生	± 4.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese cabbage	Komoda, Kashima, Iwaki	Dec-11	Cs137	9.0	Bq/Kg生	± 6.1	Bq/Kg生	9.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Oohara, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Kamikamado, Watanabe, Iwaki	Jan-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	7.2	Cs137	7.0	Bq/Kg生
			Cs134	7.2	Bq/Kg生	± 5.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Qing-geng-cai	Hirakubo, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	5.0	Bq/Kg生	± 4.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Canola flower	Minamitomioka, Onahama, Iwaki	Feb-12	Cs137	10.3	Bq/Kg生	± 6.2	Bq/Kg生	18.4	Cs137	7.0	Bq/Kg生
			Cs134	8.1	Bq/Kg生	± 5.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Tomato	Nakashio, Taira, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Pickled scallion	Kitsunozuka, Yotsukura, Iwaki	Feb-12	Cs137	5.0	Bq/Kg生	± 3.2	Bq/Kg生	9.6	Cs137	7.0	Bq/Kg生
			Cs134	4.6	Bq/Kg生	± 2.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Broccoli	Kamikajiro, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green pepper	Hosoya, Yotsukura, Iwaki	Dec-11	Cs137	3.5	Bq/Kg生	± 5.1	Bq/Kg生	4.7	Cs137	8.0	Bq/Kg生
			Cs134	1.2	Bq/Kg生	± 3.9	Bq/Kg生		Cs134	8.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
			Cs137	Bq/Kg生	±	Bq/Kg生		Cs137	Bq/Kg生	Cs134
Shiitake mushroom	Shimoyamaguchi, Taira, Iwaki	Oct-11	Cs137	2030	±	410	3,650	Cs137	7.0	Bq/Kg生
			Cs134	1620	±	320		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Nakahata, Misawa, Iwaki	Mar-11	Cs137	2280	±	460	4,240	Cs137	7.0	Bq/Kg生
			Cs134	1960	±	390		Cs134	6.0	Bq/Kg生
Citron	Kakuda, Miyagi	Dec-11	Cs137	39.3	±	9.0	73.7	Cs137	7.0	Bq/Kg生
			Cs134	34.4	±	7.5		Cs134	6.0	Bq/Kg生
Citron	Hassen, Yumoto, Iwaki	Oct-11	Cs137	55.5	±	12.5	101	Cs137	7.0	Bq/Kg生
			Cs134	45.5	±	9.9		Cs134	6.0	Bq/Kg生
Citron	Toyoma, Taira, Iwaki	Dec-11	Cs137	57.1	±	12.4	98.8	Cs137	7.0	Bq/Kg生
			Cs134	41.7	±	8.9		Cs134	6.0	Bq/Kg生
Sumo mandarin	Shimoyoshima, Yoshima, Iwaki	Feb-12	Cs137	32.1	±	7.7	65.0	Cs137	7.0	Bq/Kg生
			Cs134	32.9	±	7.2		Cs134	6.0	Bq/Kg生
Sumo mandarin	Nakoso, Iwaki	Feb-12	Cs137	90.4	±	18.5	158	Cs137	8.0	Bq/Kg生
			Cs134	67.8	±	12.1		Cs134	8.0	Bq/Kg生
Mandarin orange	Nakoso, Iwaki	Feb-12	Cs137	49.9	±	12.4	87.7	Cs137	8.0	Bq/Kg生
			Cs134	37.8	±	8.4		Cs134	8.0	Bq/Kg生
Mandarin orange	Satogaoka, Iwaki	Dec-11	Cs137	58.9	±	12.6	97.9	Cs137	8.0	Bq/Kg生
			Cs134	39.0	±	8.2		Cs134	8.0	Bq/Kg生
Navel orange (with peel)	Nakoso, Iwaki	Feb-12	Cs137	78.3	±	16.0	127	Cs137	8.0	Bq/Kg生
			Cs134	48.5	±	9.7		Cs134	8.0	Bq/Kg生
Kiwi	Kaminodera, Fukushima	Dec-11	Cs137	45.9	±	10.4	82.2	Cs137	7.0	Bq/Kg生
			Cs134	36.3	±	8.0		Cs134	6.0	Bq/Kg生
Kiwi	Koya, Uchigo, Iwaki	Oct-11	Cs137	31.8	±	9.2	54.7	Cs137	8.0	Bq/Kg生
			Cs134	22.9	±	6.7		Cs134	8.0	Bq/Kg生
Kiwi	Kamiyamaguchi, Taira, Iwaki	Nov-11	Cs137	30.3	±	7.5	58.9	Cs137	7.0	Bq/Kg生
			Cs134	28.6	±	6.5		Cs134	6.0	Bq/Kg生
Persimmon	Shirokajiro, Onahama, Iwaki	Oct-11	Cs137	7.6	±	3.3	20.0	Cs137	7.0	Bq/Kg生
			Cs134	12.4	±	3.4		Cs134	6.0	Bq/Kg生
Chinese quince	Shioki, Yotsukura, Iwaki	Nov-11	Cs137	85.4	±	18.5	153	Cs137	7.0	Bq/Kg生
			Cs134	67.6	±	14.1		Cs134	6.0	Bq/Kg生
Smelt	Tanakura, Higashishirakawa	Feb-12	Cs137	8.1	±	7.9	19.4	Cs137	5.0	Bq/Kg生
			Cs134	11.3	±	6.4		Cs134	3.0	Bq/Kg生
Greeneye	Ehime	Unknown	Cs137	7.2	±	4.0	7.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	±	0.0		Cs134	6.0	Bq/Kg生
Shrimp	Ehime	Feb-12	Cs137	0.0	±	0.0	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	±	8.7		Cs134	6.0	Bq/Kg生
Raw seaweed	Toyoma, Taira, Iwaki	Feb-12	Cs137	3.9	±	3.4	3.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	±	0.0		Cs134	6.0	Bq/Kg生
Seaweed	Onahama, Iwaki	Feb-12	Cs137	8.9	±	6.5	11.6	Cs137	8.0	Bq/Kg生
			Cs134	2.7	±	3.8		Cs134	8.0	Bq/Kg生
Egg (Silky Fowl)	Shimohirakubo, Taira, Iwaki	Jan-12	Cs137	3.3	±	2.6	7.1	Cs137	7.0	Bq/Kg生
			Cs134	3.8	±	2.3		Cs134	6.0	Bq/Kg生
Egg	Isumi, Chiba	Unknown	Cs137	0.0	±	0.0	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	±	0.0		Cs134	6.0	Bq/Kg生
Beef(thigh)	Fukushima	Unknown	Cs137	0.0	±	0.0	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	±	0.0		Cs134	6.0	Bq/Kg生
Pork(loin)	Japan	Unknown	Cs137	0.0	±	0.0	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	±	0.0		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Dried zenmai	Tono, Iwaki	Jun-11	Cs137	1122	Bq/Kg生	± 138	Bq/Kg生	1,808	Cs137	8.0	Bq/Kg生
			Cs134	686	Bq/Kg生	± 81.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Ginkgo	Oshodan, Shirakawa	Oct-11	Cs137	44.6	Bq/Kg生	± 10.5	Bq/Kg生	61.8	Cs137	8.0	Bq/Kg生
			Cs134	17.2	Bq/Kg生	± 5.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Red beans	Kamikatayose, Taira, Iwaki	Nov-11	Cs137	4.0	Bq/Kg生	± 3.2	Bq/Kg生	4.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Red beans	Kamiyamaguchi, Taira, Iwaki	Nov-11	Cs137	3.6	Bq/Kg生	± 3.4	Bq/Kg生	3.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Black beans	Iritono, Tono, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Soybeans	Hayama, Iwaki	Nov-11	Cs137	10.1	Bq/Kg生	± 4.5	Bq/Kg生	10.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Soybeans	Shimotakaku, Taira, Iwaki	Dec-11	Cs137	12.6	Bq/Kg生	± 5.1	Bq/Kg生	20.2	Cs137	7.0	Bq/Kg生
			Cs134	7.6	Bq/Kg生	± 3.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Salted plum	Tomioka, Futaba	Jun-11	Cs137	3.3	Bq/Kg生	± 2.6	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Salted plum	Ena, Iwaki	Jun-11	Cs137	82.5	Bq/Kg生	± 17.4	Bq/Kg生	150	Cs137	7.0	Bq/Kg生
			Cs134	67.2	Bq/Kg生	± 13.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Salted plums (with seed)	Tabito, Iwaki	May-11	Cs137	59.0	Bq/Kg生	± 13.0	Bq/Kg生	111	Cs137	7.0	Bq/Kg生
			Cs134	52.1	Bq/Kg生	± 10.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Miso	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Konjac	Minamidai, Kanayama, Iwaki	Nov-11	Cs137	4.6	Bq/Kg生	± 3.8	Bq/Kg生	4.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Butterbur (salted)	Tabito, Iwaki	May-11	Cs137	7.9	Bq/Kg生	± 3.9	Bq/Kg生	15.0	Cs137	7.0	Bq/Kg生
			Cs134	7.1	Bq/Kg生	± 3.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yellow pickled radish	Youkoudai, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	3.3	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Mashed potato	Fujima, Taira, Iwaki	Jul-11	Cs137	3.6	Bq/Kg生	± 2.5	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
School lunch	Unknown	Feb-12	Cs137	2.4	Bq/Kg生	± 2.0	Bq/Kg生	2.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
School lunch	Unknown	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Milk	Hokkaido	Feb-12	Cs137	1.7	Bq/Kg生	± 1.4	Bq/Kg生	1.7	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Milk	Motomiya	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Breast milk	Sanuka, Iwaki	Feb-12	Cs137	2.4	Bq/Kg生	± 1.4	Bq/Kg生	2.4	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Sekimoto, Kitaibaraki	Feb-12	Cs137	1.7	Bq/Kg生	± 1.3	Bq/Kg生	1.7	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Hosoya, Yotsukura, Iwaki	Feb-12	Cs137	1.4	Bq/Kg生	± 1.3	Bq/Kg生	1.4	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Tamatsuyu, Izumi, Iwaki	Feb-12	Cs137	2.2	Bq/Kg生	± 1.4	Bq/Kg生	2.2	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Kaminagai, Miwa, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Well water	Miwa, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 1.4	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Miwa, Iwaki	Feb-12	Cs137	1.5	Bq/Kg生	± 1.3	Bq/Kg生	1.5	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Fujima, Taira, Iwaki	Feb-12	Cs137	2.5	Bq/Kg生	± 1.4	Bq/Kg生	2.5	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Kamikamado, Watanabe, Iwaki	Feb-12	Cs137	1.4	Bq/Kg生	± 1.3	Bq/Kg生	1.4	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Well water	Tabito, Iwaki	Feb-12	Cs137	1.4	Bq/Kg生	± 1.4	Bq/Kg生	1.4	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Nishiki, Iwaki	Unknown	Cs137	2.3	Bq/Kg生	± 1.4	Bq/Kg生	2.3	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Yotsukura, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Oohara, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.0	Cs137	5.0	Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.3	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Okaona, Onahama, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	5.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.2	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Kamiyunagaya, Joban, Iwaki	Feb-12	Cs137	1.8	Bq/Kg生	± 1.4	Bq/Kg生	1.8	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Ueda, Iwaki	Feb-12	Cs137	1.7	Bq/Kg生	± 1.3	Bq/Kg生	1.7	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Mineral water	Unknown	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Mineral water	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Water of melted snow	Yamada, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Watado, Miwa, Iwaki	Feb-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Spring water	Shibahara, Ogawa, Iwaki	Feb-12	Cs137	1.4	Bq/Kg生	± 1.4	Bq/Kg生	1.4	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Spring water	Nakoso, Iwaki	Feb-12	Cs137	1.4	Bq/Kg生	± 1.4	Bq/Kg生	3.0	Cs137	5.0	Bq/Kg生
			Cs134	1.6	Bq/Kg生	± 1.3	Bq/Kg生		Cs134	3.0	Bq/Kg生
Fallen leaves②	Kashima, Iwaki	Feb-12	Cs137	438	Bq/Kg生	± 96	Bq/Kg生	760	Cs137	7.0	Bq/Kg生
			Cs134	322	Bq/Kg生	± 68	Bq/Kg生		Cs134	6.0	Bq/Kg生
Fallen leaves①	Kashima, Iwaki	Feb-12	Cs137	392	Bq/Kg生	± 86	Bq/Kg生	637	Cs137	7.0	Bq/Kg生
			Cs134	245	Bq/Kg生	± 54	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bamboo charcoal	Joban, Iwaki	Feb-12	Cs137	673	Bq/Kg生	± 135	Bq/Kg生	1,225	Cs137	7.0	Bq/Kg生
			Cs134	552	Bq/Kg生	± 110	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bamboo vinegar	Joban, Iwaki	Feb-12	Cs137	2.5	Bq/Kg生	± 1.4	Bq/Kg生	4.3	Cs137	5.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.3	Bq/Kg生		Cs134	3.0	Bq/Kg生
Field soil	Moroni, Akai, Iwaki	Feb-12	Cs137	395.0	Bq/Kg生	± 79.0	Bq/Kg生	740	Cs137	8.0	Bq/Kg乾
			Cs134	345.0	Bq/Kg生	± 69.0	Bq/Kg生		Cs134	8.0	Bq/Kg乾