



# Radiation Measurement Results of 369 Items in March 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Marumori, Igu, Miyagi	Oct-11	Cs137	6.6 Bq/Kg生	± 3.0 Bq/Kg生	14.4	Cs137	7.0 Bq/Kg生	
			Cs134	7.8 Bq/Kg生	± 2.8 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Tome, Miyagi	Unknown	Cs137	3.7 Bq/Kg生	± 2.4 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Iwate	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.8	Cs137	7.0 Bq/Kg生	
			Cs134	3.8 Bq/Kg生	± 2.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Yamagata	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Tochigi	Oct-11	Cs137	3.0 Bq/Kg生	± 2.3 Bq/Kg生	5.8	Cs137	7.0 Bq/Kg生	
			Cs134	2.8 Bq/Kg生	± 2.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Chiba	Oct-11	Cs137	1.8 Bq/Kg生	± 1.7 Bq/Kg生	1.8	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Furudono, Ishikawa	Oct-11	Cs137	3.7 Bq/Kg生	± 2.2 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Ootama, Adachi	Oct-12	Cs137	5.2 Bq/Kg生	± 2.7 Bq/Kg生	5.2	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Yabuki, Nishishirakawa	Oct-12	Cs137	2.8 Bq/Kg生	± 2.3 Bq/Kg生	2.8	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Taishin, Shirakawa	Oct-11	Cs137	4.5 Bq/Kg生	± 2.0 Bq/Kg生	4.5	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Brown rice	Niida, Sukagawa	Oct-11	Cs137	9.9 Bq/Kg生	± 3.7 Bq/Kg生	15.2	Cs137	7.0 Bq/Kg生	
			Cs134	5.3 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Soma	Oct-11	Cs137	6.5 Bq/Kg生	± 2.2 Bq/Kg生	10.1	Cs137	7.0 Bq/Kg生	
			Cs134	3.6 Bq/Kg生	± 1.6 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Ookuma, Futaba	Nov-10	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Shimokobana, Naraha, Futaba	Oct-10	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.6	Cs137	7.0 Bq/Kg生	
			Cs134	3.6 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Kawamae, Iwaki	Oct-11	Cs137	3.6 Bq/Kg生	± 2.4 Bq/Kg生	6.5	Cs137	7.0 Bq/Kg生	
			Cs134	2.9 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Kamiogawa, Ogawa, Iwaki	Nov-11	Cs137	2.8 Bq/Kg生	± 2.2 Bq/Kg生	5.4	Cs137	7.0 Bq/Kg生	
			Cs134	2.6 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Ogawa, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Brown rice	Ogawa, Iwaki	Nov-11	Cs137	2.2 Bq/Kg生	± 1.7 Bq/Kg生	4.0	Cs137	7.0 Bq/Kg生	
			Cs134	1.8 Bq/Kg生	± 1.5 Bq/Kg生		Cs134	6.0 Bq/Kg生	

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Yoshima, Iwaki	Oct-10	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Fujima, Taira, Iwaki	Oct-11	Cs137	5.5	Bq/Kg生	± 2.8	Bq/Kg生	11.1	Cs137	7.0	Bq/Kg生
			Cs134	5.6	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nishiyamashita, Taira, Iwaki	Oct-11	Cs137	3.8	Bq/Kg生	± 1.9	Bq/Kg生	6.6	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Takaku, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.4	Bq/Kg生	6.2	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	3.5	Bq/Kg生	± 1.8	Bq/Kg生	6.0	Cs137	6.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Shimotakaku, Taira, Iwaki	Sep-11	Cs137	3.0	Bq/Kg生	± 2.5	Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nakakabeya, Taira, Iwaki	Sep-11	Cs137	3.4	Bq/Kg生	± 2.5	Bq/Kg生	6.5	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Yotsukura, Iwaki	Oct-11	Cs137	3.0	Bq/Kg生	± 2.3	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Yotsukura, Iwaki	Oct-11	Cs137	6.6	Bq/Kg生	± 2.9	Bq/Kg生	11.7	Cs137	7.0	Bq/Kg生
			Cs134	5.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nagatomo, Yotsukura, Iwaki	Oct-11	Cs137	4.4	Bq/Kg生	± 2.4	Bq/Kg生	8.4	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kitsunozuka, Yotsukura, Iwaki	Oct-11	Cs137	7.1	Bq/Kg生	± 2.7	Bq/Kg生	10.9	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Noda, Onahama, Iwaki	Oct-11	Cs137	3.9	Bq/Kg生	± 2.5	Bq/Kg生	6.9	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nagasaki, Iwaki	Jun-11	Cs137	5.2	Bq/Kg生	± 3.0	Bq/Kg生	11.4	Cs137	7.0	Bq/Kg生
			Cs134	6.2	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Ishizuka, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.6	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Mizunoya, Joban, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.1	Cs137	7.0	Bq/Kg生
			Cs134	1.1	Bq/Kg生	± 0.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	2.8	Bq/Kg生	± 2.3	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	6.0	Bq/Kg生	± 2.8	Bq/Kg生	9.5	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Izumi, Iwaki	Oct-11	Cs137	3.4	Bq/Kg生	± 2.0	Bq/Kg生	3.4	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Takijiri, Izumi, Iwaki	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.6	Cs137	6.0	Bq/Kg生
			Cs134	1.6	Bq/Kg生	± 1.4	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Ueda, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.3	Bq/Kg生	5.2	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Kadono, Tono, Iwaki	Oct-11	Cs137	9.7	Bq/Kg生	± 3.4	Bq/Kg生	16.6	Cs137	7.0	Bq/Kg生
			Cs134	6.9	Bq/Kg生	± 2.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Iritono, Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.1	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Watanabe, Iwaki	Oct-11	Cs137	2.7	Bq/Kg生	± 2.2	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Watanabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Matsugoya, Watanabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.8	Cs137	6.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 1.5	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Yamanokami, Yamada, Iwaki	Oct-11	Cs137	2.8	Bq/Kg生	± 2.3	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Soeno, Iwaki	Oct-12	Cs137	4.9	Bq/Kg生	± 2.5	Bq/Kg生	4.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Nakoso, Iwakik	Oct-11	Cs137	3.1	Bq/Kg生	± 2.3	Bq/Kg生	6.6	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Azumada, Iwaki	Oct-11	Cs137	2.3	Bq/Kg生	± 1.9	Bq/Kg生	4.2	Cs137	6.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Brown rice	Aksaka, Kawabe, Iwaki	Oct-11	Cs137	7.9	Bq/Kg生	± 3.4	Bq/Kg生	11.1	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Brown rice	Unknown	Oct-11	Cs137	3.4	Bq/Kg生	± 2.9	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Hokkaido	Oct-11	Cs137	5.2	Bq/Kg生	± 2.6	Bq/Kg生	5.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yonezawa, Yamagata	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.5	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miyagi	Oct-11	Cs137	6.2	Bq/Kg生	± 2.8	Bq/Kg生	10.6	Cs137	7.0	Bq/Kg生
			Cs134	4.4	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Toyomuro, Kakuda, Miyagi	Oct-11	Cs137	3.0	Bq/Kg生	± 2.2	Bq/Kg生	3.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Oyama, Kakuda, Miyagi	Oct-11	Cs137	0.0	Bq/Kg生	± 5.6	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ibaraki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.7	Cs137	7.0	Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ibaraki	Oct-11	Cs137	2.8	Bq/Kg生	± 2.1	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Fukushima	Oct-12	Cs137	2.3	Bq/Kg生	± 2.2	Bq/Kg生	2.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nihonmatsu	Oct-11	Cs137	3.5	Bq/Kg生	± 2.1	Bq/Kg生	8.3	Cs137	6.0	Bq/Kg生
			Cs134	4.8	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Nihonmatsu	Oct-11	Cs137	7.0	Bq/Kg生	± 2.6	Bq/Kg生	15.3	Cs137	6.0	Bq/Kg生
			Cs134	8.3	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Shibukawa, Nihonmatsu	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shiozawa, Nihonmatsu	Nov-11	Cs137	37.9	Bq/Kg生	± 8.8	Bq/Kg生	67.7	Cs137	7.0	Bq/Kg生
			Cs134	29.8	Bq/Kg生	± 6.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Katahira, Koriyama	Oct-11	Cs137	3.6	Bq/Kg生	± 2.6	Bq/Kg生	6.8	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Tamura, Koriyama	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.3	Cs137	6.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Ogoe, Tamura	Oct-11	Cs137	1.7	Bq/Kg生	± 1.6	Bq/Kg生	1.7	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Iwase, Sukagawa	Oct-11	Cs137	3.2	Bq/Kg生	± 2.2	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shirakawa	Sep-11	Cs137	2.8	Bq/Kg生	± 2.0	Bq/Kg生	4.8	Cs137	7.0	Bq/Kg生
			Cs134	2.0	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Shirakawa	Oct-11	Cs137	4.7	Bq/Kg生	± 2.2	Bq/Kg生	6.9	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Tanakura, Higashishirakawa	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.4	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Aizubange, Kawanuma	Oct-11	Cs137	2.3	Bq/Kg生	± 2.1	Bq/Kg生	2.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimokobana, Naraha, Futaba	Oct-10	Cs137	2.8	Bq/Kg生	± 2.2	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Naraha, Namie	Oct-10	Cs137	2.8	Bq/Kg生	± 2.0	Bq/Kg生	2.8	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Hirata, Ishikawa	Oct-11	Cs137	2.3	Bq/Kg生	± 1.9	Bq/Kg生	2.3	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Furudono, Ishikawa	Oct-11	Cs137	3.8	Bq/Kg生	± 2.0	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miwa, Iwaki	Oct-11	Cs137	2.2	Bq/Kg生	± 2.2	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miwa, Iwaki	Nov-11	Cs137	2.9	Bq/Kg生	± 2.3	Bq/Kg生	5.4	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Miwa, Iwaki	Oct-12	Cs137	3.9	Bq/Kg生	± 1.8	Bq/Kg生	3.9	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Shimoichigaya, Miwa, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.4	Bq/Kg生	5.5	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kusawatari, Miwa, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 2.3	Bq/Kg生	4.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimookeuri, Kawamae, Iwaki	Sep-11	Cs137	4.7	Bq/Kg生	± 2.7	Bq/Kg生	7.9	Cs137	6.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Shimookeuri, Kawamae, Iwaki	Sep-11	Cs137	4.7	Bq/Kg生	± 2.7	Bq/Kg生	7.9	Cs137	6.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Takahagi, Ogawa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.1	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ogawa, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.6	Cs137	7.0	Bq/Kg生
			Cs134	2.6	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ogawa, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.4	Cs137	7.0	Bq/Kg生
			Cs134	1.4	Bq/Kg生	± 1.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nishiogawa, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.2	Cs137	7.0	Bq/Kg生
			Cs134	3.2	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Unknown	Cs137	3.4	Bq/Kg生	± 2.4	Bq/Kg生	6.3	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	6.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	2.5	Bq/Kg生	± 2.1	Bq/Kg生	4.6	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Yotsukura, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.2	Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimoniida, Yotsukura, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.5	Bq/Kg生	6.2	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nagatomo, Yotsukura, Iwaki	Oct-11	Cs137	2.2	Bq/Kg生	± 2.1	Bq/Kg生	4.9	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Rice	Komagome, Yotsukura, Iwaki	Nov-11	Cs137	2.5	Bq/Kg生	± 3.4	Bq/Kg生	7.2	Cs137	8.0	Bq/Kg生
			Cs134	4.7	Bq/Kg生	± 4.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Rice	Shimokatayose, Taira, Iwaki	Oct-11	Cs137	2.5	Bq/Kg生	± 2.5	Bq/Kg生	6.5	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kinuya, Taira, Iwaki	Oct-11	Cs137	2.4	Bq/Kg生	± 2.1	Bq/Kg生	5.8	Cs137	7.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kabeya, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.1	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kabeya, Taira, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.3	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nakakabeya, Taira, Iwaki	Sep-11	Cs137	2.3	Bq/Kg生	± 2.1	Bq/Kg生	4.8	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.5	Bq/Kg生	5.8	Cs137	7.0	Bq/Kg生
			Cs134	2.5	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimoyamaguchi, Taira, Iwaki	Oct-11	Cs137	4.0	Bq/Kg生	± 2.4	Bq/Kg生	5.9	Cs137	7.0	Bq/Kg生
			Cs134	1.9	Bq/Kg生	± 1.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Taira, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 2.4	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	4.0	Bq/Kg生	± 2.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nakahirakubo, Taira, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.3	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Shimohirakubo, Taira, Iwaki	Sep-11	Cs137	2.9	Bq/Kg生	± 2.1	Bq/Kg生	5.0	Cs137	7.0	Bq/Kg生
			Cs134	2.1	Bq/Kg生	± 1.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Ishikiriba, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.8	Cs137	6.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 1.5	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Takaku, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.7	Cs137	6.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 1.5	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Shimotakaku, Taira, Iwaki	Sep-11	Cs137	4.4	Bq/Kg生	± 2.2	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Kamiogoe, Taira, Iwaki	Oct-11	Cs137	4.9	Bq/Kg生	± 2.5	Bq/Kg生	4.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Koizumi, Taira, Iwaki	Oct-11	Cs137	3.1	Bq/Kg生	± 2.7	Bq/Kg生	6.1	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Izumizaki, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg生	± 2.3	Bq/Kg生	5.6	Cs137	8.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Rice	Aratame, Taira, Iwaki	Oct-11	Cs137	2.9	Bq/Kg生	± 2.1	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Nagasaki, Iwaki	Oct-11	Cs137	4.8	Bq/Kg生	± 2.0	Bq/Kg生	8.3	Cs137	6.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	4.0	Bq/Kg生
Rice	Sumiyoshi, Onahama, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Okaona, Onahama, Iwaki	Oct-11	Cs137	10.7	Bq/Kg生	± 3.3	Bq/Kg生	17.8	Cs137	7.0	Bq/Kg生
			Cs134	7.1	Bq/Kg生	± 2.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Okaona, Onahama, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 4.2	Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 3.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Rice	Otsurugi, Izumi, Iwaki	Mar-12	Cs137	202	Bq/Kg生	± 41.0	Bq/Kg生	340	Cs137	7.0	Bq/Kg生
			Cs134	138	Bq/Kg生	± 28.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Rice	Honya, Izumi, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Rice	Tono, Iwaki	Oct-11	Cs137	3.0 Bq/Kg生	± 2.4 Bq/Kg生	6.7	Cs137	7.0 Bq/Kg生	
			Cs134	3.7 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Tono, Iwaki	Oct-11	Cs137	2.8 Bq/Kg生	± 2.2 Bq/Kg生	2.8	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Tono, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	7.2	Cs137	7.0 Bq/Kg生	
			Cs134	7.2 Bq/Kg生	± 2.6 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.5	Cs137	6.0 Bq/Kg生	
			Cs134	2.5 Bq/Kg生	± 1.6 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	3.1 Bq/Kg生	± 2.3 Bq/Kg生	3.1	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Iritono, Tono, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Watanabe, Iwaki	Oct-11	Cs137	4.2 Bq/Kg生	± 2.3 Bq/Kg生	6.7	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kamikamado, Watanabe, Iwaki	Sep-11	Cs137	2.8 Bq/Kg生	± 2.0 Bq/Kg生	2.8	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kawabe, Iwaki	Oct-11	Cs137	3.2 Bq/Kg生	± 2.2 Bq/Kg生	5.1	Cs137	7.0 Bq/Kg生	
			Cs134	1.9 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kawabe, Iwaki	Oct-11	Cs137	3.0 Bq/Kg生	± 2.3 Bq/Kg生	3.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Nishiki, Iwaki	Oct-11	Cs137	4.1 Bq/Kg生	± 2.7 Bq/Kg生	6.5	Cs137	7.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Sanuka, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.9	Cs137	7.0 Bq/Kg生	
			Cs134	3.9 Bq/Kg生	± 2.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Yamada, Iwaki	Oct-12	Cs137	3.0 Bq/Kg生	± 1.7 Bq/Kg生	5.4	Cs137	6.0 Bq/Kg生	
			Cs134	2.4 Bq/Kg生	± 1.5 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Tabito, Iwaki	Oct-11	Cs137	3.1 Bq/Kg生	± 2.2 Bq/Kg生	5.2	Cs137	7.0 Bq/Kg生	
			Cs134	2.1 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Misawa, Iwaki	Oct-11	Cs137	4.3 Bq/Kg生	± 2.6 Bq/Kg生	8.7	Cs137	7.0 Bq/Kg生	
			Cs134	4.4 Bq/Kg生	± 2.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Kamimousu, Yamada, Iwaki	Oct-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.2	Cs137	7.0 Bq/Kg生	
			Cs134	2.2 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Azumada, Iwaki	Oct-11	Cs137	2.1 Bq/Kg生	± 1.5 Bq/Kg生	3.7	Cs137	6.0 Bq/Kg生	
			Cs134	1.6 Bq/Kg生	± 1.4 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Akakona, Ebata, Iwaki	Oct-11	Cs137	1.8 Bq/Kg生	± 1.5 Bq/Kg生	3.7	Cs137	6.0 Bq/Kg生	
			Cs134	1.9 Bq/Kg生	± 1.3 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Rice	Unknown	Oct-11	Cs137	2.8 Bq/Kg生	± 2.3 Bq/Kg生	4.9	Cs137	7.0 Bq/Kg生	
			Cs134	2.1 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Rice	Unknown	Oct-11	Cs137	6.9 Bq/Kg生	± 2.8 Bq/Kg生	6.9	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Glutinous rice	Taishin, Shirakawa	Oct-11	Cs137	1.6 Bq/Kg生	± 1.6 Bq/Kg生	1.6	Cs137	6.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生	
Glutinous rice	Shirakawa	Oct-11	Cs137	11.8 Bq/Kg生	± 3.5 Bq/Kg生	16.9	Cs137	7.0 Bq/Kg生	
			Cs134	5.1 Bq/Kg生	± 2.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Glutinous rice	Shimookeuri, Kawamae, Iwaki	Oct-11	Cs137	5.5 Bq/Kg生	± 2.7 Bq/Kg生	8.4	Cs137	7.0 Bq/Kg生	
			Cs134	2.9 Bq/Kg生	± 1.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Glutinous rice	Hosoya, Yotsukura, Iwaki	Oct-11	Cs137	17.1 Bq/Kg生	± 4.6 Bq/Kg生	33.0	Cs137	7.0 Bq/Kg生	
			Cs134	15.9 Bq/Kg生	± 4.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Glutinous rice	Kitakabeya, Taira, Iwaki	Oct-11	Cs137	3.7	Bq/Kg生	± 2.0	Bq/Kg生	5.9	Cs137	6.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Glutinous rice	Shimokatayose, Taira, Iwaki	Oct-11	Cs137	4.3	Bq/Kg生	± 2.5	Bq/Kg生	6.6	Cs137	7.0	Bq/Kg生
			Cs134	2.3	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice	Iritono, Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	7.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Glutinous rice (brown rice)	Hosoya, Yotsukura, Iwaki	Oct-12	Cs137	7.7	Bq/Kg生	± 3.3	Bq/Kg生	11.4	Cs137	7.0	Bq/Kg生
			Cs134	3.7	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Flour	Nagasaki, Iwaki	Unknown	Cs137	0.4	Bq/Kg生	± 13.0	Bq/Kg生	0.6	Cs137	8.0	Bq/Kg生
			Cs134	0.2	Bq/Kg生	± 8.4	Bq/Kg生		Cs134	8.0	Bq/Kg生
Fried rice	Kamiogoe, Taira, Iwaki	Oct-11	Cs137	2.7	Bq/Kg生	± 2.7	Bq/Kg生	2.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Miwa, Iwaki	Dec-11	Cs137	34.3	Bq/Kg生	± 11.3	Bq/Kg生	58.2	Cs137	7.0	Bq/Kg生
			Cs134	23.9	Bq/Kg生	± 8.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Yagawase, Taira, Iwaki	Mar-12	Cs137	19.6	Bq/Kg生	± 8.2	Bq/Kg生	19.6	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Ancient rice	Shimokajiro, Onahama, Iwaki	Oct-11	Cs137	4.5	Bq/Kg生	± 3.1	Bq/Kg生	4.5	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Germinated rice	Ota, Hitachi, Ibaraki	Oct-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shirakawa	Sep-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Ohisa, Iwaki	Jul-11	Cs137	6.5	Bq/Kg生	± 4.1	Bq/Kg生	19.7	Cs137	7.0	Bq/Kg生
			Cs134	13.2	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimookeuri, Kawamae, Iwaki	Aug-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Nishiogawa, Ogawa, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Koizumi, Taira, Iwaki	Jul-11	Cs137	6.0	Bq/Kg生	± 3.3	Bq/Kg生	10.2	Cs137	7.0	Bq/Kg生
			Cs134	4.2	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Shimoogoe, Taira, Iwaki	Jul-11	Cs137	3.2	Bq/Kg生	± 2.9	Bq/Kg生	11.8	Cs137	7.0	Bq/Kg生
			Cs134	8.6	Bq/Kg生	± 2.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kamiyamaguchi, Taira, Iwaki	Jun-11	Cs137	2.9	Bq/Kg生	± 2.7	Bq/Kg生	6.8	Cs137	7.0	Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kouya, Uchigo, Iwaki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kouya, Uchigo, Iwaki	Jul-12	Cs137	5.1	Bq/Kg生	± 3.2	Bq/Kg生	5.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Nagasaki, Iwaki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Ena, Iwaki	Jun-11	Cs137	3.0	Bq/Kg生	± 2.6	Bq/Kg生	7.1	Cs137	7.0	Bq/Kg生
			Cs134	4.1	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Seto, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.9	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Ishizuka, Iwaki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.5	Cs137	7.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 3.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Okaona, Onahama, Iwaki	Jul-11	Cs137	4.1	Bq/Kg生	± 2.2	Bq/Kg生	4.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Potato	Okaona, Onahama, Iwaki	Jul-11	Cs137	5.1	Bq/Kg生	± 3.1	Bq/Kg生	5.1	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Kamikajiro, Onahama, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Yanagi, Ushiroda, Iwaki	Apr-11	Cs137	4.0	Bq/Kg生	± 3.3	Bq/Kg生	7.5	Cs137	7.0	Bq/Kg生
			Cs134	3.5	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potato	Nakoso, Iwaki	Jun-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Taro	Kamiyamaguchi, Taira, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Taro	Tanmatsuyu, Izumi, Iwaki	Sep-11	Cs137	2.3	Bq/Kg生	± 6.8	Bq/Kg生	4.1	Cs137	8.0	Bq/Kg生
			Cs134	1.8	Bq/Kg生	± 5.3	Bq/Kg生		Cs134	8.0	Bq/Kg生
Taro	Mizunoya, Joban, Iwaki	Mar-12	Cs137	5.2	Bq/Kg生	± 4.4	Bq/Kg生	5.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Taro	Okaona, Onahama, Iwaki	Nov-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Sweet potato	Hosoya, Yotsukura, Iwaki	Nov-11	Cs137	4.1	Bq/Kg生	± 3.7	Bq/Kg生	10.4	Cs137	7.0	Bq/Kg生
			Cs134	6.3	Bq/Kg生	± 3.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried sweet potato	Hitachiota, Ibaraki	Dec-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chinese yam	Kadono, Tono, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.0	Cs137	8.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Jerusalem artichoke	Nakashio, Taira, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Jerusalem artichoke	Nakashio, Taira, Iwaki	Mar-12	Cs137	4.3	Bq/Kg生	± 3.4	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Pumpkin	Kamiyamaguchi, Taira, Iwaki	Jul-11	Cs137	3.4	Bq/Kg生	± 3.0	Bq/Kg生	3.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Ohisa, Iwaki	Mar-12	Cs137	10.8	Bq/Kg生	± 4.8	Bq/Kg生	18.7	Cs137	7.0	Bq/Kg生
			Cs134	7.9	Bq/Kg生	± 3.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Koizumi, Taira, Iwaki	Mar-12	Cs137	4.3	Bq/Kg生	± 3.3	Bq/Kg生	7.9	Cs137	7.0	Bq/Kg生
			Cs134	3.6	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Kamiogoe, Taira, Iwaki	May-11	Cs137	6.1	Bq/Kg生	± 4.1	Bq/Kg生	11.4	Cs137	7.0	Bq/Kg生
			Cs134	5.3	Bq/Kg生	± 3.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shimohirakubo, Taira, Iwaki	Mar-12	Cs137	3.7	Bq/Kg生	± 2.4	Bq/Kg生	3.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Kouya, Uchigo, Iwaki	Dec-11	Cs137	3.8	Bq/Kg生	± 2.2	Bq/Kg生	3.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Shimoyada, Kashima, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Moegidai, Izumi, Iwaki	Mar-12	Cs137	3.3	Bq/Kg生	± 3.0	Bq/Kg生	3.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Nishiki, Iwaki	Mar-12	Cs137	4.7	Bq/Kg生	± 2.4	Bq/Kg生	4.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese white radish	Nishiki, Iwaki	Mar-12	Cs137	4.7	Bq/Kg生	± 2.4	Bq/Kg生	4.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Japanese white radish	Nakoso, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.7	Cs137	7.0 Bq/Kg生	
			Cs134	3.7 Bq/Kg生	± 3.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish	Namesawa, Yamada, Iwaki	Mar-12	Cs137	4.4 Bq/Kg生	± 2.8 Bq/Kg生	4.4	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Round radish	Kitakabeya, Taira, Iwaki	Mar-12	Cs137	3.6 Bq/Kg生	± 5.3 Bq/Kg生	6.6	Cs137	8.0 Bq/Kg生	
			Cs134	3.0 Bq/Kg生	± 4.4 Bq/Kg生		Cs134	8.0 Bq/Kg生	
Japanese white radish (with leaves)	Nakoso, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.1	Cs137	7.0 Bq/Kg生	
			Cs134	4.1 Bq/Kg生	± 3.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Japanese white radish(leaves)	Satogaoka, Iwaki	Mar-12	Cs137	15.7 Bq/Kg生	± 6.9 Bq/Kg生	24.5	Cs137	7.0 Bq/Kg生	
			Cs134	8.8 Bq/Kg生	± 5.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Dried Japanese radish	Kamikajiro, Onahama, Iwaki	Dec-12	Cs137	28.5 Bq/Kg生	± 11.6 Bq/Kg生	41.0	Cs137	7.0 Bq/Kg生	
			Cs134	12.5 Bq/Kg生	± 8.8 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Carrot	Kishi, Tairakamatat, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Carrot	Ori, Yoshima, Iwaki	Mar-12	Cs137	5.9 Bq/Kg生	± 3.6 Bq/Kg生	9.7	Cs137	7.0 Bq/Kg生	
			Cs134	3.8 Bq/Kg生	± 3.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese cabbage	Kubo, Kashima, Iwaki	Feb-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	2.2	Cs137	7.0 Bq/Kg生	
			Cs134	2.2 Bq/Kg生	± 2.1 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese cabbage	Okaona, Onahama, Iwaki	Mar-12	Cs137	5.6 Bq/Kg生	± 5.4 Bq/Kg生	5.6	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese cabbage	Yokodai, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese cabbage	Izumigaoka, Iwaki	Feb-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.8	Cs137	7.0 Bq/Kg生	
			Cs134	4.8 Bq/Kg生	± 4.4 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese cabbage	Shimokawa, Izumi, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Chinese cabbage	Sanuka, Iwaki	Dec-11	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	3.4	Cs137	7.0 Bq/Kg生	
			Cs134	3.4 Bq/Kg生	± 3.2 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Ohisa, Iwaki	Jul-11	Cs137	6.6 Bq/Kg生	± 4.0 Bq/Kg生	6.6	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Shimotakaku, Taira, Iwaki	Mar-12	Cs137	4.5 Bq/Kg生	± 3.4 Bq/Kg生	8.6	Cs137	7.0 Bq/Kg生	
			Cs134	4.1 Bq/Kg生	± 2.9 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Nakashio, Taira, Iwaki	Mar-12	Cs137	5.1 Bq/Kg生	± 3.0 Bq/Kg生	8.2	Cs137	7.0 Bq/Kg生	
			Cs134	3.1 Bq/Kg生	± 2.7 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Shio, Taira, Iwaki	Mar-12	Cs137	3.6 Bq/Kg生	± 3.6 Bq/Kg生	3.6	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Ishizuka, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Okaona, Onahama, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	4.4	Cs137	7.0 Bq/Kg生	
			Cs134	4.4 Bq/Kg生	± 3.3 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Mizunoya, Joban, Iwaki	Mar-12	Cs137	5.8 Bq/Kg生	± 4.5 Bq/Kg生	12.5	Cs137	7.0 Bq/Kg生	
			Cs134	6.7 Bq/Kg生	± 3.7 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Kaneyama, Numabe, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Soeno, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	
Green onion	Higashida, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	7.0 Bq/Kg生	
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	6.0 Bq/Kg生	

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Onion	Nakashio, Taira, Iwaki	Jun-12	Cs137	4.8	Bq/Kg生	± 2.8	Bq/Kg生	4.8	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Kamikatayose, Taira, Iwaki	Jul-01	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Kabeya, Taira, Iwaki	Jul-11	Cs137	7.1	Bq/Kg生	± 3.9	Bq/Kg生	11.6	Cs137	7.0	Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 3.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Miyama, Uchigo, Iwaki	Jul-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.4	Cs137	8.0	Bq/Kg生
			Cs134	2.4	Bq/Kg生	± 6.1	Bq/Kg生		Cs134	8.0	Bq/Kg生
Onion	Nagasaki, Iwaki	Jul-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	7.3	Cs137	7.0	Bq/Kg生
			Cs134	7.3	Bq/Kg生	± 4.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Onion	Yanagi, Ushiroda, Iwaki	Mar-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.4	Cs137	7.0	Bq/Kg生
			Cs134	3.4	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spring onion	Higashida, Iwaki	Feb-12	Cs137	2.3	Bq/Kg生	± 2.2	Bq/Kg生	2.3	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Shibahara, Ogawa, Iawki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	7.4	Cs137	7.0	Bq/Kg生
			Cs134	7.4	Bq/Kg生	± 5.9	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Ori, Yoshima, Iwaki	Mar-12	Cs137	3.8	Bq/Kg生	± 7.6	Bq/Kg生	16.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Yoshima, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.2	Cs137	8.0	Bq/Kg生
			Cs134	1.2	Bq/Kg生	± 4.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Spinach	Shimotakaku, Taira, Iwaki	Mar-12	Cs137	6.7	Bq/Kg生	± 4.7	Bq/Kg生	6.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Kamiyunagaya, Joban, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Spinach	Soeno, Iwaki	Mar-12	Cs137	4.0	Bq/Kg生	± 6.0	Bq/Kg生	7.9	Cs137	8.0	Bq/Kg生
			Cs134	3.9	Bq/Kg生	± 4.8	Bq/Kg生		Cs134	8.0	Bq/Kg生
Qing-geng-cai	Namesawa, Yamada, Iwaki	Mar-12	Cs137	4.7	Bq/Kg生	± 3.8	Bq/Kg生	4.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese mustard spinach	Inoue, Yamada, Iwaki	Mar-12	Cs137	4.0	Bq/Kg生	± 3.9	Bq/Kg生	4.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Japanese mustard spinach	Sawagami, Kawabe, Iwaki	Mar-12	Cs137	15.8	Bq/Kg生	± 6.8	Bq/Kg生	27.3	Cs137	7.0	Bq/Kg生
			Cs134	11.5	Bq/Kg生	± 5.5	Bq/Kg生		Cs134	6.0	Bq/Kg生
Canola flower	Yokodai, Iwaki	Mar-12	Cs137	6.7	Bq/Kg生	± 9.7	Bq/Kg生	11.7	Cs137	8.0	Bq/Kg生
			Cs134	5.0	Bq/Kg生	± 6.6	Bq/Kg生		Cs134	8.0	Bq/Kg生
Canola flower	Akai, Taira, Iwaki	Mar-12	Cs137	4.4	Bq/Kg生	± 4.1	Bq/Kg生	4.4	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Canola flower	Shisawa, Nakoso, Iwaki	Jan-11	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.2	Cs137	7.0	Bq/Kg生
			Cs134	5.2	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potherb mustard	Nakashio, Taira, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	6.2	Cs137	7.0	Bq/Kg生
			Cs134	6.2	Bq/Kg生	± 4.1	Bq/Kg生		Cs134	6.0	Bq/Kg生
Potherb mustard	Nakashio, Taira, Iwaki	Mar-12	Cs137	3.6	Bq/Kg生	± 3.3	Bq/Kg生	6.6	Cs137	7.0	Bq/Kg生
			Cs134	3.0	Bq/Kg生	± 2.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Angelica keiskei	Wakabadai, Iwaki	Mar-12	Cs137	6.3	Bq/Kg生	± 5.6	Bq/Kg生	11.8	Cs137	8.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 3.8	Bq/Kg生		Cs134	8.0	Bq/Kg生
Broccoli	Tamatsuyu, Izumi, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Canola flower	Numabe, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Canola flower	Miya, Uchigo, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Canola flower	Kamiyagyu, Yotsukura, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	5.5	Cs137	7.0	Bq/Kg生
			Cs134	5.5	Bq/Kg生	± 4.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Cherry tomato	Haramachi, Minamisoma	Aug-11	Cs137	3.6	Bq/Kg生	± 2.4	Bq/Kg生	6.5	Cs137	7.0	Bq/Kg生
			Cs134	2.9	Bq/Kg生	± 2.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Red beans	Shiraiwa, Yotsukura, Iwaki	Nov-11	Cs137	12.5	Bq/Kg生	± 4.6	Bq/Kg生	17.0	Cs137	7.0	Bq/Kg生
			Cs134	4.5	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Snap pea	Sumiyoshi, Onahama, Iwaki	May-11	Cs137	8.8	Bq/Kg生	± 4.8	Bq/Kg生	17.6	Cs137	7.0	Bq/Kg生
			Cs134	8.8	Bq/Kg生	± 4.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
Soybean	Okaona, Onahama, Iwaki	Oct-11	Cs137	5.7	Bq/Kg生	± 3.9	Bq/Kg生	5.7	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Soybean	Miwa, Iwaki	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Aobata bean	Yosawa, Nakoso, Iwaki	Nov-11	Cs137	4.7	Bq/Kg生	± 3.8	Bq/Kg生	8.5	Cs137	7.0	Bq/Kg生
			Cs134	3.8	Bq/Kg生	± 3.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Green bean	Kamiyamauchi, Taira, Iwaki	Aug-11	Cs137	6.3	Bq/Kg生	± 3.9	Bq/Kg生	11.2	Cs137	7.0	Bq/Kg生
			Cs134	4.9	Bq/Kg生	± 3.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Oita	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	3.1	Cs137	7.0	Bq/Kg生
			Cs134	3.1	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Shimoyamauchi, Taira, Iwaki	Mar-12	Cs137	437	Bq/Kg生	± 87.0	Bq/Kg生	769	Cs137	7.0	Bq/Kg生
			Cs134	332	Bq/Kg生	± 66.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Akai, Taira, Iwaki	Mar-12	Cs137	904	Bq/Kg生	± 181	Bq/Kg生	1,624	Cs137	7.0	Bq/Kg生
			Cs134	720	Bq/Kg生	± 144	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Shimoyunagaya, Joban, Iwaki	Oct-11	Cs137	522	Bq/Kg生	± 104	Bq/Kg生	963	Cs137	7.0	Bq/Kg生
			Cs134	441	Bq/Kg生	± 88.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	2136	Bq/Kg生	± 216	Bq/Kg生	3,248	Cs137	8.0	Bq/Kg生
			Cs134	1112	Bq/Kg生	± 115	Bq/Kg生		Cs134	8.0	Bq/Kg生
Shiitake mushroom	Kamikajiro, Onahama, Iwaki	Mar-12	Cs137	496	Bq/Kg生	± 99.0	Bq/Kg生	906	Cs137	7.0	Bq/Kg生
			Cs134	410	Bq/Kg生	± 82.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Tamatsuyu, Izumi, Iwaki	Mar-12	Cs137	92.4	Bq/Kg生	± 20.3	Bq/Kg生	168	Cs137	7.0	Bq/Kg生
			Cs134	75.4	Bq/Kg生	± 15.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Tamatsuyu, Izumi, Iwaki	Mar-12	Cs137	327	Bq/Kg生	± 66.0	Bq/Kg生	581	Cs137	7.0	Bq/Kg生
			Cs134	254	Bq/Kg生	± 51.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Shimokawa, Izumi, Iwaki	Mar-12	Cs137	319	Bq/Kg生	± 64.0	Bq/Kg生	572	Cs137	7.0	Bq/Kg生
			Cs134	253	Bq/Kg生	± 51.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Shiitake mushroom	Nakakamado, Watanabe, Iwaki	Mar-12	Cs137	439	Bq/Kg生	± 88.0	Bq/Kg生	778	Cs137	6.0	Bq/Kg生
			Cs134	339	Bq/Kg生	± 68.0	Bq/Kg生		Cs134	4.0	Bq/Kg生
Shitake mushroom grown in log	Miyamada, Tono, Iwaki	Mar-12	Cs137	203	Bq/Kg生	± 41.0	Bq/Kg生	364	Cs137	7.0	Bq/Kg生
			Cs134	161	Bq/Kg生	± 32.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Dried shiitake mushroom	Shimookeuri, Kawamae, Iwaki	Apr-11	Cs137	1518	Bq/Kg生	± 177	Bq/Kg生	2,126	Cs137	8.0	Bq/Kg生
			Cs134	608	Bq/Kg生	± 81.6	Bq/Kg生		Cs134	8.0	Bq/Kg生
Butterbur sprout	Numabe, Iwaki	Mar-12	Cs137	12.3	Bq/Kg生	± 5.6	Bq/Kg生	27.8	Cs137	7.0	Bq/Kg生
			Cs134	15.5	Bq/Kg生	± 4.8	Bq/Kg生		Cs134	6.0	Bq/Kg生
Butterbur	Tamatsuyu, Izumi, Iwaki	Mar-12	Cs137	15.6	Bq/Kg生	± 12.8	Bq/Kg生	29.1	Cs137	8.0	Bq/Kg生
			Cs134	13.5	Bq/Kg生	± 8.9	Bq/Kg生		Cs134	8.0	Bq/Kg生
Zenmai	Nishiaizu, Yama	Unknown	Cs137	3.0	Bq/Kg生	± 3.7	Bq/Kg生	4.3	Cs137	8.0	Bq/Kg生
			Cs134	1.3	Bq/Kg生	± 3.9	Bq/Kg生		Cs134	8.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
			Cs137	Bq/Kg生	±	Bq/Kg生		Cs137	Bq/Kg生
Bamboo shoot (boiled)	Unknown	Unknown	Cs137	57.7	±	13.9	108.9	Cs137	7.0
			Cs134	51.2	±	11.7		Cs134	6.0
Bamboo shoot (boiled twice)	Unknown	Unknown	Cs137	28.1	±	8.0	47.7	Cs137	7.0
			Cs134	19.6	±	5.7		Cs134	6.0
Kiwi	Kubo, Kashima, Iwaki	Dec-11	Cs137	26.7	±	6.5	55.6	Cs137	7.0
			Cs134	28.9	±	6.4		Cs134	6.0
Chinese citron	Mobara, Chiba	Mar-12	Cs137	16.6	±	6.9	27.3	Cs137	8.0
			Cs134	10.7	±	5.5		Cs134	8.0
Mandarin orange	Ohisa, Iwaki	Mar-12	Cs137	220	±	30.4	349	Cs137	8.0
			Cs134	129	±	17.5		Cs134	8.0
Mandarin orange	Nishiki, Iwaki	Feb-12	Cs137	14.7	±	4.3	28.2	Cs137	7.0
			Cs134	13.5	±	3.6		Cs134	6.0
Mandarin orange	Kishi, Tairakamata, Iwaki	Mar-12	Cs137	28.2	±	6.8	50.6	Cs137	7.0
			Cs134	22.4	±	5.3		Cs134	6.0
Hassaku orange	Akai, Taira, Iwaki	Mar-12	Cs137	26.4	±	6.5	6.7	Cs137	7.0
			Cs134	23.3	±	5.3		Cs134	6.0
Blueberry	Akai, Taira, Iwaki	Jul-11	Cs137	25.7	±	6.3	46.3	Cs137	7.0
			Cs134	20.6	±	4.8		Cs134	6.0
Blueberry	Saizuchikoji, Taira, Iwaki	Unknown	Cs137	28.2	±	7.2	49.7	Cs137	7.0
			Cs134	21.5	±	5.3		Cs134	6.0
Kiwi	Haramachi, Soma	Nov-11	Cs137	57.7	±	12.7	100	Cs137	7.0
			Cs134	42.3	±	9.0		Cs134	6.0
Chinese citron	Okaona, Onahama, Iwaki	Mar-12	Cs137	74.0	±	16.4	141	Cs137	7.0
			Cs134	67.0	±	13.9		Cs134	6.0
Chinese citron	Ojima, Taira	Mar-12	Cs137	40.6	±	9.7	68.5	Cs137	7.0
			Cs134	27.9	±	6.7		Cs134	6.0
Citron	Shibahara, Ogawa, Iwaki	Dec-11	Cs137	61.1	±	13.2	108	Cs137	7.0
			Cs134	46.4	±	9.7		Cs134	6.0
Chestnut	Miyama, Uchigo, Iwaki	Sep-11	Cs137	26.6	±	10.3	52.3	Cs137	8.0
			Cs134	25.7	±	6.9		Cs134	8.0
Dried persimmon	Aratame, Taira, Iwaki	Nov-11	Cs137	54.6	±	12.0	95.4	Cs137	7.0
			Cs134	40.8	±	8.8		Cs134	6.0
Saury	Hokkaido	Unknown	Cs137	2.9	±	3.9	5.6	Cs137	8.0
			Cs134	2.7	±	2.7		Cs134	8.0
Saury	Hokkaido	Unknown	Cs137	4.3	±	3.0	4.3	Cs137	7.0
			Cs134	0.0	±	0.0		Cs134	6.0
Dark sleeper	Misaki, Onahama, Iwaki	Feb-12	Cs137	33.1	±	7.8	56.1	Cs137	7.0
			Cs134	23.0	±	5.4		Cs134	6.0
Octopus	Onahama, Iwaki	Oct-11	Cs137	2.8	±	5.3	4.7	Cs137	8.0
			Cs134	1.9	±	3.8		Cs134	8.0
Wild boar meat	Kawauchi, Futaba	Unknown	Cs137	300	±	60.0	534	Cs137	7.0
			Cs134	234	±	47.0		Cs134	6.0
Wild boar meat	Miwa, Iwaki	Unknown	Cs137	60.7	±	13.3	108	Cs137	7.0
			Cs134	47.7	±	10.0		Cs134	6.0
Wild boar meat	Iwaki	Mar-12	Cs137	251	±	50.0	459	Cs137	7.0
			Cs134	208	±	42.0		Cs134	6.0
Wild boar meat	Iwaki	Jan-12	Cs137	110	±	23.0	204	Cs137	7.0
			Cs134	93.5	±	18.7		Cs134	6.0



## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Egg	Suginami, Taira, Iwaki	Mar-12	Cs137	4.9	Bq/Kg生	± 4.4	Bq/Kg生	4.9	Cs137	8.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Chicken thigh	Japan	Unknown	Cs137	4.2	Bq/Kg生	± 2.1	Bq/Kg生	4.2	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Minced pork meat	Japan	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Chicken breast (minced)	Japan	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.8	Cs137	7.0	Bq/Kg生
			Cs134	2.8	Bq/Kg生	± 2.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Formula	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	4.3	Cs137	7.0	Bq/Kg生
			Cs134	4.3	Bq/Kg生	± 3.4	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yogurt	Unknown	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	8.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	8.0	Bq/Kg生
Chinese citron marmalade	Uchimachimaeda, Uchigo, Iwaki	Feb-12	Cs137	38.0	Bq/Kg生	± 8.1	Bq/Kg生	70.5	Cs137	6.0	Bq/Kg生
			Cs134	32.5	Bq/Kg生	± 6.7	Bq/Kg生		Cs134	4.0	Bq/Kg生
Salted plum	Seto, Iwaki	Jun-11	Cs137	85.9	Bq/Kg生	± 15.6	Bq/Kg生	148	Cs137	8.0	Bq/Kg生
			Cs134	62.3	Bq/Kg生	± 10.5	Bq/Kg生		Cs134	8.0	Bq/Kg生
Salted plum	Kamikatayose, Taira, Iwaki	Jun-11	Cs137	83.1	Bq/Kg生	± 17.3	Bq/Kg生	156	Cs137	7.0	Bq/Kg生
			Cs134	72.8	Bq/Kg生	± 14.3	Bq/Kg生		Cs134	6.0	Bq/Kg生
School lunch	Tsuzura, Uchigo, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.7	Cs137	7.0	Bq/Kg生
			Cs134	2.7	Bq/Kg生	± 1.7	Bq/Kg生		Cs134	6.0	Bq/Kg生
School lunch	Tsuzura, Uchigo, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	2.2	Cs137	7.0	Bq/Kg生
			Cs134	2.2	Bq/Kg生	± 1.6	Bq/Kg生		Cs134	6.0	Bq/Kg生
Miso	Kawamata, Date	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Miso	Nihonmatsu	Unknown	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	7.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Bran	Unknown	Mar-12	Cs137	15.8	Bq/Kg生	± 4.4	Bq/Kg生	27.8	Cs137	7.0	Bq/Kg生
			Cs134	12.0	Bq/Kg生	± 3.2	Bq/Kg生		Cs134	6.0	Bq/Kg生
Yellow pickled radish	Ohisa, Iwaki	Nov-11	Cs137	12.6	Bq/Kg生	± 3.8	Bq/Kg生	22.9	Cs137	7.0	Bq/Kg生
			Cs134	10.3	Bq/Kg生	± 3.0	Bq/Kg生		Cs134	6.0	Bq/Kg生
Houttuynia cordata(dried)	Satogaoka, Iwaki	Jul-11	Cs137	756	Bq/Kg生	± 159	Bq/Kg生	1,397	Cs137	7.0	Bq/Kg生
			Cs134	641	Bq/Kg生	± 129	Bq/Kg生		Cs134	6.0	Bq/Kg生
Grass	Ono, Tamura	Unknown	Cs137	48.2	Bq/Kg生	± 28.5	Bq/Kg生	80.5	Cs137	8.0	Bq/Kg生
			Cs134	32.3	Bq/Kg生	± 20.3	Bq/Kg生		Cs134	8.0	Bq/Kg生
Grass	Kawamae, Iwaki	Unknown	Cs137	9625	Bq/Kg生	± 936	Bq/Kg生	16,604	Cs137	8.0	Bq/Kg生
			Cs134	6979	Bq/Kg生	± 640	Bq/Kg生		Cs134	8.0	Bq/Kg生
Grass	Kawamae, Iwaki	Unknown	Cs137	1559	Bq/Kg生	± 198	Bq/Kg生	2,808	Cs137	8.0	Bq/Kg生
			Cs134	1249	Bq/Kg生	± 136	Bq/Kg生		Cs134	8.0	Bq/Kg生
Charcoal	Shimookeuri, Kawamae, Iwaki	2011年	Cs137	12840	Bq/Kg生	± 1160	Bq/Kg生	21,060	Cs137	8.0	Bq/Kg生
			Cs134	8220	Bq/Kg生	± 720	Bq/Kg生		Cs134	8.0	Bq/Kg生
Tap water	Taira, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Taira, Iwaki	Mar-12	Cs137	1.2	Bq/Kg生	± 1.2	Bq/Kg生	1.2	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Wakabadai, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	0.0	Cs137	5.0	Bq/Kg生
			Cs134	0.0	Bq/Kg生	± 0.0	Bq/Kg生		Cs134	3.0	Bq/Kg生
Tap water	Kusakidai, Iwaki	Mar-12	Cs137	0.0	Bq/Kg生	± 0.0	Bq/Kg生	1.2	Cs137	5.0	Bq/Kg生
			Cs134	1.2	Bq/Kg生	± 1.0	Bq/Kg生		Cs134	3.0	Bq/Kg生

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Tap water	Sumiyoshi, Onahama, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	6.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	4.0 Bq/Kg生		
Tap water	Tamatsuyu, Izumi, Iwaki	Mar-12	Cs137	2.2 Bq/Kg生	± 1.4 Bq/Kg生	2.2	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Hisanohama, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Filtered water	Hisanohama, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Tap water	Namie, Futaba	Feb-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.4	Cs137	5.0 Bq/Kg生		
			Cs134	1.4 Bq/Kg生	± 1.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Shimookeuri, Kawamae, Iwaki	Mar-12	Cs137	1.7 Bq/Kg生	± 1.2 Bq/Kg生	3.9	Cs137	5.0 Bq/Kg生		
			Cs134	2.2 Bq/Kg生	± 1.1 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Ebata, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Kouya, Uchigo, Iwaki	Mar-12	Cs137	1.2 Bq/Kg生	± 1.2 Bq/Kg生	1.2	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Kamikajiro, Onahama, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.3	Cs137	5.0 Bq/Kg生		
			Cs134	1.3 Bq/Kg生	± 1.1 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Okaona, Onahama, Iwaki	Mar-12	Cs137	2.3 Bq/Kg生	± 1.5 Bq/Kg生	2.3	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Kamikajiro, Onahama, Iwaki	Unknown	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Shimoniida, Yotsukura, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Misawa, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Well water	Nakoso, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.3	Cs137	5.0 Bq/Kg生		
			Cs134	1.3 Bq/Kg生	± 1.1 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Groundwater	Tamura, Koriyama	Feb-12	Cs137	2.1 Bq/Kg生	± 1.3 Bq/Kg生	2.1	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Groundwater	Haragoya, Taira, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Groundwater	Okaona, Onahama, Iwaki	Mar-12	Cs137	2.5 Bq/Kg生	± 1.4 Bq/Kg生	2.5	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Shimookeuri, Kawamae, Iwaki	Oct-11	Cs137	1.7 Bq/Kg生	± 1.3 Bq/Kg生	3.0	Cs137	5.0 Bq/Kg生		
			Cs134	1.3 Bq/Kg生	± 1.2 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Fujiwara, Joban, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.8	Cs137	5.0 Bq/Kg生		
			Cs134	1.8 Bq/Kg生	± 1.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Fujiwara, Joban, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	1.2	Cs137	5.0 Bq/Kg生		
			Cs134	1.2 Bq/Kg生	± 1.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Spring water	Kaminagai, Miwa, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Hot spring water	Kusakidai, Iwaki	Mar-12	Cs137	0.0 Bq/Kg生	± 0.0 Bq/Kg生	0.0	Cs137	5.0 Bq/Kg生		
			Cs134	0.0 Bq/Kg生	± 0.0 Bq/Kg生		Cs134	3.0 Bq/Kg生		
Soil	Nihonmatsu	Mar-12	Cs137	389 Bq/Kg乾	± 47.0 Bq/Kg乾	660	Cs137	8.0 Bq/Kg乾		
			Cs134	271 Bq/Kg乾	± 30.0 Bq/Kg乾		Cs134	8.0 Bq/Kg乾		
Soil	Hirono, Futaba	Mar-12	Cs137	974 Bq/Kg乾	± 99.0 Bq/Kg乾	1,651	Cs137	8.0 Bq/Kg乾		
			Cs134	677 Bq/Kg乾	± 65.0 Bq/Kg乾		Cs134	8.0 Bq/Kg乾		

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Soil	Hirono, Futaba	Mar-12	Cs137	836	Bq/Kg乾	± 88.0	Bq/Kg乾	1,451	Cs137	8.0	Bq/Kg乾
			Cs134	615	Bq/Kg乾	± 60.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Yamada, Ogawa, Iwaki	Mar-12	Cs137	81.0	Bq/Kg乾	± 15.0	Bq/Kg乾	143	Cs137	8.0	Bq/Kg乾
			Cs134	62.0	Bq/Kg乾	± 10.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Ori, Yoshima, Iwaki	Mar-12	Cs137	163	Bq/Kg乾	± 25.0	Bq/Kg乾	288	Cs137	8.0	Bq/Kg乾
			Cs134	125	Bq/Kg乾	± 17.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Ori, Yoshima, Iwaki	Mar-12	Cs137	273	Bq/Kg乾	± 35.0	Bq/Kg乾	458	Cs137	8.0	Bq/Kg乾
			Cs134	185	Bq/Kg乾	± 23.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Shimoogoe, Taira, Iwaki	Mar-12	Cs137	421	Bq/Kg乾	± 49.0	Bq/Kg乾	727	Cs137	8.0	Bq/Kg乾
			Cs134	306	Bq/Kg乾	± 32.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Taira, Iwaki	Mar-12	Cs137	802	Bq/Kg乾	± 85.0	Bq/Kg乾	1,358	Cs137	8.0	Bq/Kg乾
			Cs134	556	Bq/Kg乾	± 56.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Kitashirado, Taira, Iwaki	Mar-12	Cs137	401	Bq/Kg乾	± 49.0	Bq/Kg乾	704	Cs137	8.0	Bq/Kg乾
			Cs134	303	Bq/Kg乾	± 33.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Fujima, Taira, Iwaki	Mar-12	Cs137	151	Bq/Kg乾	± 21.0	Bq/Kg乾	249	Cs137	8.0	Bq/Kg乾
			Cs134	98.0	Bq/Kg乾	± 13.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Iino, Chuodai, Iwaki	Mar-12	Cs137	308	Bq/Kg乾	± 38.0	Bq/Kg乾	531	Cs137	8.0	Bq/Kg乾
			Cs134	223	Bq/Kg乾	± 25.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Satogaoka, Iwaki	Mar-12	Cs137	619	Bq/Kg乾	± 67.0	Bq/Kg乾	1,067	Cs137	8.0	Bq/Kg乾
			Cs134	448	Bq/Kg乾	± 45.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Miya, Uchigo, Iwaki	Mar-12	Cs137	145	Bq/Kg乾	± 22.0	Bq/Kg乾	248	Cs137	8.0	Bq/Kg乾
			Cs134	103	Bq/Kg乾	± 14.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Shimoyunagawa, Joban, Iwaki	Mar-12	Cs137	343	Bq/Kg乾	± 42.0	Bq/Kg乾	578	Cs137	8.0	Bq/Kg乾
			Cs134	235	Bq/Kg乾	± 26.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Shimoyunagawa, Joban, Iwaki	Mar-12	Cs137	289	Bq/Kg乾	± 36.0	Bq/Kg乾	486	Cs137	8.0	Bq/Kg乾
			Cs134	197	Bq/Kg乾	± 23.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Onahama, Iwaki	Mar-12	Cs137	122	Bq/Kg乾	± 19.0	Bq/Kg乾	211	Cs137	8.0	Bq/Kg乾
			Cs134	89.0	Bq/Kg乾	± 13.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	33.0	Bq/Kg乾	± 9.0	Bq/Kg乾	59.0	Cs137	8.0	Bq/Kg乾
			Cs134	26.0	Bq/Kg乾	± 7.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	337	Bq/Kg乾	± 41.0	Bq/Kg乾	574	Cs137	8.0	Bq/Kg乾
			Cs134	237	Bq/Kg乾	± 27.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Okaona, Onahama, Iwaki	Mar-12	Cs137	95.0	Bq/Kg乾	± 17.0	Bq/Kg乾	167	Cs137	8.0	Bq/Kg乾
			Cs134	72.0	Bq/Kg乾	± 11.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Okaona, Onahama, Iwaki	Mar-12	Cs137	479	Bq/Kg乾	± 56.0	Bq/Kg乾	835	Cs137	8.0	Bq/Kg乾
			Cs134	356	Bq/Kg乾	± 38.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Okaona, Onahama, Iwaki	Mar-12	Cs137	520	Bq/Kg乾	± 59.0	Bq/Kg乾	867	Cs137	8.0	Bq/Kg乾
			Cs134	347	Bq/Kg乾	± 37.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Shiroyone, Nakoso, Iwaki	Mar-12	Cs137	166	Bq/Kg乾	± 23.0	Bq/Kg乾	281	Cs137	8.0	Bq/Kg乾
			Cs134	115	Bq/Kg乾	± 15.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Soil	Sekita, Nakoso, Iwaki	Mar-12	Cs137	47.0	Bq/Kg乾	± 12.0	Bq/Kg乾	80.0	Cs137	8.0	Bq/Kg乾
			Cs134	33.0	Bq/Kg乾	± 8.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Ogawa, Iwaki	Mar-12	Cs137	70.0	Bq/Kg乾	± 14.0	Bq/Kg乾	123	Cs137	8.0	Bq/Kg乾
			Cs134	53.0	Bq/Kg乾	± 10.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Shimoarakawa, Taira, Iwaki	Mar-12	Cs137	368	Bq/Kg乾	± 46.0	Bq/Kg乾	669	Cs137	8.0	Bq/Kg乾
			Cs134	301	Bq/Kg乾	± 33.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Shimotakaku, Taira, Iwaki	Mar-12	Cs137	357	Bq/Kg乾	± 44.0	Bq/Kg乾	608	Cs137	8.0	Bq/Kg乾
			Cs134	251	Bq/Kg乾	± 28.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
			Cs137	Cs134	±	±		Cs137	Cs134		
Field soil	Iino, Chuodai, Iwaki	Mar-12	Cs137	146	Bq/Kg乾	± 23.0	Bq/Kg乾	254	Cs137	8.0	Bq/Kg乾
			Cs134	108	Bq/Kg乾	± 16.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Shimoyunagaya, Joban, Iwaki	Mar-12	Cs137	247	Bq/Kg乾	± 33.0	Bq/Kg乾	419	Cs137	8.0	Bq/Kg乾
			Cs134	172	Bq/Kg乾	± 21.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Kamiyunagaya, Joban, Iwaki	Mar-12	Cs137	277	Bq/Kg乾	± 36.0	Bq/Kg乾	488	Cs137	8.0	Bq/Kg乾
			Cs134	211	Bq/Kg乾	± 24.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Kamikajiro, Onahama, Iwaki	Mar-12	Cs137	293	Bq/Kg乾	± 38.0	Bq/Kg乾	502	Cs137	8.0	Bq/Kg乾
			Cs134	209	Bq/Kg乾	± 25.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	492	Bq/Kg乾	± 56.0	Bq/Kg乾	885	Cs137	8.0	Bq/Kg乾
			Cs134	393	Bq/Kg乾	± 40.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	261	Bq/Kg乾	± 35.0	Bq/Kg乾	458	Cs137	8.0	Bq/Kg乾
			Cs134	197	Bq/Kg乾	± 24.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Nagasaki, Iwaki	Mar-12	Cs137	114	Bq/Kg乾	± 18.0	Bq/Kg乾	191	Cs137	8.0	Bq/Kg乾
			Cs134	77.0	Bq/Kg乾	± 12.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Field soil	Nishiki, Iwaki	Mar-12	Cs137	413	Bq/Kg乾	± 49.0	Bq/Kg乾	727	Cs137	8.0	Bq/Kg乾
			Cs134	314	Bq/Kg乾	± 33.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Rice field soil	Kamimosu, Yamadama, Iwaki	Mar-12	Cs137	358	Bq/Kg乾	± 42.0	Bq/Kg乾	608	Cs137	8.0	Bq/Kg乾
			Cs134	250	Bq/Kg乾	± 27.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Rice field soil	Kamimosu, Yamadama, Iwaki	Mar-12	Cs137	387	Bq/Kg乾	± 45.0	Bq/Kg乾	612	Cs137	8.0	Bq/Kg乾
			Cs134	225	Bq/Kg乾	± 26.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Garden soil	Kitashirado, Taira, Iwaki	Mar-12	Cs137	1043	Bq/Kg乾	± 107	Bq/Kg乾	1,795	Cs137	8.0	Bq/Kg乾
			Cs134	752	Bq/Kg乾	± 72.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Garden soil	Iino, Chuodai, Iwaki	Mar-12	Cs137	290	Bq/Kg乾	± 36.0	Bq/Kg乾	500	Cs137	8.0	Bq/Kg乾
			Cs134	210	Bq/Kg乾	± 24.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Garden soil	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	237	Bq/Kg乾	± 31.0	Bq/Kg乾	408	Cs137	8.0	Bq/Kg乾
			Cs134	171	Bq/Kg乾	± 21.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Turf grass/soil	Kouya, Uchigo, Iwaki	Mar-12	Cs137	2400	Bq/Kg乾	± 240	Bq/Kg乾	4,100	Cs137	8.0	Bq/Kg乾
			Cs134	1700	Bq/Kg乾	± 160	Bq/Kg乾		Cs134	8.0	Bq/Kg乾
Turf grass	Kouya, Uchigo, Iwaki	Mar-12	Cs137	620	Bq/Kg乾	± 70.0	Bq/Kg乾	1,060	Cs137	8.0	Bq/Kg乾
			Cs134	440	Bq/Kg乾	± 45.0	Bq/Kg乾		Cs134	8.0	Bq/Kg乾