



Radiation Measurement Results of 522 Items in April 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Miyazaki	Oct-11	Cs137	0.0	Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Fukushima	2010	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Maenarusawa, Fukushima	Oct-11	Cs137	0.0	Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Matsukawa, Fukushima	Oct-11	Cs137	6.6	Bq/Kg raw	15.6	Cs137	7.0 Bq/Kg raw
			Cs134	9.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Minamiaizu	Oct-11	Cs137	0.0	Bq/Kg raw	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Aizutadami	Nov-11	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Tanakura, Higashishirakawa	Oct-11	Cs137	3.2	Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Date, Date	Oct-11	Cs137	0.0	Bq/Kg raw	5.8	Cs137	7.0 Bq/Kg raw
			Cs134	5.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Nakajima, Nametsu, Shirakawa	Oct-11	Cs137	4.5	Bq/Kg raw	7.1	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Otama, Adachi	Unknown	Cs137	2.2	Bq/Kg raw	4.4	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Nihonmatsu	Unknown	Cs137	4.3	Bq/Kg raw	8.3	Cs137	7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Mihota, Koriyama	Oct-11	Cs137	0.0	Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Sukagawa	Oct-11	Cs137	6.1	Bq/Kg raw	12.0	Cs137	7.0 Bq/Kg raw
			Cs134	5.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Furudono, Ishikawa	Oct-11	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Furudono, Ishikawa	Oct-11	Cs137	0.0	Bq/Kg raw	2.4	Cs137	7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Hiwatashi, Namie, Futaba	Oct-10	Cs137	0.0	Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kamiogawa, Ogawa, Iwaki	Oct-11	Cs137	3.5	Bq/Kg raw	7.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Akai, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Akai, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.0	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Kamiyamaguchi, Taira, Iwaki	Oct-11	Cs137	4.5	Bq/Kg raw	± 2.0	Bq/Kg raw	7.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 1.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Kabeya, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yoshima, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Natsui, Iwaki	Oct-11	Cs137	6.5	Bq/Kg raw	± 2.5	Bq/Kg raw	6.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Natsui, Iwaki	Oct-11	Cs137	5.7	Bq/Kg raw	± 2.4	Bq/Kg raw	7.9	Cs137 7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Numanouchi, Iwaki	Oct-11	Cs137	4.1	Bq/Kg raw	± 2.7	Bq/Kg raw	8.3	Cs137 7.0 Bq/Kg raw
			Cs134	4.2	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Nagatomo, Yotsukura, Iwaki	Oct-11	Cs137	3.0	Bq/Kg raw	± 2.5	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Nishiki, Iwaki	Oct-11	Cs137	4.3	Bq/Kg raw	± 2.5	Bq/Kg raw	7.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Iritono, Tono, Iwaki	Oct-11	Cs137	4.6	Bq/Kg raw	± 2.5	Bq/Kg raw	4.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamada, Iwaki	Oct-11	Cs137	9.5	Bq/Kg raw	± 3.0	Bq/Kg raw	19.4	Cs137 6.0 Bq/Kg raw
			Cs134	9.9	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Brown rice	Yamada, Iwaki	Unknown	Cs137	4.0	Bq/Kg raw	± 2.2	Bq/Kg raw	4.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Akita	Oct-11	Cs137	3.2	Bq/Kg raw	± 4.4	Bq/Kg raw	4.5	Cs137 8.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 4.2	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
White rice	Daisen, Akita	Oct-11	Cs137	2.6	Bq/Kg raw	± 2.4	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Omagari, Akita	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamagata	Oct-01	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamagata	Oct-11	Cs137	2.5	Bq/Kg raw	± 2.2	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Shiroishi, Miyagi	Oct-11	Cs137	2.1	Bq/Kg raw	± 2.0	Bq/Kg raw	4.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kurokawa, Miyagi	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.9	Cs137 7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Takakiyo, Kurihara, Miyagi	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.8	Cs137 7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Takakura, Kakuda, Miyagi	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kokunou, Katori, Chiba	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.3	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Morido, Katori, Chiba	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Niigata	Unknown	Cs137	0.0	Bq/Kg raw	± 2.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Niigata	Oct-11	Cs137	2.2	Bq/Kg raw	± 2.2	Bq/Kg raw	2.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Juou, Hitachi, Ibaraki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.3	Cs137 7.0 Bq/Kg raw
			Cs134	4.3	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ibaraki	Oct-11	Cs137	4.3	Bq/Kg raw	± 2.2	Bq/Kg raw	6.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Aizumisato, Onuma	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.8	Cs137 7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Aizuwakamatsu	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Koriyama	Oct-11	Cs137	2.6	Bq/Kg raw	± 2.3	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tamura, Koriyama	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.8	Cs137 7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nakata, Koriyama	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Outama, Adachi	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Outama, Adachi	Oct-11	Cs137	3.9	Bq/Kg raw	± 2.1	Bq/Kg raw	3.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Funahiki, Tamura	Oct-11	Cs137	2.7	Bq/Kg raw	± 2.4	Bq/Kg raw	5.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Taguchi, Furudono, Ishikawa	Oct-11	Cs137	2.4	Bq/Kg raw	± 2.1	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ishikawa, Ishikawa	Oct-11	Cs137	2.6	Bq/Kg raw	± 2.1	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ishikawa, Ishikawa	Nov-11	Cs137	3.0	Bq/Kg raw	± 2.0	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soma	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Saiso, Miwa, Iwaki	Oct-11	Cs137	3.2	Bq/Kg raw	± 2.1	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Miwa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ogawa, Iwaki	Oct-11	Cs137	2.9	Bq/Kg raw	± 2.1	Bq/Kg raw	5.5	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ogawa, Iwaki	Oct-11	Cs137	2.8	Bq/Kg raw	± 2.6	Bq/Kg raw	8.8	Cs137 7.0 Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ogawa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.7	Cs137 7.0 Bq/Kg raw
			Cs134	4.7	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
White rice	Yotsukura, Iwaki	Oct-11	Cs137	3.2	Bq/Kg raw	± 2.2	Bq/Kg raw	5.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	2.6	Bq/Kg raw	± 2.2	Bq/Kg raw	6.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	4.0	Bq/Kg raw	± 2.3	Bq/Kg raw	7.7	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Hosoya, Yotsukura, Iwaki	Oct-11	Cs137	4.3	Bq/Kg raw	± 2.5	Bq/Kg raw	7.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Toda, Yotsukura, Iwaki	Oct-11	Cs137	3.2	Bq/Kg raw	± 1.7	Bq/Kg raw	5.6	Cs137	6.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.5	Bq/Kg raw		Cs134	4.0 Bq/Kg raw
White rice	Yagyu, Yotsukura, Iwaki	Oct-11	Cs137	3.5	Bq/Kg raw	± 2.5	Bq/Kg raw	7.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Kusano, Iwaki	Oct-11	Cs137	4.1	Bq/Kg raw	± 2.5	Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yoshima, Iwaki	Oct-11	Cs137	2.9	Bq/Kg raw	± 2.9	Bq/Kg raw	5.1	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Akai, Taira, Iwaki	Nov-11	Cs137	3.4	Bq/Kg raw	± 2.1	Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Akai, Taira, Iwaki	Sep-11	Cs137	5.2	Bq/Kg raw	± 2.4	Bq/Kg raw	5.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Kinuya, Taira, Iwaki	Nov-12	Cs137	2.7	Bq/Kg raw	± 2.2	Bq/Kg raw	5.3	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yoshinoya, Taira, Iwaki	Oct-11	Cs137	2.3	Bq/Kg raw	± 2.2	Bq/Kg raw	4.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Kitashirado, Taira, Iwaki	Oct-11	Cs137	2.8	Bq/Kg raw	± 1.9	Bq/Kg raw	5.9	Cs137	7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Takaku, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	3.6	Bq/Kg raw	± 2.1	Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Shimohirakubo, Taira, Iwaki	Oct-11	Cs137	3.5	Bq/Kg raw	± 2.1	Bq/Kg raw	5.4	Cs137	7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Shimohirakubo, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg raw	± 2.1	Bq/Kg raw	5.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Mizushina, Taira, Iwaki	Oct-11	Cs137	2.6	Bq/Kg raw	± 2.3	Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg raw	± 2.4	Bq/Kg raw	6.1	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Shimoogoe, Taira, Iwaki	Oct-11	Cs137	5.0	Bq/Kg raw	± 2.5	Bq/Kg raw	8.8	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Kujiraoka, Taira, Iwaki	Oct-11	Cs137	2.9	Bq/Kg raw	± 2.1	Bq/Kg raw	5.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Takasaka, Uchigo, Iwaki	Oct-11	Cs137	2.6	Bq/Kg raw	± 2.1	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kashima, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kashima, Iwaki	Oct-11	Cs137	2.7	Bq/Kg raw	± 4.4	Bq/Kg raw	5.0	Cs137 8.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 4.1	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
White rice	Sumiyoshi, Onahama, Iwaki	Nov-11	Cs137	3.9	Bq/Kg raw	± 2.5	Bq/Kg raw	7.9	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nishigo, Joban, Iwaki	Oct-11	Cs137	3.7	Bq/Kg raw	± 2.2	Bq/Kg raw	6.5	Cs137 7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nishigo, Joban, Iwaki	Nov-11	Cs137	3.0	Bq/Kg raw	± 2.2	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	3.2	Bq/Kg raw	± 2.5	Bq/Kg raw	7.2	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	4.2	Bq/Kg raw	± 2.2	Bq/Kg raw	4.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Fujiwara, Joban, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Shiratori, Joban, Iwaki	Oct-11	Cs137	3.1	Bq/Kg raw	± 2.1	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kadono, Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamada, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Misawa, Iwaki	Oct-11	Cs137	3.9	Bq/Kg raw	± 2.7	Bq/Kg raw	3.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ebata, Ueda, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.5	Cs137 6.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 1.5	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
White rice	Watanabe, Iwaki	Oct-11	Cs137	2.5	Bq/Kg raw	± 2.4	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Watanabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tamachi, Nakoso, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.3	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Iwaki	Unknown	Cs137	2.6	Bq/Kg raw	± 2.2	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Unknown	Unknown	Cs137	3.1	Bq/Kg raw	± 2.0	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sticky rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sticky rice	Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.1	Cs137 7.0 Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sticky rice	Watanabe, Iwaki	Oct-11	Cs137	3.9	Bq/Kg raw	± 2.3	Bq/Kg raw	3.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Sticky rice (brown)	Inawashiro, Yama	Oct-11	Cs137	2.2	Bq/Kg raw	± 1.8	2.2	Cs137	6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	4.0 Bq/Kg raw
Rice (Half-milled)	Yotsukura, Iwaki	Oct-11	Cs137	6.7	Bq/Kg raw	± 3.0	10.9	Cs137	7.0 Bq/Kg raw
			Cs134	4.2	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Fried rice	Toyotomi, Kakuda, Miyagi	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
Fried rice	Toyotomi, Kakuda, Miyagi	Oct-11	Cs137	2.8	Bq/Kg raw	± 2.1	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Fried rice	Inaoki, Kakuda, Miyagi	Oct-11	Cs137	1.9	Bq/Kg raw	± 2.1	4.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Fried rice	Shimoyagyu, Yotsukura, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
Potatoes	Taira, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.4		Cs134	6.0 Bq/Kg raw
Potatoes	Shirado, Taira, Iwaki	May-11	Cs137	0.0	Bq/Kg raw	± 0.0	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.9		Cs134	6.0 Bq/Kg raw
Potatoes	Kitashirado, Taira, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Shimoogoe, Taira, Iwaki	Aug-11	Cs137	0.0	Bq/Kg raw	± 0.0	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.5		Cs134	6.0 Bq/Kg raw
Potatoes	Akai, Taira, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.2		Cs134	6.0 Bq/Kg raw
Potatoes	Kamata, Taira, Iwaki	Jul-11	Cs137	5.2	Bq/Kg raw	± 2.7	5.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Nakashio, Taira, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Ishimori, Taira, Iwaki	Jun-11	Cs137	4.1	Bq/Kg raw	± 2.7	7.9	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.4		Cs134	6.0 Bq/Kg raw
Potatoes	Nakakabeya, Taira, Iwaki	Aug-11	Cs137	4.4	Bq/Kg raw	± 2.2	8.1	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
Potatoes	Shimoyoshima, Yoshima, Iwaki	Jul-11	Cs137	1.9	Bq/Kg raw	± 1.4	3.3	Cs137	5.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.2		Cs134	3.0 Bq/Kg raw
Potatoes	Mimaya, Uchigo, Iwaki	Jul-11	Cs137	2.7	Bq/Kg raw	± 2.4	6.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Potatoes	Miya, Uchigo, Iwaki	Aug-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Misawa, Joban, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.3		Cs134	6.0 Bq/Kg raw
Potatoes	Onahamashima, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.3	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Potatoes	Okaona, Onahama, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Tamagawa, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Shimokawa, Izumi, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Potatoes	Kamikamado, Watanabe, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potatoes	Miyamada, Tono, Iwaki	Aug-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.7	Cs137 7.0 Bq/Kg raw
			Cs134	4.7	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potatoes	Sakai, Nakoso, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potatoes	Ibaraki	Unknown	Cs137	7.5	Bq/Kg raw	± 3.0	Bq/Kg raw	10.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potatoes	Numabe, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic	Shibahara, Ogawa, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic	Ori, Yoshima, Iwaki	Jun-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	7.7	Cs137 8.0 Bq/Kg raw
			Cs134	7.7	Bq/Kg raw	± 5.4	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Chinese yams	Kashima, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Yams	Tamagawa, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Konjak yams	Kuroda, Tabito, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.3	Cs137 7.0 Bq/Kg raw
			Cs134	4.3	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Arios	Oseki, Yoshima, Iwaki	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Yacon	Toyoma, Taira, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137 7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Burdock roots	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	2.0	Bq/Kg raw	± 2.5	Bq/Kg raw	10.4	Cs137 8.0 Bq/Kg raw
			Cs134	8.4	Bq/Kg raw	± 6.0	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Japanese radish	Moegidai, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Moegidai, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Tamagawa, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Mimaya, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.8	Cs137 7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Akai, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Iino, Chuodai, Iwaki	Mar-12	Cs137	2.6	Bq/Kg raw	± 2.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Tamagawa, Onahama, Iwaki	Mar-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks, Japanese radish & potatoes	Sumiyoshi, Onahama, Iwaki	Unknown	Cs137	7.1	Bq/Kg raw	± 5.0	Bq/Kg raw	7.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Dried Japanese radish	Odaka, Nakoso, Iwaki	Jan-12	Cs137	18.9	Bq/Kg raw	± 9.8	Bq/Kg raw	35.7	Cs137 7.0 Bq/Kg raw
			Cs134	16.8	Bq/Kg raw	± 8.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish leaves	Nagasaki, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Carrots	Shimohirakubo, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.4	Cs137 7.0 Bq/Kg raw
			Cs134	5.4	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Carrots	Mimaya, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shallots	Jigiri, nagasaki, Iwaki	Apr-12	Cs137	8.9	Bq/Kg raw	± 5.0	Bq/Kg raw	15.8	Cs137 7.0 Bq/Kg raw
			Cs134	6.9	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onions	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onions	Kitashirado, Taira, Iwaki	Jul-11	Cs137	5.6	Bq/Kg raw	± 3.2	Bq/Kg raw	5.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onions	Ori, Yoshima, Iwaki	Jun-11	Cs137	2.8	Bq/Kg raw	± 4.3	Bq/Kg raw	7.0	Cs137 8.0 Bq/Kg raw
			Cs134	4.2	Bq/Kg raw	± 1.3	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Spring onions	Numanouchi, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Spring onions	Shimoogoe, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137 7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Spring onions	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Spring onions	Kamiyunagaya, Joban, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137 7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Spring onions	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.6	Cs137 7.0 Bq/Kg raw
			Cs134	4.6	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Spring onions	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137 8.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 4.8	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Leeks	Kamata, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Kamiyamaguchi, Taira, Iwaki	Apr-12	Cs137	5.5	Bq/Kg raw	± 3.4	Bq/Kg raw	5.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Shimoogoe, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.5	Cs137 7.0 Bq/Kg raw
			Cs134	4.5	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	6.4	Bq/Kg raw	± 3.3	Bq/Kg raw	12.8	Cs137 7.0 Bq/Kg raw
			Cs134	6.4	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Takahagi, Ogawa, Iwaki	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Shibahara, Ogawa, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Koya, Uchigo, Iwaki	Apr-12	Cs137	5.9	Bq/Kg raw	± 4.0	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Miya, Uchigo, Iwaki	Apr-12	Cs137	5.6	Bq/Kg raw	± 3.3	Bq/Kg raw	5.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Nagasaki, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Onahamashima, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Kamiyada, Joban, Iwaki	Apr-12	Cs137	3.8	Bq/Kg raw	± 2.7	Bq/Kg raw	3.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Leeks	Moegidai, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Kadono, Tono, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Leeks	Tanabe, Watanabe, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leeks	Numabe, Iwaki	Apr-12	Cs137	2.5	Bq/Kg raw	± 2.0	Bq/Kg raw	6.5	Cs137 8.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 3.8	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Leeks	Yamada, Iwaki	Apr-12	Cs137	0.3	Bq/Kg raw	± 0.0	Bq/Kg raw	5.0	Cs137 8.0 Bq/Kg raw
			Cs134	4.7	Bq/Kg raw	± 3.8	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Garlic chives	Fujima, Taira, Iwaki	Apr-12	Cs137	4.2	Bq/Kg raw	± 4.0	Bq/Kg raw	8.0	Cs137 7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic chives	Shimoarakawa, Taira, Iwaki	Apr-12	Cs137	4.9	Bq/Kg raw	± 4.5	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic chives	Shimoogoe, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.5	Cs137 7.0 Bq/Kg raw
			Cs134	5.5	Bq/Kg raw	± 3.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic chives	Shimoyamaguchi, Taiara, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic chives	Iida, Kashima, Iwaki	Apr-12	Cs137	25.8	Bq/Kg raw	± 8.1	Bq/Kg raw	41.3	Cs137 7.0 Bq/Kg raw
			Cs134	15.5	Bq/Kg raw	± 5.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic chives	Sumiyoshi, Onahama, Iwaki	Apr-12	Cs137	7.4	Bq/Kg raw	± 5.0	Bq/Kg raw	18.6	Cs137 7.0 Bq/Kg raw
			Cs134	11.2	Bq/Kg raw	± 4.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Garlic chives	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.8	Cs137 7.0 Bq/Kg raw
			Cs134	4.8	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shallot	Nishiki, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shallot	Shimoarakawa, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chives	Kashima, Chuodai, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	6.6	Cs137 7.0 Bq/Kg raw
			Cs134	6.6	Bq/Kg raw	± 6.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shallots	Kashima, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.7	Cs137 7.0 Bq/Kg raw
			Cs134	5.7	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shallots	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Azuki beans	Obihiro, Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Black beans	Nichibu, Tabito, Iwaki	Dec-11	Cs137	4.4	Bq/Kg raw	± 2.7	Bq/Kg raw	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbages	Miyama, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.1	Cs137 8.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 4.5	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Chinese cabbages	Yokodai, Iwaki	Apr-12	Cs137	5.7	Bq/Kg raw	± 4.8	Bq/Kg raw	10.6	Cs137 7.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 3.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbages	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	5.5	Bq/Kg raw	± 3.0	Bq/Kg raw	5.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbages	Kamiyamaguchi, Taira, Iwaki	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Chinese cabbages	Moegidai, Izumi, Iwaki	Feb-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.3	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbages	Sumiyoshi, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.9	Cs137 7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbages	Miwa, Iwaki	Sep-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Aichi	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Nakashio, Taira, Iwaki	Apr-12	Cs137	3.0	Bq/Kg raw	± 2.8	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Mizunoya, Joban, Iwaki	Apr-12	Cs137	7.0	Bq/Kg raw	± 6.5	Bq/Kg raw	7.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	6.6	Cs137 7.0 Bq/Kg raw
			Cs134	6.6	Bq/Kg raw	± 3.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Shimokawa, Izumi, Iwaki	Apr-12	Cs137	3.3	Bq/Kg raw	± 3.3	Bq/Kg raw	7.7	Cs137 7.0 Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Moegidai, Izumi, Iwaki	Apr-12	Cs137	3.0	Bq/Kg raw	± 2.4	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Otani, Yamada, Iwaki	Apr-12	Cs137	3.6	Bq/Kg raw	± 3.1	Bq/Kg raw	3.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cabbages	Hanawa, Watanabe, Iwaki	Apr-12	Cs137	4.2	Bq/Kg raw	± 2.4	Bq/Kg raw	4.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Natsui, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	5.9	Bq/Kg raw	± 3.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	4.9	Bq/Kg raw	± 0.4	Bq/Kg raw	8.1	Cs137 8.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 8.2	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Broccoli	Kinuya, Taira, Iwaki	Apr-12	Cs137	3.8	Bq/Kg raw	± 3.4	Bq/Kg raw	8.7	Cs137 6.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Broccoli	Fukimatsu, Onahama, Iwaki	Mar-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.0	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Moegidai, Izumi, Iwaki	Apr-12	Cs137	6.3	Bq/Kg raw	± 3.7	Bq/Kg raw	6.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Shimokawa, Izumi, Iwaki	Mar-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Hanawa, Watanabe, Iwaki	Apr-12	Cs137	3.5	Bq/Kg raw	± 3.4	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Misawa, Iwaki	Apr-12	Cs137	8.4	Bq/Kg raw	± 7.5	Bq/Kg raw	16.9	Cs137 7.0 Bq/Kg raw
			Cs134	8.5	Bq/Kg raw	± 6.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Asparagus	Shimookeuri, Kawamae, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Canola flower	Sakuragaoka, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Canola flower	Sakuragaoka, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Canola flower	Ozeki, Kamiyoshima, Iwaki	Apr-12	Cs137	4.6	Bq/Kg raw	± 2.6	Bq/Kg raw	4.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Canola flower	Moegidai, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Shimohirakubo, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Kamiyamaguchi, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Numanouchi, Taira, Iwaki	Apr-12	Cs137	4.0	Bq/Kg raw	± 2.6	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.3	Bq/Kg raw	
Spinach	Jiyugaoka, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	5.6	Bq/Kg raw	± 4.4	Bq/Kg raw	
Spinach	Shimoyoshima, Yoshima, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 3.2	Bq/Kg raw	
Spinach	Shimoyoshima, Yoshima, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Koya, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	6.7	Bq/Kg raw	± 6.6	Bq/Kg raw	
Spinach	Iida, Kashima, Iwaki	Apr-12	Cs137	7.2	Bq/Kg raw	± 5.3	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	6.7	Bq/Kg raw	± 4.5	Bq/Kg raw	
Spinach	Ohara, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Tamagawa, Onahama, Iwaki	Apr-12	Cs137	5.1	Bq/Kg raw	± 5.1	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Okaona, Onahama, Iwaki	Apr-12	Cs137	6.5	Bq/Kg raw	± 7.4	Bq/Kg raw	Cs137 8.0 Bq/Kg raw Cs134 8.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 5.9	Bq/Kg raw	
Spinach	Tamagawa, Onahama, Iwaki	Apr-12	Cs137	5.3	Bq/Kg raw	± 3.8	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	5.8	Bq/Kg raw	± 4.4	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Moegidai, Izumi, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Spinach	Yamada, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Canola flower	Okaona, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	5.7	Bq/Kg raw	± 3.3	Bq/Kg raw	
Canola flower	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	13.9	Bq/Kg raw	± 6.5	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	10.7	Bq/Kg raw	± 5.0	Bq/Kg raw	
Canola flower	Shioki, Yotsukura, Iwaki	Apr-12	Cs137	11.7	Bq/Kg raw	± 4.9	Bq/Kg raw	Cs137 6.0 Bq/Kg raw Cs134 4.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 3.8	Bq/Kg raw	
Canola flower	Kamiyunagaya, Joban, Iwaki	Apr-12	Cs137	7.2	Bq/Kg raw	± 4.6	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Canola flower	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	3.3	Bq/Kg raw	± 2.5	Bq/Kg raw	Cs137 5.0 Bq/Kg raw Cs134 3.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.4	Bq/Kg raw	
Canola flower	Okaona, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	7.5	Bq/Kg raw	± 3.8	Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Canola flower	Shimokawa, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	12.2	Cs137	7.0 Bq/Kg raw
			Cs134	12.2	Bq/Kg raw	± 6.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Mizuna	Miyamada, Tono, Iwaki	Apr-12	Cs137	3.4	Bq/Kg raw	± 3.1	Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Leaf mustard	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	5.7	Bq/Kg raw	± 14.5	Bq/Kg raw	9.3	Cs137	8.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 4.0	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Leaf mustard	Jiyugaoka, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Leaf mustard	Kitayoshima, Yoshima, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	7.6	Cs137	7.0 Bq/Kg raw
			Cs134	7.6	Bq/Kg raw	± 3.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Leaf mustard	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	2.4	Bq/Kg raw	± 2.2	Bq/Kg raw	2.4	Cs137	6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0 Bq/Kg raw
Leaf mustard	Taki, Tono, Iwaki	Apr-12	Cs137	4.7	Bq/Kg raw	± 3.1	Bq/Kg raw	9.0	Cs137	7.0 Bq/Kg raw
			Cs134	4.3	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Leaf mustard	Kubota, Nakoso, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shungiku	Shimohirakubo, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	6.7	Cs137	7.0 Bq/Kg raw
			Cs134	6.7	Bq/Kg raw	± 4.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shungiku	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	7.4	Cs137	7.0 Bq/Kg raw
			Cs134	7.4	Bq/Kg raw	± 5.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Kakina	Fudaba, Takakura, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Oishina	Kaneyama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Turnips	Koya, Uchigo, Iwaki	Apr-12	Cs137	4.6	Bq/Kg raw	± 3.7	Bq/Kg raw	8.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Turnips	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.1	Cs137	7.0 Bq/Kg raw
			Cs134	5.1	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Turnip Leaves	Nishiki, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Wasabi leaves	Shioki, Yotsukura, Iwaki	Apr-12	Cs137	17.6	Bq/Kg raw	± 11.6	Bq/Kg raw	29.7	Cs137	8.0 Bq/Kg raw
			Cs134	12.1	Bq/Kg raw	± 7.1	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Norabona	Nakashio, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese mustard spinach	Jigiri, Nagasaki, Iwaki	Apr-12	Cs137	7.4	Bq/Kg raw	± 6.8	Bq/Kg raw	18.6	Cs137	7.0 Bq/Kg raw
			Cs134	11.2	Bq/Kg raw	± 5.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese mustard spinach	Taki, Tono, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.0	Cs137	7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese mustard spinach	Shimokawa, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese mustard spinach	Kamiyada, Joban, Iwaki	Apr-12	Cs137	7.4	Bq/Kg raw	± 4.7	Bq/Kg raw	7.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Watercress	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	8.6	Bq/Kg raw	± 4.8	Bq/Kg raw	16.8	Cs137	7.0 Bq/Kg raw
			Cs134	8.2	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese parsley	Shimofunao, Joban, Iwaki	Apr-12	Cs137	10.3	Bq/Kg raw	± 5.0	Bq/Kg raw	22.1	Cs137	7.0 Bq/Kg raw
			Cs134	11.8	Bq/Kg raw	± 4.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Japanese parsley	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	13.3 Bq/Kg raw	± 5.7 Bq/Kg raw	18.3	Cs137	7.0 Bq/Kg raw
			Cs134	5.0 Bq/Kg raw	± 4.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Mitsuba	Yamada, Iwaki	Apr-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Greens	Kitayoshima, Yoshima, Iwaki	Apr-12	Cs137	5.9 Bq/Kg raw	± 4.7 Bq/Kg raw	5.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Tsubomina	Shimoogoe, Taira, Iwaki	Apr-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Tsubomina	Koya, Uchigo, Iwaki	Apr-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	7.4	Cs137	7.0 Bq/Kg raw
			Cs134	7.4 Bq/Kg raw	± 5.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Akai, Taira, Iwaki	Apr-12	Cs137	6.8 Bq/Kg raw	± 3.0 Bq/Kg raw	10.9	Cs137	7.0 Bq/Kg raw
			Cs134	4.1 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Shimoogoe, Taira, Iwaki	Apr-11	Cs137	9.9 Bq/Kg raw	± 4.2 Bq/Kg raw	15.8	Cs137	7.0 Bq/Kg raw
			Cs134	5.9 Bq/Kg raw	± 3.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	8.2	Cs137	7.0 Bq/Kg raw
			Cs134	8.2 Bq/Kg raw	± 4.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kubosorimachi, Kashima, Iwaki	Apr-12	Cs137	7.5 Bq/Kg raw	± 4.5 Bq/Kg raw	13.6	Cs137	7.0 Bq/Kg raw
			Cs134	6.1 Bq/Kg raw	± 3.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kumabashiri, Kashima, Iwaki	Apr-12	Cs137	6.2 Bq/Kg raw	± 3.3 Bq/Kg raw	13.7	Cs137	7.0 Bq/Kg raw
			Cs134	7.5 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	3.8 Bq/Kg raw	± 3.6 Bq/Kg raw	7.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.8 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kanari, Onahama, Iwaki	Apr-12	Cs137	6.0 Bq/Kg raw	± 3.6 Bq/Kg raw	11.1	Cs137	7.0 Bq/Kg raw
			Cs134	5.1 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Okaona, Onahama, Iwaki	Apr-12	Cs137	6.1 Bq/Kg raw	± 4.9 Bq/Kg raw	12.3	Cs137	7.0 Bq/Kg raw
			Cs134	6.2 Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Fujiwara, Joban, Iwaki	Apr-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	3.4 Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kamiyada, Joban, Iwaki	Apr-12	Cs137	12.2 Bq/Kg raw	± 4.3 Bq/Kg raw	15.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.0 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Shimoyunagaya, Joban, Iwaki	Apr-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	5.3 Bq/Kg raw	± 3.0 Bq/Kg raw	8.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.9 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Soeno, Iwaki	Apr-12	Cs137	2.7 Bq/Kg raw	± 2.6 Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw
			Cs134	4.2 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur	Kaneyama, Iwaki	Apr-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.0 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Butterbur Sprout	Okaona, Onahama, Iwaki	Apr-12	Cs137	12.4 Bq/Kg raw	± 16.6 Bq/Kg raw	24.9	Cs137	8.0 Bq/Kg raw
			Cs134	12.5 Bq/Kg raw	± 10.6 Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Fatsia sprouts	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	9.4 Bq/Kg raw	± 4.5 Bq/Kg raw	19.8	Cs137	7.0 Bq/Kg raw
			Cs134	10.4 Bq/Kg raw	± 3.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Fatsia sprouts	Okaona, Onahama, Iwaki	Apr-12	Cs137	155 Bq/Kg raw	± 31.6 Bq/Kg raw	245	Cs137	8.0 Bq/Kg raw
			Cs134	90.3 Bq/Kg raw	± 17.3 Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Fatsia sprouts	Onahama, Iwaki	Apr-12	Cs137	8.1 Bq/Kg raw	± 8.9 Bq/Kg raw	33.2	Cs137	8.0 Bq/Kg raw
			Cs134	25.1 Bq/Kg raw	± 9.0 Bq/Kg raw		Cs134	8.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Fatsia sprouts	Okaona, Onahama, Iwaki	Apr-12	Cs137	8.3	Bq/Kg raw	± 4.3	Bq/Kg raw	13.8	Cs137 7.0 Bq/Kg raw
			Cs134	5.5	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Fatsia sprouts	Shimokawa, Izumi, Iwaki	Apr-12	Cs137	6.5	Bq/Kg raw	± 4.1	Bq/Kg raw	10.1	Cs137 7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Kogomi	Sekimoto, Kitaiabaraki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken	Ogawa, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.4	Cs137 8.0 Bq/Kg raw
			Cs134	0.4	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Bracken	Yotsukura, Iwaki	Apr-12	Cs137	4.3	Bq/Kg raw	± 8.1	Bq/Kg raw	7.8	Cs137 8.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 4.8	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Bracken	Fujiwara, Joban, Iwaki	Apr-12	Cs137	8.7	Bq/Kg raw	± 4.5	Bq/Kg raw	23.0	Cs137 7.0 Bq/Kg raw
			Cs134	14.3	Bq/Kg raw	± 4.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken	Miyamaguchi, Tono, Iwaki	Apr-12	Cs137	5.0	Bq/Kg raw	± 4.4	Bq/Kg raw	17.4	Cs137 7.0 Bq/Kg raw
			Cs134	12.4	Bq/Kg raw	± 4.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken	Kaneyama, Iwaki	Apr-12	Cs137	6.1	Bq/Kg raw	± 3.2	Bq/Kg raw	12.3	Cs137 7.0 Bq/Kg raw
			Cs134	6.2	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken	Soeno, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken	Minamiodaira, Tabito, Iwaki	Apr-12	Cs137	54.9	Bq/Kg raw	± 12.7	Bq/Kg raw	92.4	Cs137 7.0 Bq/Kg raw
			Cs134	37.5	Bq/Kg raw	± 8.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Pickled Bracken	Furudono, Ishikawa	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137 7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Takahagi, Ibaraki	Apr-12	Cs137	151	Bq/Kg raw	± 31.0	Bq/Kg raw	272	Cs137 7.0 Bq/Kg raw
			Cs134	121	Bq/Kg raw	± 24.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Nakashio, Taira, Iwaki	Apr-12	Cs137	14.8	Bq/Kg raw	± 4.2	Bq/Kg raw	26.1	Cs137 7.0 Bq/Kg raw
			Cs134	11.3	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	39.1	Bq/Kg raw	± 9.5	Bq/Kg raw	61.9	Cs137 7.0 Bq/Kg raw
			Cs134	22.8	Bq/Kg raw	± 5.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Akai, Taira, Iwaki	Apr-12	Cs137	45.0	Bq/Kg raw	± 10.7	Bq/Kg raw	84.3	Cs137 7.0 Bq/Kg raw
			Cs134	39.3	Bq/Kg raw	± 8.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Komoda, Kashima, Iwaki	Apr-12	Cs137	54.9	Bq/Kg raw	± 12.8	Bq/Kg raw	109	Cs137 7.0 Bq/Kg raw
			Cs134	53.6	Bq/Kg raw	± 11.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Hashirikuma, Kashima, Iwaki	Apr-12	Cs137	41.9	Bq/Kg raw	± 10.0	Bq/Kg raw	77.9	Cs137 7.0 Bq/Kg raw
			Cs134	36.0	Bq/Kg raw	± 8.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	42.8	Bq/Kg raw	± 9.9	Bq/Kg raw	71.2	Cs137 7.0 Bq/Kg raw
			Cs134	28.4	Bq/Kg raw	± 6.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimoyunagaya, Joban, Iwaki	Apr-12	Cs137	13.2	Bq/Kg raw	± 4.8	Bq/Kg raw	23.8	Cs137 7.0 Bq/Kg raw
			Cs134	10.6	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimoyunagaya, Joban, Iwaki	Apr-12	Cs137	15.9	Bq/Kg raw	± 5.2	Bq/Kg raw	35.2	Cs137 7.0 Bq/Kg raw
			Cs134	19.3	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimoyunagaya, Joban, Iwaki	Apr-12	Cs137	39.1	Bq/Kg raw	± 9.1	Bq/Kg raw	62.8	Cs137 7.0 Bq/Kg raw
			Cs134	23.7	Bq/Kg raw	± 5.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Fujiwara, Joban, Iwaki	Apr-12	Cs137	18.1	Bq/Kg raw	± 6.0	Bq/Kg raw	34.8	Cs137 7.0 Bq/Kg raw
			Cs134	16.7	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Mizunoya, Joban, Iwaki	Apr-12	Cs137	50.7	Bq/Kg raw	± 12.1	Bq/Kg raw	89.8	Cs137 7.0 Bq/Kg raw
			Cs134	39.1	Bq/Kg raw	± 9.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Bamboo shoot	Shimoyunagaya, Joban, Iwaki	Apr-12	Cs137	32.7	Bq/Kg raw	± 8.3	Bq/Kg raw	58.8	Cs137	7.0 Bq/Kg raw
			Cs134	26.1	Bq/Kg raw	± 6.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Izumi, Iwaki	Apr-12	Cs137	68.7	Bq/Kg raw	± 15.9	Bq/Kg raw	118	Cs137	7.0 Bq/Kg raw
			Cs134	49.2	Bq/Kg raw	± 11.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Fujima, Taira, Iwaki	Apr-12	Cs137	340	Bq/Kg raw	± 68	Bq/Kg raw	607	Cs137	7.0 Bq/Kg raw
			Cs134	267	Bq/Kg raw	± 53	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Idekura, Nakoso, Iwaki	Apr-12	Cs137	211	Bq/Kg raw	± 42.0	Bq/Kg raw	373	Cs137	7.0 Bq/Kg raw
			Cs134	162	Bq/Kg raw	± 32.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Boiled bamboo shoot	Kamiyunagaya, Joban, Iwaki	May-11	Cs137	4.5	Bq/Kg raw	± 2.3	Bq/Kg raw	9.0	Cs137	7.0 Bq/Kg raw
			Cs134	4.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Nishiki, Iwaki	Apr-12	Cs137	36.1	Bq/Kg raw	± 17.0	Bq/Kg raw	58.4	Cs137	8.0 Bq/Kg raw
			Cs134	22.3	Bq/Kg raw	± 8.2	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Bamboo shoot	Ebata, Iwaki	Apr-12	Cs137	16.1	Bq/Kg raw	± 4.6	Bq/Kg raw	30.3	Cs137	7.0 Bq/Kg raw
			Cs134	14.2	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Soeno, Iwaki	Apr-12	Cs137	16.0	Bq/Kg raw	± 4.7	Bq/Kg raw	33.0	Cs137	7.0 Bq/Kg raw
			Cs134	17.0	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Iwaki	Apr-12	Cs137	78.4	Bq/Kg raw	± 17.2	Bq/Kg raw	139	Cs137	7.0 Bq/Kg raw
			Cs134	60.1	Bq/Kg raw	± 12.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Sekimoto, Kitaibaraki	Apr-12	Cs137	58.0	Bq/Kg raw	± 12.6	Bq/Kg raw	100	Cs137	7.0 Bq/Kg raw
			Cs134	41.8	Bq/Kg raw	± 8.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Takahagi, Kitaibaraki	Apr-12	Cs137	554	Bq/Kg raw	± 111	Bq/Kg raw	993	Cs137	7.0 Bq/Kg raw
			Cs134	439	Bq/Kg raw	± 88	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Ohisa, Iwaki	Apr-12	Cs137	1900	Bq/Kg raw	± 380	Bq/Kg raw	3,350	Cs137	7.0 Bq/Kg raw
			Cs134	1450	Bq/Kg raw	± 290	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Komagome, Yotsukura, Iwaki	Unknown	Cs137	287	Bq/Kg raw	± 57.0	Bq/Kg raw	509	Cs137	7.0 Bq/Kg raw
			Cs134	222	Bq/Kg raw	± 44.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Yotsukura, Iwaki	Apr-12	Cs137	646	Bq/Kg raw	± 129	Bq/Kg raw	1,158	Cs137	7.0 Bq/Kg raw
			Cs134	512	Bq/Kg raw	± 102	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Shibahara, Ogawa, Iwaki	Mar-12	Cs137	94.8	Bq/Kg raw	± 20.3	Bq/Kg raw	176	Cs137	7.0 Bq/Kg raw
			Cs134	81.6	Bq/Kg raw	± 16.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Nakakabeya, Taira, Iwaki	Apr-12	Cs137	136	Bq/Kg raw	± 28.0	Bq/Kg raw	249	Cs137	7.0 Bq/Kg raw
			Cs134	113	Bq/Kg raw	± 23.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Taira, Iwaki	Apr-12	Cs137	286	Bq/Kg raw	± 36.2	Bq/Kg raw	466	Cs137	8.0 Bq/Kg raw
			Cs134	180	Bq/Kg raw	± 21.7	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Shiitake mushrooms	Akai, Taira, Iwaki	Apr-12	Cs137	266	Bq/Kg raw	± 55.0	Bq/Kg raw	482	Cs137	7.0 Bq/Kg raw
			Cs134	216	Bq/Kg raw	± 43.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Akai, Taira, Iwaki	Apr-12	Cs137	623	Bq/Kg raw	± 125	Bq/Kg raw	1,100	Cs137	7.0 Bq/Kg raw
			Cs134	477	Bq/Kg raw	± 95.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Kamiogoe, Taira, Iwaki	Apr-12	Cs137	614	Bq/Kg raw	± 123	Bq/Kg raw	1,100	Cs137	7.0 Bq/Kg raw
			Cs134	486	Bq/Kg raw	± 97.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Manome, Taira, Iwaki	Unknown	Cs137	1820	Bq/Kg raw	± 360	Bq/Kg raw	3,240	Cs137	7.0 Bq/Kg raw
			Cs134	1420	Bq/Kg raw	± 280	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Kamikatayose, Taira, Iwaki	Apr-12	Cs137	129	Bq/Kg raw	± 26.0	Bq/Kg raw	234	Cs137	7.0 Bq/Kg raw
			Cs134	105	Bq/Kg raw	± 21.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Koizumi, Taira, Iwaki	Apr-12	Cs137	403	Bq/Kg raw	± 81.0	Bq/Kg raw	716	Cs137	7.0 Bq/Kg raw
			Cs134	313	Bq/Kg raw	± 63.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Shiitake mushrooms	Natsui, Taira, Iwaki	Apr-12	Cs137	1130	Bq/Kg raw	± 230	Bq/Kg raw	1,972	Cs137	7.0 Bq/Kg raw
			Cs134	842	Bq/Kg raw	± 168	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Numanouchi, Iwaki	Unknown	Cs137	547	Bq/Kg raw	± 109	Bq/Kg raw	966	Cs137	7.0 Bq/Kg raw
			Cs134	419	Bq/Kg raw	± 84.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Miwa, Iwaki	Unknown	Cs137	186	Bq/Kg raw	± 37.0	Bq/Kg raw	326	Cs137	7.0 Bq/Kg raw
			Cs134	140	Bq/Kg raw	± 28.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Kitayoshima, Yoshima, Iwaki	Apr-12	Cs137	750	Bq/Kg raw	± 83.8	Bq/Kg raw	1,264	Cs137	8.0 Bq/Kg raw
			Cs134	514	Bq/Kg raw	± 52.8	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Shiitake mushrooms	Miya, Uchigo, Iwaki	Apr-12	Cs137	263	Bq/Kg raw	± 53.0	Bq/Kg raw	467	Cs137	7.0 Bq/Kg raw
			Cs134	204	Bq/Kg raw	± 41.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Mimaya, Uchigo, Iwaki	Apr-12	Cs137	823	Bq/Kg raw	± 165	Bq/Kg raw	1,479	Cs137	7.0 Bq/Kg raw
			Cs134	656	Bq/Kg raw	± 131	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Toyoma, Taira, Iwaki	Apr-12	Cs137	1080	Bq/Kg raw	± 222	Bq/Kg raw	1,908	Cs137	7.0 Bq/Kg raw
			Cs134	828	Bq/Kg raw	± 166	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Shimoyunagaya, Joban, Iwaki	Apr-12	Cs137	107	Bq/Kg raw	± 20.7	Bq/Kg raw	177	Cs137	8.0 Bq/Kg raw
			Cs134	69.6	Bq/Kg raw	± 12.4	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Shiitake mushrooms	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	367	Bq/Kg raw	± 73.0	Bq/Kg raw	646	Cs137	7.0 Bq/Kg raw
			Cs134	279	Bq/Kg raw	± 56.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Shimokajiro, Onahama, Iwaki	Mar-12	Cs137	258	Bq/Kg raw	± 55.0	Bq/Kg raw	490	Cs137	7.0 Bq/Kg raw
			Cs134	232	Bq/Kg raw	± 46.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Okaona, Onahama, Iwaki	Apr-12	Cs137	178	Bq/Kg raw	± 36.0	Bq/Kg raw	323	Cs137	7.0 Bq/Kg raw
			Cs134	145	Bq/Kg raw	± 29.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Ohara, Onahama, Iwaki	Apr-12	Cs137	354	Bq/Kg raw	± 71.0	Bq/Kg raw	629	Cs137	7.0 Bq/Kg raw
			Cs134	275	Bq/Kg raw	± 55.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Tamagawa, Onahama, Iwaki	Apr-12	Cs137	190	Bq/Kg raw	± 40.0	Bq/Kg raw	324	Cs137	7.0 Bq/Kg raw
			Cs134	134	Bq/Kg raw	± 27.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	492	Bq/Kg raw	± 98.0	Bq/Kg raw	891	Cs137	7.0 Bq/Kg raw
			Cs134	399	Bq/Kg raw	± 80.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	918	Bq/Kg raw	± 184	Bq/Kg raw	1,621	Cs137	7.0 Bq/Kg raw
			Cs134	703	Bq/Kg raw	± 141	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Shimokawa, Izumi, Iwaki	Apr-12	Cs137	83.3	Bq/Kg raw	± 17.3	Bq/Kg raw	154	Cs137	7.0 Bq/Kg raw
			Cs134	70.8	Bq/Kg raw	± 14.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Izumigaoka, Iwaki	Apr-12	Cs137	788	Bq/Kg raw	± 158	Bq/Kg raw	1,382	Cs137	7.0 Bq/Kg raw
			Cs134	594	Bq/Kg raw	± 119	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Ebata, Iwaki	Apr-12	Cs137	222	Bq/Kg raw	± 44.0	Bq/Kg raw	397	Cs137	7.0 Bq/Kg raw
			Cs134	175	Bq/Kg raw	± 35.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Idekura, Nakoso, Iwaki	Apr-12	Cs137	492	Bq/Kg raw	± 98.0	Bq/Kg raw	882	Cs137	7.0 Bq/Kg raw
			Cs134	390	Bq/Kg raw	± 78.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Izumi, Iwaki	Apr-12	Cs137	408	Bq/Kg raw	± 82.0	Bq/Kg raw	718	Cs137	7.0 Bq/Kg raw
			Cs134	310	Bq/Kg raw	± 62.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Tanabe, Watanabe, Iwaki	Apr-12	Cs137	57.4	Bq/Kg raw	± 12.9	Bq/Kg raw	98.1	Cs137	7.0 Bq/Kg raw
			Cs134	40.7	Bq/Kg raw	± 9.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Fudaba, Takakura, Iwaki	Apr-12	Cs137	393	Bq/Kg raw	± 48.6	Bq/Kg raw	672	Cs137	8.0 Bq/Kg raw
			Cs134	279	Bq/Kg raw	± 31.3	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Shiitake mushrooms	Kuroda, Tabito, Iwaki	Apr-12	Cs137	171	Bq/Kg raw	± 35.0	Bq/Kg raw	173	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 25.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Shiitake mushrooms	Tabito, Iwaki	Apr-12	Cs137	130	Bq/Kg raw	± 28.0	Bq/Kg raw	243	Cs137	7.0 Bq/Kg raw
			Cs134	113	Bq/Kg raw	± 23.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Kaneyama, Iwaki	Apr-12	Cs137	616	Bq/Kg raw	± 123	Bq/Kg raw	1,103	Cs137	7.0 Bq/Kg raw
			Cs134	487	Bq/Kg raw	± 97.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Nishiki, Iwaki	Apr-12	Cs137	121	Bq/Kg raw	± 25.0	Bq/Kg raw	216	Cs137	7.0 Bq/Kg raw
			Cs134	95.4	Bq/Kg raw	± 19.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Yamada, Iwaki	Apr-12	Cs137	245	Bq/Kg raw	± 50.0	Bq/Kg raw	435	Cs137	7.0 Bq/Kg raw
			Cs134	190	Bq/Kg raw	± 38.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushrooms	Yamada, Iwaki	Apr-12	Cs137	198	Bq/Kg raw	± 30.7	Bq/Kg raw	336	Cs137	8.0 Bq/Kg raw
			Cs134	138	Bq/Kg raw	± 19.4	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Enoki mushrooms	Esashi, Iwate	Unknown	Cs137	4.7	Bq/Kg raw	± 3.0	Bq/Kg raw	4.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried shiitake mushrooms	Tabito, Iwaki	Apr-12	Cs137	875	Bq/Kg raw	± 177	Bq/Kg raw	1,614	Cs137	7.0 Bq/Kg raw
			Cs134	739	Bq/Kg raw	± 148	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried shiitake mushrooms (reconstituted)	Tabito, Iwaki	Apr-12	Cs137	19.1	Bq/Kg raw	± 5.1	Bq/Kg raw	40.7	Cs137	7.0 Bq/Kg raw
			Cs134	21.6	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried shiitake mushrooms	Minamiodaira, Tabito, Iwaki	Apr-12	Cs137	1680	Bq/Kg raw	± 340	Bq/Kg raw	3,010	Cs137	7.0 Bq/Kg raw
			Cs134	1330	Bq/Kg raw	± 270	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Blueberries	Wakabadai, Iwaki	Aug-11	Cs137	55.2	Bq/Kg raw	± 12.0	Bq/Kg raw	97.7	Cs137	7.0 Bq/Kg raw
			Cs134	42.5	Bq/Kg raw	± 8.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Blueberries	Komoda, Kashima, Iwaki	Aug-11	Cs137	66.3	Bq/Kg raw	± 14.3	Bq/Kg raw	113	Cs137	7.0 Bq/Kg raw
			Cs134	46.7	Bq/Kg raw	± 9.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Blueberries	Takaku, Chuodai, Iwaki	Aug-11	Cs137	137	Bq/Kg raw	± 28.0	Bq/Kg raw	252	Cs137	7.0 Bq/Kg raw
			Cs134	115	Bq/Kg raw	± 23.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Apples	Hobara, Date	Oct-11	Cs137	30.8	Bq/Kg raw	± 7.2	Bq/Kg raw	52.4	Cs137	7.0 Bq/Kg raw
			Cs134	21.6	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Oranges	Kamiyunagaya, Joban, Iwaki	Mar-12	Cs137	83.2	Bq/Kg raw	± 18.3	Bq/Kg raw	152	Cs137	7.0 Bq/Kg raw
			Cs134	69.2	Bq/Kg raw	± 14.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Mandarin oranges	Sakuragaoka, Iwaki	Feb-12	Cs137	11.9	Bq/Kg raw	± 3.5	Bq/Kg raw	19.5	Cs137	7.0 Bq/Kg raw
			Cs134	7.6	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Mandarin oranges	Kamiyada, Joban, Iwaki	Feb-12	Cs137	179	Bq/Kg raw	± 28.3	Bq/Kg raw	282	Cs137	8.0 Bq/Kg raw
			Cs134	103	Bq/Kg raw	± 16.3	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Natsumikan	Nakashio, Taira, Iwaki	Apr-12	Cs137	95.3	Bq/Kg raw	± 19.6	Bq/Kg raw	173	Cs137	7.0 Bq/Kg raw
			Cs134	77.9	Bq/Kg raw	± 15.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Natsumikan	Sakuragaoka, Iwaki	Feb-12	Cs137	12.5	Bq/Kg raw	± 3.7	Bq/Kg raw	23.2	Cs137	7.0 Bq/Kg raw
			Cs134	10.7	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Natsumikan	Shonandai, Iwaki	Apr-12	Cs137	28.9	Bq/Kg raw	± 9.3	Bq/Kg raw	51.8	Cs137	8.0 Bq/Kg raw
			Cs134	22.9	Bq/Kg raw	± 6.8	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Kiwi fruit	Ori, Yoshima, Iwaki	Dec-11	Cs137	32.5	Bq/Kg raw	± 7.7	Bq/Kg raw	54.9	Cs137	7.0 Bq/Kg raw
			Cs134	22.4	Bq/Kg raw	± 5.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Kumquat	Kubo, Taira, Iwaki	Jan-12	Cs137	7.7	Bq/Kg raw	± 4.5	Bq/Kg raw	12.3	Cs137	8.0 Bq/Kg raw
			Cs134	4.6	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Yuzu	Moegidai, Izumi, Iwaki	Dec-11	Cs137	70.0	Bq/Kg raw	± 15.4	Bq/Kg raw	128	Cs137	7.0 Bq/Kg raw
			Cs134	57.9	Bq/Kg raw	± 12.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Yuzu	Ohara, Onahama, Iwaki	Apr-12	Cs137	64.2	Bq/Kg raw	± 14.4	Bq/Kg raw	119	Cs137	7.0 Bq/Kg raw
			Cs134	54.4	Bq/Kg raw	± 11.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Persimmons	Kaminodera, Fukushima	Nov-11	Cs137	14.8	Bq/Kg raw	± 4.4	Bq/Kg raw	26.7	Cs137	7.0 Bq/Kg raw
			Cs134	11.9	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Persimmons	Shimoogawa, Ogawa, Iwaki	Nov-11	Cs137	4.8	Bq/Kg raw	± 2.6	Bq/Kg raw	9.2	Cs137	7.0 Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried persimmons	Moegidai, Izumi, Iwaki	Nov-11	Cs137	15.7	Bq/Kg raw	± 4.6	Bq/Kg raw	27.1	Cs137	7.0 Bq/Kg raw
			Cs134	11.4	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried persimmons	Shimoyoshima, Yoshima, Iwaki	Nov-11	Cs137	14.4	Bq/Kg raw	± 7.0	Bq/Kg raw	24.5	Cs137	8.0 Bq/Kg raw
			Cs134	10.1	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Dried persimmons	Akai, Taira, Iwaki	Dec-11	Cs137	42.5	Bq/Kg raw	± 9.6	Bq/Kg raw	77.8	Cs137	7.0 Bq/Kg raw
			Cs134	35.3	Bq/Kg raw	± 7.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Plums	Kubo, Taira, Iwaki	Jun-11	Cs137	5.9	Bq/Kg raw	± 2.7	Bq/Kg raw	8.4	Cs137	7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried plums	Wakayama	Unknown	Cs137	2.0	Bq/Kg raw	± 1.8	Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried plums	Okaona, Onahama, Iwaki	Jun-11	Cs137	64.2	Bq/Kg raw	± 13.9	Bq/Kg raw	117	Cs137	7.0 Bq/Kg raw
			Cs134	52.4	Bq/Kg raw	± 10.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Ibaraki	Unknown	Cs137	3.1	Bq/Kg raw	± 2.4	Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Naraha, Futaba	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Miwa, Iwaki	Apr-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Tairanakayama, Iwaki	Nov-11	Cs137	1.0	Bq/Kg raw	± 3.4	Bq/Kg raw	2.0	Cs137	8.0 Bq/Kg raw
			Cs134	1.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Soybean paste	Tabito, Iwaki	Sep-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Kaneyama, Numabe, Iwaki	May-10	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Unknown	Unknown	Cs137	3.0	Bq/Kg raw	± 2.0	Bq/Kg raw	5.3	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Nakakabeya, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Soybean paste	Kawabe, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Eggs	Kita, Nigata	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Eggs	Kamiogawa, Ogawa, Iwaki	Mar-12	Cs137	1.9	Bq/Kg raw	± 1.0	Bq/Kg raw	3.6	Cs137	5.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 0.9	Bq/Kg raw		Cs134	3.0 Bq/Kg raw
Eggs	Kamiogawa, Ogawa, Iwaki	Mar-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Eggs	Kaneyama, Numabe, Iwaki	Apr-12	Cs137	2.3	Bq/Kg raw	± 2.3	Bq/Kg raw	5.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Wild boa meat	Ogawa, Iwaki	Unknown	Cs137	78.3	Bq/Kg raw	± 16.5	Bq/Kg raw	137	Cs137	7.0 Bq/Kg raw
			Cs134	58.4	Bq/Kg raw	± 12.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Wild boa meat	Shimoichigaya, Miwa, Iwaki	Apr-12	Cs137	31.7	Bq/Kg raw	± 7.4	Bq/Kg raw	59.8	Cs137	7.0 Bq/Kg raw
			Cs134	28.1	Bq/Kg raw	± 6.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Wild boa meat	Nakadera, Miwa, Iwaki	Apr-12	Cs137	41.6	Bq/Kg raw	± 9.3	Bq/Kg raw	72.5	Cs137	7.0 Bq/Kg raw
			Cs134	30.9	Bq/Kg raw	± 6.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Wild boa meat	Nakadera, Miwa, Iwaki	Apr-12	Cs137	68.9	Bq/Kg raw	± 14.6	Bq/Kg raw	120	Cs137	7.0 Bq/Kg raw
			Cs134	51.5	Bq/Kg raw	± 10.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Minced pork	Samegawa, Higashishirakawa	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137	8.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 4.4	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Egoma minced meat cutlet	Fukushima	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Minced Japanese beef cutlet	Fukushima	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Fried chickens	Fukushima	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Curry (prepared)	Unknown	Apr-12	Cs137	2.2	Bq/Kg raw	± 1.8	Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Blueberry jam	Komoda, Kashima, Iwaki	Aug-11	Cs137	57.6	Bq/Kg raw	± 12.6	Bq/Kg raw	108	Cs137	7.0 Bq/Kg raw
			Cs134	50.0	Bq/Kg raw	± 10.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
School lunch	Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
School lunch	Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
School lunch	Joban, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Pickled Japanese radish	Kanari, Onahama, Iwaki	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Plum syrup	Shimokawa, Izumi, Iwaki	Jun-11	Cs137	12.2	Bq/Kg raw	± 3.6	Bq/Kg raw	26.6	Cs137	7.0 Bq/Kg raw
			Cs134	14.4	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Crab stick flakes	Unknown	Apr-12	Cs137	3.4	Bq/Kg raw	± 2.0	Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Crab stick	Unknown	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Green tea drink	Unknown	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.1	Cs137	7.0 Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Milk	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	3.0 Bq/Kg raw
Milk	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137	5.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134	3.0 Bq/Kg raw
Yogurt	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Cod and sole (mixed)	Iwaki	Apr-12	Cs137	4.1	Bq/Kg raw	± 2.8	Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Eel	Shimoogoe, Taira, Iwaki	Apr-12	Cs137	287	Bq/Kg raw	± 36.4	Bq/Kg raw	464	Cs137	7.0 Bq/Kg raw
			Cs134	177	Bq/Kg raw	± 22.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese mitten crab	Nasuigawa, Iwaki	Apr-12	Cs137	34.0	Bq/Kg raw	± 8.0	Bq/Kg raw	69.7	Cs137	7.0 Bq/Kg raw
			Cs134	35.7	Bq/Kg raw	± 7.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Seaweed	Onahama, Iwaki	Mar-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Seaweed	Nagasaki, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Seaweed	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	11.1	Bq/Kg raw	± 3.7	Bq/Kg raw	18.2	Cs137 7.0 Bq/Kg raw
			Cs134	7.1	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Seaweed	Onahama, Iwaki	Apr-12	Cs137	2.7	Bq/Kg raw	± 0.2	Bq/Kg raw	6.0	Cs137 8.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 4.9	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Seaweed	Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Seaweed	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137 7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sea Kelp	Korea	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.3	Cs137 7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shellfish (Mussels/Shuri mussels)	Shimokajiro, Onahama, Iwaki	Apr-12	Cs137	5.4	Bq/Kg raw	± 2.8	Bq/Kg raw	11.8	Cs137 7.0 Bq/Kg raw
			Cs134	6.4	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Hijiki seaweed	Toyoma, Iwaki	Apr-12	Cs137	15.0	Bq/Kg raw	± 4.9	Bq/Kg raw	20.2	Cs137 7.0 Bq/Kg raw
			Cs134	5.2	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Alkaline ionized water	Unknown	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.1	Cs137 5.0 Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Mineral water	Tonami, Toyama	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Mineral water	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137 5.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Pure water	Unknown	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 5.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Water	Unknown	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137 5.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Water	Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Water	Taki, Tono, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137 5.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap Water	Hirakata, Kitaibaraki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 5.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap Water	Kamiarakawa, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.2	Cs137 7.0 Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tap Water	Takijiri, Izumi, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap Water	Kaminiida, Yotsukura, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tap Water	Koya, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap Water	Takasaka, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137 5.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap Water	Kojima, Uchigo, Iwaki	Apr-12	Cs137	1.1	Bq/Kg raw	± 1.1	Bq/Kg raw	1.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tap Water	Miya, Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.2	Cs137 5.0 Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Tap Water	Uchigo, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 5.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap Water	Nishiki, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137 5.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Naraha, Futaba	Apr-12	Cs137	1.4	Bq/Kg raw	± 1.3	Bq/Kg raw	2.8	Cs137 5.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Kitsunezuka, Yotsukura, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 5.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Kamihirakubo, Taira, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137 5.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Iritono, Tono, Iwaki	Apr-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Samegawa, Higashishirakawa	Feb-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.1	Cs137 5.0 Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Godo, Miwa, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137 5.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.3	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Kawabe, Iwaki	Apr-12	Cs137	1.2	Bq/Kg raw	± 1.1	Bq/Kg raw	1.2	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Rain water (storaged)	Shonandai, Iwaki	Apr-12	Cs137	1.6	Bq/Kg raw	± 1.5	Bq/Kg raw	3.0	Cs137 6.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.4	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Groundwater	Saiso, Miwa, Iwaki	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Nikko maple	Onahama, Iwaki	Oct-11	Cs137	481	Bq/Kg raw	± 107	Bq/Kg raw	898	Cs137 7.0 Bq/Kg raw
			Cs134	417	Bq/Kg raw	± 87.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese torreya	Moegidai, Izumi, Iwaki	Mar-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	15.5	Cs137 7.0 Bq/Kg raw
			Cs134	15.5	Bq/Kg raw	± 11.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sawdust	Unknown	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	13.5	Cs137 7.0 Bq/Kg raw
			Cs134	13.5	Bq/Kg raw	± 11.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Compost (cow manure)	Sukagawa	Apr-12	Cs137	97.0	Bq/Kg raw	± 23.0	Bq/Kg raw	170	Cs137 8.0 Bq/Kg raw
			Cs134	73.0	Bq/Kg raw	± 14.0	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Compost	Moegidai, Izumi, Iwaki	Mar-12	Cs137	501	Bq/Kg raw	± 60.0	Bq/Kg raw	842	Cs137 8.0 Bq/Kg raw
			Cs134	341	Bq/Kg raw	± 38.0	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Ashes	Saizuchikoji, Taira, Iwaki	Apr-12	Cs137	3659.5	Bq/Kg raw	± 333.0	Bq/Kg raw	9076	Cs137 8.0 Bq/Kg raw
			Cs134	5416.4	Bq/Kg raw	± 508.1	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Soil	Maehara, Ogawa, Iwaki	Apr-12	Cs137	240	Bq/Kg dry	± 33.0	Bq/Kg dry	417	Cs137 8.0 Bq/Kg dry
			Cs134	177	Bq/Kg dry	± 23.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Maebara, Shimoogawa, Iwaki	Apr-12	Cs137	371	Bq/Kg dry	± 46.0	Bq/Kg dry	583	Cs137 8.0 Bq/Kg dry
			Cs134	212	Bq/Kg dry	± 27.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shimokabeya, Taira, Iwaki	Apr-12	Cs137	408	Bq/Kg dry	± 48.0	Bq/Kg dry	692	Cs137 8.0 Bq/Kg dry
			Cs134	284	Bq/Kg dry	± 31.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Jiyugaoka, Taira, Iwaki	Apr-12	Cs137	111	Bq/Kg dry	± 18.0	Bq/Kg dry	191	Cs137 8.0 Bq/Kg dry
			Cs134	80.0	Bq/Kg dry	± 13.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Yagawase, Taira, Iwaki	Apr-12	Cs137	161	Bq/Kg dry	± 27.0	Bq/Kg dry	270	Cs137 8.0 Bq/Kg dry
			Cs134	109	Bq/Kg dry	± 17.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shimohirakubo, Taira, Iwaki	Apr-12	Cs137	60.0	Bq/Kg dry	± 14.0	Bq/Kg dry	105	Cs137 8.0 Bq/Kg dry
			Cs134	45.0	Bq/Kg dry	± 9.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Soil	Shimoarakawa, Taira, Iwaki	Apr-12	Cs137	379	Bq/Kg dry	± 47.0	644	Cs137	8.0	Bq/Kg dry
			Cs134	265	Bq/Kg dry	± 30.0		Cs134	8.0	Bq/Kg dry
Soil	Kashima, Chuodai, Iwaki	Apr-12	Cs137	180	Bq/Kg dry	± 26.0	304	Cs137	8.0	Bq/Kg dry
			Cs134	124	Bq/Kg dry	± 17.0		Cs134	8.0	Bq/Kg dry
Soil	Takaku, Chuodai, Iwaki	Apr-12	Cs137	808	Bq/Kg dry	± 87.0	1,355	Cs137	8.0	Bq/Kg dry
			Cs134	547	Bq/Kg dry	± 56.0		Cs134	8.0	Bq/Kg dry
Soil	Takasaka, Uchigo, Iwaki	Apr-12	Cs137	101	Bq/Kg dry	± 18.0	163	Cs137	8.0	Bq/Kg dry
			Cs134	62.0	Bq/Kg dry	± 11.0		Cs134	8.0	Bq/Kg dry
Soil	Tsuzura, Uchigo, Iwaki	Apr-12	Cs137	367	Bq/Kg dry	± 45.0	597	Cs137	8.0	Bq/Kg dry
			Cs134	230	Bq/Kg dry	± 27.0		Cs134	8.0	Bq/Kg dry
Soil	Hashirikuma, Kashima, Iwaki	Apr-12	Cs137	108	Bq/Kg dry	± 19.0	190	Cs137	8.0	Bq/Kg dry
			Cs134	82.0	Bq/Kg dry	± 13.0		Cs134	8.0	Bq/Kg dry
Soil	Kamiyada, Joban, Iwaki	Apr-12	Cs137	217	Bq/Kg dry	± 30.0	336	Cs137	8.0	Bq/Kg dry
			Cs134	119	Bq/Kg dry	± 17.0		Cs134	8.0	Bq/Kg dry
Soil	Sekifune, Joban, Iwaki	Apr-12	Cs137	216	Bq/Kg dry	± 30.0	370	Cs137	8.0	Bq/Kg dry
			Cs134	154	Bq/Kg dry	± 20.0		Cs134	8.0	Bq/Kg dry
Soil	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	157	Bq/Kg dry	± 27.0	272	Cs137	8.0	Bq/Kg dry
			Cs134	115	Bq/Kg dry	± 17.0		Cs134	8.0	Bq/Kg dry
Soil	Moegidai, Izumi, Iwaki	Mar-12	Cs137	224	Bq/Kg dry	± 29.0	387	Cs137	8.0	Bq/Kg dry
			Cs134	163	Bq/Kg dry	± 19.0		Cs134	8.0	Bq/Kg dry
Soil	Miyamada, Tono, Iwaki	Apr-12	Cs137	52.0	Bq/Kg dry	± 14.0	95.0	Cs137	8.0	Bq/Kg dry
			Cs134	43.0	Bq/Kg dry	± 9.0		Cs134	8.0	Bq/Kg dry
Soil	Kubota, Nakoso, Iwaki	Apr-12	Cs137	316	Bq/Kg dry	± 40.0	525	Cs137	8.0	Bq/Kg dry
			Cs134	209	Bq/Kg dry	± 26.0		Cs134	8.0	Bq/Kg dry
Field soil	Nakadera, Miwa, Iwaki	Apr-12	Cs137	148	Bq/Kg dry	± 23.0	232	Cs137	8.0	Bq/Kg dry
			Cs134	84.0	Bq/Kg dry	± 14.0		Cs134	8.0	Bq/Kg dry
Field soil	Kitayoshima, Yoshima, Iwaki	Apr-12	Cs137	220	Bq/Kg dry	± 31.0	353	Cs137	8.0	Bq/Kg dry
			Cs134	133	Bq/Kg dry	± 19.0		Cs134	8.0	Bq/Kg dry
Field soil	Akai, Taira, Iwaki	Apr-12	Cs137	252	Bq/Kg dry	± 35.0	421	Cs137	8.0	Bq/Kg dry
			Cs134	169	Bq/Kg dry	± 22.0		Cs134	8.0	Bq/Kg dry
Field soil	Akai, Taira, Iwaki	Mar-12	Cs137	176	Bq/Kg dry	± 24.0	300	Cs137	8.0	Bq/Kg dry
			Cs134	124	Bq/Kg dry	± 16.0		Cs134	8.0	Bq/Kg dry
Field soil	Akai, Taira, Iwaki	Mar-12	Cs137	48.0	Bq/Kg dry	± 12.0	84.0	Cs137	8.0	Bq/Kg dry
			Cs134	36.0	Bq/Kg dry	± 8.0		Cs134	8.0	Bq/Kg dry
Field soil	Shimohirakubo, Taira, Iwaki	Apr-12	Cs137	668	Bq/Kg dry	± 72.0	1,133	Cs137	8.0	Bq/Kg dry
			Cs134	465	Bq/Kg dry	± 48.0		Cs134	8.0	Bq/Kg dry
Field soil	Shimohirakubo, Taira, Iwaki	Apr-12	Cs137	63.0	Bq/Kg dry	± 14.0	109	Cs137	8.0	Bq/Kg dry
			Cs134	46.0	Bq/Kg dry	± 10.0		Cs134	8.0	Bq/Kg dry
Field soil	Shimotakaku, Taira, Iwaki	Apr-12	Cs137	516	Bq/Kg dry	± 60.0	904	Cs137	8.0	Bq/Kg dry
			Cs134	388	Bq/Kg dry	± 40.0		Cs134	8.0	Bq/Kg dry
Field soil	Shimotakaku, Taira, Iwaki	Apr-12	Cs137	265	Bq/Kg dry	± 35.0	438	Cs137	8.0	Bq/Kg dry
			Cs134	173	Bq/Kg dry	± 21.0		Cs134	8.0	Bq/Kg dry
Field soil	Shimotakaku, Taira, Iwaki	Apr-12	Cs137	179	Bq/Kg dry	± 25.0	285	Cs137	8.0	Bq/Kg dry
			Cs134	106	Bq/Kg dry	± 15.0		Cs134	8.0	Bq/Kg dry
Field soil	Shimotakaku, Taira, Iwaki	Apr-12	Cs137	212	Bq/Kg dry	± 31.0	382	Cs137	8.0	Bq/Kg dry
			Cs134	170	Bq/Kg dry	± 21.0		Cs134	8.0	Bq/Kg dry

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Field soil	Kitashirado, Taira, Iwaki	Apr-12	Cs137	207	Bq/Kg dry	± 29.0	Bq/Kg dry	360	Cs137	8.0 Bq/Kg dry
			Cs134	153	Bq/Kg dry	± 20.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Kitashirado, Taira, Iwaki	Apr-12	Cs137	218	Bq/Kg dry	± 30.0	Bq/Kg dry	380	Cs137	8.0 Bq/Kg dry
			Cs134	162	Bq/Kg dry	± 21.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Fujima, Taira, Iwaki	Apr-12	Cs137	114	Bq/Kg dry	± 19.0	Bq/Kg dry	181	Cs137	8.0 Bq/Kg dry
			Cs134	67.0	Bq/Kg dry	± 11.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Numanouchi, Taira, Iwaki	Apr-12	Cs137	245	Bq/Kg dry	± 32.0	Bq/Kg dry	386	Cs137	8.0 Bq/Kg dry
			Cs134	141	Bq/Kg dry	± 19.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Takaku, Chuodai, Iwaki	Apr-12	Cs137	40.0	Bq/Kg dry	± 10.0	Bq/Kg dry	67.0	Cs137	8.0 Bq/Kg dry
			Cs134	27.0	Bq/Kg dry	± 8.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Takaku, Chuodai, Iwaki	Apr-12	Cs137	801	Bq/Kg dry	± 86.0	Bq/Kg dry	1,310	Cs137	8.0 Bq/Kg dry
			Cs134	509	Bq/Kg dry	± 53.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Iino, Chuodai, Iwaki	Apr-12	Cs137	228	Bq/Kg dry	± 32.0	Bq/Kg dry	400	Cs137	8.0 Bq/Kg dry
			Cs134	172	Bq/Kg dry	± 22.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Koya, Uchigo, Iwaki	Apr-12	Cs137	178	Bq/Kg dry	± 26.0	Bq/Kg dry	303	Cs137	8.0 Bq/Kg dry
			Cs134	125	Bq/Kg dry	± 17.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Miya, Uchigo, Iwaki	Apr-12	Cs137	262	Bq/Kg dry	± 34.0	Bq/Kg dry	425	Cs137	8.0 Bq/Kg dry
			Cs134	163	Bq/Kg dry	± 21.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Okaona, Onahama, Iwaki	Apr-12	Cs137	79.0	Bq/Kg dry	± 15.0	Bq/Kg dry	133	Cs137	8.0 Bq/Kg dry
			Cs134	54.0	Bq/Kg dry	± 10.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Sumiyoshi, Onahama, Iwaki	Apr-12	Cs137	100	Bq/Kg dry	± 18.0	Bq/Kg dry	155	Cs137	8.0 Bq/Kg dry
			Cs134	55.0	Bq/Kg dry	± 11.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Sumiyoshi, Onahama, Iwaki	Apr-12	Cs137	66.0	Bq/Kg dry	± 14.0	Bq/Kg dry	102	Cs137	8.0 Bq/Kg dry
			Cs134	36.0	Bq/Kg dry	± 8.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	202	Bq/Kg dry	± 28.0	Bq/Kg dry	317	Cs137	8.0 Bq/Kg dry
			Cs134	115	Bq/Kg dry	± 17.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Kamikajiro, Onahama, Iwaki	Apr-12	Cs137	81.0	Bq/Kg dry	± 15.0	Bq/Kg dry	137	Cs137	8.0 Bq/Kg dry
			Cs134	56.0	Bq/Kg dry	± 10.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Nishigo, Joban, Iwaki	Apr-12	Cs137	82.0	Bq/Kg dry	± 17.0	Bq/Kg dry	143	Cs137	8.0 Bq/Kg dry
			Cs134	61.0	Bq/Kg dry	± 11.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Misawa, Joban, Iwaki	Mar-12	Cs137	201	Bq/Kg dry	± 28.0	Bq/Kg dry	330	Cs137	8.0 Bq/Kg dry
			Cs134	129	Bq/Kg dry	± 17.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Moegidai, Izumi, Iwaki	Mar-12	Cs137	105	Bq/Kg dry	± 17.0	Bq/Kg dry	177	Cs137	8.0 Bq/Kg dry
			Cs134	72.0	Bq/Kg dry	± 11.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Tamatsuyu, Izumi, Iwaki	Apr-12	Cs137	128	Bq/Kg dry	± 21.0	Bq/Kg dry	220	Cs137	8.0 Bq/Kg dry
			Cs134	92.0	Bq/Kg dry	± 14.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Yamada, Iwaki	Apr-12	Cs137	190	Bq/Kg dry	± 42.0	Bq/Kg dry	306	Cs137	8.0 Bq/Kg dry
			Cs134	116	Bq/Kg dry	± 26.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Shiroyone, Nakoso, Iwaki	Apr-12	Cs137	328	Bq/Kg dry	± 42.0	Bq/Kg dry	543	Cs137	8.0 Bq/Kg dry
			Cs134	215	Bq/Kg dry	± 26.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Paddy field soil	Nakadera, Miwa, Iwaki	Apr-12	Cs137	149	Bq/Kg dry	± 24.0	Bq/Kg dry	258	Cs137	8.0 Bq/Kg dry
			Cs134	109	Bq/Kg dry	± 15.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Paddy field soil	Manome, Taira, Iwaki	Apr-12	Cs137	395	Bq/Kg dry	± 49.0	Bq/Kg dry	654	Cs137	8.0 Bq/Kg dry
			Cs134	259	Bq/Kg dry	± 30.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry