



Radiation Measurement Results of 302 items in May 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Kurihara, Miyagi	Oct-11	Cs137	0.0	Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Higashimatsush ima, Miyagi	Oct-11	Cs137	2.2	Bq/Kg raw	4.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Fukui	Oct-11	Cs137	3.6	Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Fukushima	Oct-11	Cs137	2.8	Bq/Kg raw	6.8	Cs137	7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Naraha, Futaba	Oct-11	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Ishikawa, Ishikawa	Oct-11	Cs137	5.6	Bq/Kg raw	5.6	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Imaniida, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kashima, Iwaki	Oct-11	Cs137	2.5	Bq/Kg raw	5.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kashima, Iwaki	Oct-11	Cs137	3.1	Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Iwaki	Oct-11	Cs137	5.7	Bq/Kg raw	8.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Iwaki	Oct-11	Cs137	3.7	Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Unknown	Oct-11	Cs137	4.2	Bq/Kg raw	6.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Hokkaido	Oct-11	Cs137	0.0	Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Hokkaido	Oct-11	Cs137	2.5	Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Akita	Oct-11	Cs137	0.0	Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Akita	Oct-11	Cs137	3.0	Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Yamagata	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Marumori, Igu, Miyagi	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.3	Cs137 7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Niigata	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.7	Cs137 7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Mie	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tochigi	Oct-11	Cs137	2.9	Bq/Kg raw	± 2.2	Bq/Kg raw	5.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ibaraki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ibaraki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ibaraki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nakamachi, Naka, Ibaraki	Oct-11	Cs137	2.3	Bq/Kg raw	± 2.3	Bq/Kg raw	5.6	Cs137 7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Fukushima	Oct-10	Cs137	2.1	Bq/Kg raw	± 2.1	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Fukushima	Oct-11	Cs137	2.9	Bq/Kg raw	± 2.2	Bq/Kg raw	5.2	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tadami, Minamiaizu	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Aizu	Unknown	Cs137	2.8	Bq/Kg raw	± 2.1	Bq/Kg raw	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Aizu	Unknown	Cs137	2.8	Bq/Kg raw	± 2.1	Bq/Kg raw	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Sukagawa	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ono, Tamura	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.3	Cs137 7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ishikawa, Ishikawa	Oct-11	Cs137	2.4	Bq/Kg raw	± 1.9	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Godo, Miwa, Iwaki	Oct-11	Cs137	2.5	Bq/Kg raw	± 2.1	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ogawa, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kamihirakubo, Taira, Iwaki	Oct-11	Cs137	3.4	Bq/Kg raw	± 2.2	Bq/Kg raw	5.7	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kinuya, Taira, Iwaki	Oct-11	Cs137	2.5	Bq/Kg raw	± 2.3	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nakashio, Taira, Iwaki	Oct-11	Cs137	3.5	Bq/Kg raw	± 2.1	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Hosoya, Yotsukura, Iwaki	Nov-11	Cs137	3.5	Bq/Kg raw	± 2.0	Bq/Kg raw	5.8	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Yoshima, Iwaki	Oct-11	Cs137	3.9	Bq/Kg raw	± 2.1	3.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Kashima, Iwaki	Oct-11	Cs137	3.9	Bq/Kg raw	± 2.1	6.3	Cs137	7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
White rice	Kashima, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
White rice	Mizunoya, Joban, Iwaki	Oct-11	Cs137	5.5	Bq/Kg raw	± 2.7	8.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
White rice	Reiba, Takakura, Iwaki	Nov-11	Cs137	6.4	Bq/Kg raw	± 2.8	13.3	Cs137	7.0 Bq/Kg raw
			Cs134	6.9	Bq/Kg raw	± 2.4		Cs134	6.0 Bq/Kg raw
White rice	Izumi, Iwaki	Oct-11	Cs137	2.2	Bq/Kg raw	± 2.1	4.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
White rice	Honya, Izumi, Iwaki	Oct-11	Cs137	1.8	Bq/Kg raw	± 1.5	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.3		Cs134	6.0 Bq/Kg raw
White rice	Numabe, Iwaki	Oct-11	Cs137	3.8	Bq/Kg raw	± 2.2	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Tono, Iwaki	Oct-11	Cs137	4.3	Bq/Kg raw	± 2.2	4.3	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Yamada, Iwaki	Oct-11	Cs137	3.2	Bq/Kg raw	± 2.1	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Yamada, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Yamada, Iwaki	Nov-11	Cs137	4.4	Bq/Kg raw	± 2.3	8.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
White rice	Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	3.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
White rice	Iwaki	Unknown	Cs137	3.0	Bq/Kg raw	± 2.2	6.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
White rice	Iwaki	Oct-11	Cs137	3.5	Bq/Kg raw	± 2.1	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Unknown	Oct-11	Cs137	2.5	Bq/Kg raw	± 2.1	2.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Unknown	Unknown	Cs137	3.3	Bq/Kg raw	± 2.4	3.3	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Honya, Izumi, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Yamada, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
Pre-washed white rice	Misawa, Iwaki	Oct-11	Cs137	2.2	Bq/Kg raw	± 2.2	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pre-washed white rice	Misawa, Iwaki	Oct-11	Cs137	2.2	Bq/Kg raw	± 2.2	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Sticky rice	Ono, Tamura	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.7		Cs134	6.0 Bq/Kg raw
Sticky rice	Akai, Taira, Iwaki	Oct-11	Cs137	4.6	Bq/Kg raw	± 2.4	8.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Sticky rice	Akita	Oct-11	Cs137	3.6	Bq/Kg raw	± 2.4	6.0	Cs137	7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
Fried rice	Kakuda, Miyagi	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Kamiyoshima, Yoshima, Iwaki	Jul-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potatoes	Kitsunezuka, Yotsukura, Iwaki	Jul-11	Cs137	7.8	Bq/Kg raw	± 2.9	11.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Onions	Kamiyoshima, Yoshima, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onions	Makunouchi, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onions	Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Onions	Akai, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onions	Shimoogoe, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.4		Cs134	6.0 Bq/Kg raw
Onions	Shimoarakawa, Taira, Iwaki	May-12	Cs137	3.1	Bq/Kg raw	± 2.0	3.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onions	Shimokajiro, Onahama, Iwaki	May-11	Cs137	3.9	Bq/Kg raw	± 2.8	3.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onions	Shimokajiro, Onahama, Iwaki	May-11	Cs137	3.9	Bq/Kg raw	± 2.8	3.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onions	Yamada, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Spring onions	Kitsunezuka, Yotsukura, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	4.3	Cs137	7.0 Bq/Kg raw
			Cs134	4.3	Bq/Kg raw	± 3.9		Cs134	6.0 Bq/Kg raw
Leeks	Okubo, Ohisa, Iwaki	May-12	Cs137	4.5	Bq/Kg raw	± 3.9	4.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Leeks	Nakashio, Taira, Iwaki	Oct-11	Cs137	3.6	Bq/Kg raw	± 2.6	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Leeks	Suginami, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Leeks	Kashima, Iwaki	May-12	Cs137	2.5	Bq/Kg raw	± 2.0	2.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Leeks	Okaona, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Leeks	Miyamada, Tono, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Scallion	Jinba, Ohisa, Iwaki	May-12	Cs137	4.3	Bq/Kg raw	± 2.7	7.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.4		Cs134	6.0 Bq/Kg raw
Myoga	Numabe, Iwaki	May-12	Cs137	7.1	Bq/Kg raw	± 3.9	11.1	Cs137	7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 3.1		Cs134	6.0 Bq/Kg raw
Myoga	Ohisa, Ohisa, Iwaki	May-12	Cs137	5.8	Bq/Kg raw	± 4.1	9.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 3.7		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Japanese radish	Shimoarakawa, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese radish	Shimokajiro, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese radish	Kamikajiro, Onahama, Iwaki	May-12	Cs137	3.5	Bq/Kg raw	± 2.0	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese radish	Kamiyada, Joban, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Turnip	Shimoarakawa, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Turnip	Kamikuramochi, Kashima, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Turnip	Shimogawa, Izumi, Iwaki	May-12	Cs137	5.6	Bq/Kg raw	± 3.5	5.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Red turnip	Onahama, Iwaki	May-12	Cs137	3.9	Bq/Kg raw	± 2.9	3.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Shimoyoshima, Yoshima, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Shimoogoe, Taira, Iwaki	May-12	Cs137	5.6	Bq/Kg raw	± 4.7	5.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Jiyugaoka, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Suginami, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Shimoarakawa, Taira, Iwaki	May-12	Cs137	5.5	Bq/Kg raw	± 5.3	5.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Shimokajiro, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Okaona, Onahama, Iwaki	May-12	Cs137	3.6	Bq/Kg raw	± 3.1	3.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Watanabe, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Cabbages	Unknown	Unknown	Cs137	2.1	Bq/Kg raw	± 1.8	4.5	Cs137 8.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 4.7		Cs134 8.0 Bq/Kg raw
Cabbages (boiled)	Unknown	Unknown	Cs137	2.4	Bq/Kg raw	± 4.4	5.0	Cs137 8.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 4.4		Cs134 8.0 Bq/Kg raw
Asparagus	Kawamae, Iwaki	May-12	Cs137	3.9	Bq/Kg raw	± 2.5	7.6	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.2		Cs134 6.0 Bq/Kg raw
Asparagus	Kamiyoshima, Yoshima, Iwaki	May-12	Cs137	2.7	Bq/Kg raw	± 2.2	2.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Broccoli	Kamikajiro, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Broccoli	Unkonwn	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Chinese chive	Okubo, Ohisa, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 4.3		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Chinese chive	Watanabe, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Chinese chive	Ohara, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Chinese chive	Shimonagai, Miwa, Iwaki	May-12	Cs137	5.3	Bq/Kg raw	± 3.6	14.1	Cs137 8.0 Bq/Kg raw
			Cs134	8.8	Bq/Kg raw	± 7.8		Cs134 8.0 Bq/Kg raw
Chinese chive	Shimoogoe, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Chinese chive	Yamada, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Chinese chive	Okaona, Onahama, Iwaki	May-12	Cs137	2.0	Bq/Kg raw	± 1.9	2.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Vitamin-na	Kitsunezuka, Yotsukura, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.3		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Kitsunezuka, Yotsukura, Iwaki	May-12	Cs137	3.8	Bq/Kg raw	± 3.2	6.5	Cs137 7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 2.7		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Mizunoya, Joban, Iwaki	May-12	Cs137	5.8	Bq/Kg raw	± 3.5	5.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Kamiyada, Joban, Iwaki	May-12	Cs137	5.0	Bq/Kg raw	± 4.9	5.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Kashima, Iwaki	May-12	Cs137	4.0	Bq/Kg raw	± 3.0	8.8	Cs137 7.0 Bq/Kg raw
			Cs134	4.8	Bq/Kg raw	± 2.6		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Shima, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Shimogawa, Izumi, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Unknown	Unknown	Cs137	3.7	Bq/Kg raw	± 2.9	6.3	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.5		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	4.0	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 3.0		Cs134 6.0 Bq/Kg raw
Spinach	Kitsunezuka, Yotsukura, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Spinach	Shima, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Spinach	Okaona, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Spinach	Unknown	Unknown	Cs137	3.6	Bq/Kg raw	± 2.4	6.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.1		Cs134 6.0 Bq/Kg raw
Ashitaba	Kashimayama, Koriyama	Apr-12	Cs137	3.8	Bq/Kg raw	± 2.5	7.3	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.1		Cs134 6.0 Bq/Kg raw
Mitsuba	Shimokajiro, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134 6.0 Bq/Kg raw
Green chillies	Shimogawa, Izumi, Iwaki	Sep-11	Cs137	3.1	Bq/Kg raw	± 2.3	5.2	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.9		Cs134 6.0 Bq/Kg raw
Apple mint	Shimokajiro, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	5.6	Cs137 7.0 Bq/Kg raw
			Cs134	5.6	Bq/Kg raw	± 4.0		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Snap peas	Oyasaku, Yoshima, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 3.1		Cs134	6.0 Bq/Kg raw
Snap peas	Kamikajiro, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.5	Cs137	7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 2.4		Cs134	6.0 Bq/Kg raw
Snap peas	Yamada, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	5.5	Cs137	7.0 Bq/Kg raw
			Cs134	5.5	Bq/Kg raw	± 3.5		Cs134	6.0 Bq/Kg raw
Butterbur	Naraha, Futaba	May-12	Cs137	99.5	Bq/Kg raw	± 21.6	164	Cs137	7.0 Bq/Kg raw
			Cs134	64.1	Bq/Kg raw	± 13.9		Cs134	6.0 Bq/Kg raw
Butterbur	Kawamae, Iwaki	May-12	Cs137	33.6	Bq/Kg raw	± 9.3	70.6	Cs137	7.0 Bq/Kg raw
			Cs134	37.0	Bq/Kg raw	± 8.8		Cs134	6.0 Bq/Kg raw
Butterbur	Oyasaku, Yoshima, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.9		Cs134	6.0 Bq/Kg raw
Butterbur	Kitashirado, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Butterbur	Midaisakai, Uchigo, Iwaki	Unknown	Cs137	7.2	Bq/Kg raw	± 4.7	11.8	Cs137	7.0 Bq/Kg raw
			Cs134	4.6	Bq/Kg raw	± 3.9		Cs134	6.0 Bq/Kg raw
Butterbur	Yamada, Iwaki	May-12	Cs137	5.5	Bq/Kg raw	± 3.9	5.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Butterbur	Tabito, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Butterbur	Tabito, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	5.1	Cs137	7.0 Bq/Kg raw
			Cs134	5.1	Bq/Kg raw	± 3.6		Cs134	6.0 Bq/Kg raw
Butterbur	Shiroyone, Nakoso, Iwaki	May-12	Cs137	10.1	Bq/Kg raw	± 5.9	26.2	Cs137	7.0 Bq/Kg raw
			Cs134	16.1	Bq/Kg raw	± 5.5		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Shibahara, Ogawa, Iwaki	May-12	Cs137	3.2	Bq/Kg raw	± 2.7	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Shimoyamauchi, Taira, Iwaki	May-12	Cs137	7.0	Bq/Kg raw	± 2.6	12.7	Cs137	7.0 Bq/Kg raw
			Cs134	5.7	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Motoyashiki, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Kanari, Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Tabito, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
Butterbur	Tenei, Iwase	Apr-12	Cs137	14.1	Bq/Kg raw	± 6.7	32.0	Cs137	8.0 Bq/Kg raw
			Cs134	17.9	Bq/Kg raw	± 9.3		Cs134	8.0 Bq/Kg raw
Udo	Shidamyo, Kawamae, Iwaki	May-12	Cs137	73.5	Bq/Kg raw	± 16.4	121	Cs137	7.0 Bq/Kg raw
			Cs134	47.8	Bq/Kg raw	± 10.7		Cs134	6.0 Bq/Kg raw
Udo	Yotsukura, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Udo	Kitsunezuka, Yotsukura, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	5.7	Cs137	7.0 Bq/Kg raw
			Cs134	5.7	Bq/Kg raw	± 5.4		Cs134	6.0 Bq/Kg raw
Udo	Godo, Miwa, Iwaki	May-12	Cs137	5.2	Bq/Kg raw	± 2.8	5.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Udo	Shimogawa, Izumi, Iwaki	May-12	Cs137	3.0	Bq/Kg raw	± 2.4	5.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Fatsia Sprouts	Sekimoto, Kitaibaraki	May-12	Cs137	4.7	Bq/Kg raw	± 2.4	7.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 2.1		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Samegawa, Higashishirakawa	May-12	Cs137	67.3	Bq/Kg raw	± 16.3	139	Cs137	7.0	Bq/Kg raw
			Cs134	71.6	Bq/Kg raw	± 15.6		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Watarasekoemushi, Samegawa, Higashishirakawa	May-12	Cs137	45.4	Bq/Kg raw	± 10.2	78.5	Cs137	7.0	Bq/Kg raw
			Cs134	33.1	Bq/Kg raw	± 7.2		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Ide, Naraha, Futaba	May-15	Cs137	105	Bq/Kg raw	± 7.0	181	Cs137	7.0	Bq/Kg raw
			Cs134	75.6	Bq/Kg raw	± 6.0		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Godo, Miwa, Iwaki	May-12	Cs137	5.6	Bq/Kg raw	± 3.8	9.5	Cs137	7.0	Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 3.2		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Miwa, Iwaki	May-12	Cs137	184	Bq/Kg raw	± 38.0	322	Cs137	7.0	Bq/Kg raw
			Cs134	138	Bq/Kg raw	± 28.0		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Shima, Onahama, Iwaki	May-12	Cs137	8.7	Bq/Kg raw	± 5.3	14.9	Cs137	7.0	Bq/Kg raw
			Cs134	6.2	Bq/Kg raw	± 4.5		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Shima, Onahama, Iwaki	May-12	Cs137	8.7	Bq/Kg raw	± 5.3	14.9	Cs137	7.0	Bq/Kg raw
			Cs134	6.2	Bq/Kg raw	± 4.5		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Tono, Iwaki	May-12	Cs137	7.7	Bq/Kg raw	± 4.0	14.0	Cs137	7.0	Bq/Kg raw
			Cs134	6.3	Bq/Kg raw	± 3.3		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Tono, Iwaki	Apr-12	Cs137	58.5	Bq/Kg raw	± 14.0	116	Cs137	7.0	Bq/Kg raw
			Cs134	57.2	Bq/Kg raw	± 12.5		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Tabito, Iwaki	Apr-12	Cs137	138	Bq/Kg raw	± 30.0	247	Cs137	7.0	Bq/Kg raw
			Cs134	109	Bq/Kg raw	± 23.0		Cs134	6.0	Bq/Kg raw
Fatsia Sprouts	Nakoso, Iwaki	Apr-12	Cs137	61.7	Bq/Kg raw	± 13.4	107	Cs137	7.0	Bq/Kg raw
			Cs134	45.5	Bq/Kg raw	± 9.6		Cs134	6.0	Bq/Kg raw
Bracken	Watarasekoemushi, Samegawa, Higashishirakawa	May-12	Cs137	3.9	Bq/Kg raw	± 2.0	7.6	Cs137	7.0	Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 1.8		Cs134	6.0	Bq/Kg raw
Bracken	Samegawa, Higashishirakawa	May-12	Cs137	24.4	Bq/Kg raw	± 6.7	44.0	Cs137	7.0	Bq/Kg raw
			Cs134	19.6	Bq/Kg raw	± 5.2		Cs134	6.0	Bq/Kg raw
Bracken	Miwa, Iwaki	May-12	Cs137	2.7	Bq/Kg raw	± 2.5	6.3	Cs137	7.0	Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.1		Cs134	6.0	Bq/Kg raw
Bracken	Miwa, Iwaki	May-12	Cs137	2.7	Bq/Kg raw	± 2.5	6.3	Cs137	7.0	Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.1		Cs134	6.0	Bq/Kg raw
Bracken	Shidamyo, Kawamae, Iwaki	May-12	Cs137	126	Bq/Kg raw	± 27.0	231	Cs137	7.0	Bq/Kg raw
			Cs134	105	Bq/Kg raw	± 21.0		Cs134	6.0	Bq/Kg raw
Bracken	Kitayoshima, Yoshima, Iwaki	Apr-12	Cs137	10.6	Bq/Kg raw	± 4.5	20.8	Cs137	7.0	Bq/Kg raw
			Cs134	10.2	Bq/Kg raw	± 3.8		Cs134	6.0	Bq/Kg raw
Bracken	Shimogawa, Izumi, Iwaki	May-12	Cs137	12.6	Bq/Kg raw	± 4.0	21.8	Cs137	7.0	Bq/Kg raw
			Cs134	9.2	Bq/Kg raw	± 3.0		Cs134	6.0	Bq/Kg raw
Bracken	Ueda, Iwaki	Apr-12	Cs137	9.5	Bq/Kg raw	± 3.3	15.5	Cs137	7.0	Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 2.4		Cs134	6.0	Bq/Kg raw
Bracken	Yamada, Iwaki	May-12	Cs137	3.7	Bq/Kg raw	± 3.5	3.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0	Bq/Kg raw
Bracken	Tono, Iwaki	Apr-12	Cs137	3.5	Bq/Kg raw	± 2.2	6.7	Cs137	7.0	Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.9		Cs134	6.0	Bq/Kg raw
Bracken	Tono, Iwaki	Apr-12	Cs137	3.5	Bq/Kg raw	± 2.2	6.7	Cs137	7.0	Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.9		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Bracken	Tabiuto, Tabito, Iwaki	May-12	Cs137	58.6	Bq/Kg raw	± 13.8	Bq/Kg raw	104	Cs137 7.0 Bq/Kg raw
			Cs134	45.0	Bq/Kg raw	± 10.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken (boiled)	Shimonagai, Miwa, Iwaki (top)	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken (boiled)	Shimonagai, Miwa, Iwaki (bottom)	May-12	Cs137	4.1	Bq/Kg raw	± 2.1	Bq/Kg raw	4.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken (boiled)	Nagasaki, Iwaki	May-12	Cs137	2.7	Bq/Kg raw	± 2.0	Bq/Kg raw	5.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bracken (preserved)	Iwaki	May-08	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.7	Cs137 7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shidoki	Tono, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Kogomi	Tadami, Minamiaizu	May-12	Cs137	59.0	Bq/Kg raw	± 14.1	Bq/Kg raw	110.0	Cs137 7.0 Bq/Kg raw
			Cs134	51.0	Bq/Kg raw	± 11.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Koshiabura	Saiso, Miwa, Iwaki	May-12	Cs137	105	Bq/Kg raw	± 23.0	Bq/Kg raw	190	Cs137 7.0 Bq/Kg raw
			Cs134	84.6	Bq/Kg raw	± 17.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Tomioka, Futaba	May-12	Cs137	31.2	Bq/Kg raw	± 7.9	Bq/Kg raw	56.3	Cs137 7.0 Bq/Kg raw
			Cs134	25.1	Bq/Kg raw	± 6.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Ide, Naraha, Futaba	May-12	Cs137	190	Bq/Kg raw	± 38.0	Bq/Kg raw	331	Cs137 7.0 Bq/Kg raw
			Cs134	141	Bq/Kg raw	± 28.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shibahara, Ogawa, Iwaki	May-12	Cs137	4.5	Bq/Kg raw	± 2.5	Bq/Kg raw	7.9	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Kawanago, Yoshima, Iwaki	May-12	Cs137	11.4	Bq/Kg raw	± 4.5	Bq/Kg raw	23.0	Cs137 7.0 Bq/Kg raw
			Cs134	11.6	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Komagome, Yotsukura, Iwaki	Apr-12	Cs137	2.5	Bq/Kg raw	± 2.5	Bq/Kg raw	6.0	Cs137 7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimoyamauchi, Taira, Iwaki	May-12	Cs137	74.6	Bq/Kg raw	± 16.4	Bq/Kg raw	135.3	Cs137 7.0 Bq/Kg raw
			Cs134	60.7	Bq/Kg raw	± 12.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Suginami, Taira, Iwaki	May-12	Cs137	94.5	Bq/Kg raw	± 20.3	Bq/Kg raw	163	Cs137 7.0 Bq/Kg raw
			Cs134	68.7	Bq/Kg raw	± 14.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Makunouchi, Taira, Iwaki	May-12	Cs137	28.9	Bq/Kg raw	± 7.4	Bq/Kg raw	49.6	Cs137 7.0 Bq/Kg raw
			Cs134	20.7	Bq/Kg raw	± 5.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Rokunincho, Taira, Iwaki	May-12	Cs137	22.3	Bq/Kg raw	± 6.2	Bq/Kg raw	37.1	Cs137 7.0 Bq/Kg raw
			Cs134	14.8	Bq/Kg raw	± 4.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Yagawase, taira, Iwaki	Apr-12	Cs137	42.5	Bq/Kg raw	± 10.9	Bq/Kg raw	75.3	Cs137 7.0 Bq/Kg raw
			Cs134	32.8	Bq/Kg raw	± 7.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Joban, Iwaki	May-12	Cs137	47.2	Bq/Kg raw	± 11.0	Bq/Kg raw	87.6	Cs137 7.0 Bq/Kg raw
			Cs134	40.4	Bq/Kg raw	± 8.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Yumoto, Joban, Iwaki	May-12	Cs137	15.3	Bq/Kg raw	± 4.5	Bq/Kg raw	31.7	Cs137 7.0 Bq/Kg raw
			Cs134	16.4	Bq/Kg raw	± 4.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Kamiyunagaya, Joban, Iwaki	May-12	Cs137	14.0	Bq/Kg raw	± 4.3	Bq/Kg raw	25.0	Cs137 7.0 Bq/Kg raw
			Cs134	11.0	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Miyo, Kashima, Iwaki	Apr-12	Cs137	4.3	Bq/Kg raw	± 9.5	Bq/Kg raw	8.8	Cs137 8.0 Bq/Kg raw
			Cs134	4.5	Bq/Kg raw	± 5.1	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Bamboo shoot	Miyo, Kashima, Iwaki	Apr-12	Cs137	15.9	Bq/Kg raw	± 4.2	Bq/Kg raw	24.7	Cs137 8.0 Bq/Kg raw
			Cs134	8.8	Bq/Kg raw	± 5.6	Bq/Kg raw		Cs134 8.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Bamboo shoot	Noda, Onahama, Iwaki	Apr-12	Cs137	112	Bq/Kg raw	± 24.0	Bq/Kg raw	196	Cs137 7.0 Bq/Kg raw
			Cs134	83.7	Bq/Kg raw	± 17.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Okaona, Onahama, Iwaki	May-12	Cs137	21.0	Bq/Kg raw	± 6.3	Bq/Kg raw	41.6	Cs137 7.0 Bq/Kg raw
			Cs134	20.6	Bq/Kg raw	± 5.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimokajiro, Onahama, Iwaki	May-12	Cs137	107	Bq/Kg raw	± 22.0	Bq/Kg raw	196	Cs137 7.0 Bq/Kg raw
			Cs134	89.1	Bq/Kg raw	± 17.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Kaneyama, Numabe, Iwaki	May-12	Cs137	54.5	Bq/Kg raw	± 12.3	Bq/Kg raw	91.0	Cs137 7.0 Bq/Kg raw
			Cs134	36.5	Bq/Kg raw	± 8.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Reiba, Takakura, Iwaki	Apr-12	Cs137	6.6	Bq/Kg raw	± 2.4	Bq/Kg raw	9.8	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Obama, Iwaki	May-12	Cs137	69.3	Bq/Kg raw	± 15.4	Bq/Kg raw	125	Cs137 7.0 Bq/Kg raw
			Cs134	55.7	Bq/Kg raw	± 11.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shimogawa, Izumi, Iwaki	May-12	Cs137	41.9	Bq/Kg raw	± 9.5	Bq/Kg raw	69.6	Cs137 7.0 Bq/Kg raw
			Cs134	27.7	Bq/Kg raw	± 6.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Honya, Izumi, Iwaki	Apr-12	Cs137	17.8	Bq/Kg raw	± 4.6	Bq/Kg raw	30.4	Cs137 7.0 Bq/Kg raw
			Cs134	12.6	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Yamadama, Iwaki	Apr-12	Cs137	45.1	Bq/Kg raw	± 10.4	Bq/Kg raw	83.7	Cs137 7.0 Bq/Kg raw
			Cs134	38.6	Bq/Kg raw	± 8.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Watanabe, Iwaki	May-12	Cs137	40.1	Bq/Kg raw	± 9.6	Bq/Kg raw	70.3	Cs137 7.0 Bq/Kg raw
			Cs134	30.2	Bq/Kg raw	± 7.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Kamado, Watanabe, Iwaki	May-12	Cs137	30.0	Bq/Kg raw	± 7.5	Bq/Kg raw	54.4	Cs137 7.0 Bq/Kg raw
			Cs134	24.4	Bq/Kg raw	± 5.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Yamadama, Iwaki	May-12	Cs137	22.3	Bq/Kg raw	± 6.6	Bq/Kg raw	44.2	Cs137 7.0 Bq/Kg raw
			Cs134	21.9	Bq/Kg raw	± 5.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Idekura, Nakoso, Iwaki	May-12	Cs137	57.7	Bq/Kg raw	± 12.6	Bq/Kg raw	99.9	Cs137 7.0 Bq/Kg raw
			Cs134	42.2	Bq/Kg raw	± 9.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot	Shiroyone, Nakoso, Iwaki	May-12	Cs137	54.7	Bq/Kg raw	± 12.5	Bq/Kg raw	99.2	Cs137 7.0 Bq/Kg raw
			Cs134	44.5	Bq/Kg raw	± 9.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Ibaraki	May-12	Cs137	19.5	Bq/Kg raw	± 5.1	Bq/Kg raw	32.0	Cs137 7.0 Bq/Kg raw
			Cs134	12.5	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Ibaraki	May-12	Cs137	19.5	Bq/Kg raw	± 5.1	Bq/Kg raw	32.0	Cs137 7.0 Bq/Kg raw
			Cs134	12.5	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Ogoe, Tamura	May-12	Cs137	13.8	Bq/Kg raw	± 4.1	Bq/Kg raw	17.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Rokunincho, Taira, Iwaki	May-12	Cs137	13.4	Bq/Kg raw	± 3.9	Bq/Kg raw	24.8	Cs137 7.0 Bq/Kg raw
			Cs134	11.4	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Shimoyamauchi, Taira, Iwaki	May-12	Cs137	41.2	Bq/Kg raw	± 9.3	Bq/Kg raw	73.2	Cs137 7.0 Bq/Kg raw
			Cs134	32.0	Bq/Kg raw	± 6.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Koya, Uchigo, Iwaki	May-12	Cs137	72.7	Bq/Kg raw	± 15.6	Bq/Kg raw	134	Cs137 7.0 Bq/Kg raw
			Cs134	60.9	Bq/Kg raw	± 12.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Joban, Iwaki	Apr-12	Cs137	16.0	Bq/Kg raw	± 4.3	Bq/Kg raw	26.2	Cs137 7.0 Bq/Kg raw
			Cs134	10.2	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Kamiyunagaya, Joban, Iwaki	May-12	Cs137	4.2	Bq/Kg raw	± 2.4	Bq/Kg raw	11.5	Cs137 7.0 Bq/Kg raw
			Cs134	7.3	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bamboo shoot (boiled)	Nagasaki, Iwaki	May-12	Cs137	36.5	Bq/Kg raw	± 8.6	Bq/Kg raw	60.9	Cs137 7.0 Bq/Kg raw
			Cs134	24.4	Bq/Kg raw	± 5.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Bamboo shoot (boiled)	Kanari, Onahama, Iwaki	May-12	Cs137	59.3	Bq/Kg raw	± 13.0	Bq/Kg raw	112	Cs137	7.0 Bq/Kg raw
			Cs134	52.5	Bq/Kg raw	± 10.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot (boiled)	Soeno, Iwaki	Unknown	Cs137	11.6	Bq/Kg raw	± 3.6	Bq/Kg raw	15.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot (boiled)	Kaneyama, Numabe, Iwaki	May-12	Cs137	12.5	Bq/Kg raw	± 3.7	Bq/Kg raw	24.4	Cs137	7.0 Bq/Kg raw
			Cs134	11.9	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Bamboo shoot (boiled)	Yamadama, Iwaki	Apr-12	Cs137	22.2	Bq/Kg raw	± 5.5	Bq/Kg raw	38.0	Cs137	7.0 Bq/Kg raw
			Cs134	15.8	Bq/Kg raw	± 3.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Azuki beans	Takasaka, Uchigo, Iwaki	Aug-11	Cs137	6.4	Bq/Kg raw	± 3.3	Bq/Kg raw	10.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Black soybeans	Tabito, Iwaki	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushroom	Samegawa, Higashishirakawa	Apr-12	Cs137	108	Bq/Kg raw	± 23.0	Bq/Kg raw	195	Cs137	7.0 Bq/Kg raw
			Cs134	87.2	Bq/Kg raw	± 17.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushroom	Godō, Miwa, Iwaki	May-12	Cs137	81.9	Bq/Kg raw	± 17.2	Bq/Kg raw	141	Cs137	7.0 Bq/Kg raw
			Cs134	58.9	Bq/Kg raw	± 12.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushroom	Shimoyada, Kashima, Iwaki	May-12	Cs137	205	Bq/Kg raw	± 42.0	Bq/Kg raw	354	Cs137	7.0 Bq/Kg raw
			Cs134	149	Bq/Kg raw	± 30.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushroom	Onahama, Iwaki	Apr-12	Cs137	477	Bq/Kg raw	± 95.0	Bq/Kg raw	844	Cs137	7.0 Bq/Kg raw
			Cs134	367	Bq/Kg raw	± 73.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushroom	Yamada, Iwaki	Apr-12	Cs137	46.4	Bq/Kg raw	± 10.5	Bq/Kg raw	83.2	Cs137	7.0 Bq/Kg raw
			Cs134	36.8	Bq/Kg raw	± 8.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Shiitake mushroom	Nichibu, Tabito, Iwaki	Apr-12	Cs137	88.7	Bq/Kg raw	± 18.4	Bq/Kg raw	155	Cs137	7.0 Bq/Kg raw
			Cs134	66.4	Bq/Kg raw	± 13.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried shiitake mushroom	Shiratori, Joban, Iwaki	Mar-12	Cs137	1150	Bq/Kg raw	± 230	Bq/Kg raw	2,052	Cs137	7.0 Bq/Kg raw
			Cs134	902	Bq/Kg raw	± 180	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Yuzu	Numabe, Iwaki	Apr-12	Cs137	183	Bq/Kg raw	± 37.0	Bq/Kg raw	323	Cs137	7.0 Bq/Kg raw
			Cs134	140	Bq/Kg raw	± 28.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Dried persimmon	Ide, Naraha, Futaba	Oct-11	Cs137	360	Bq/Kg raw	± 72.0	Bq/Kg raw	638	Cs137	7.0 Bq/Kg raw
			Cs134	278	Bq/Kg raw	± 56.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Strawberries	Iwaki Sakuranbo Nursery	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137	6.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	4.0 Bq/Kg raw
Japanese bitterling	Nagasaki, Iwaki	May-12	Cs137	34.5	Bq/Kg raw	± 8.5	Bq/Kg raw	64.9	Cs137	7.0 Bq/Kg raw
			Cs134	30.4	Bq/Kg raw	± 7.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese bitterling	Onahama, Iwaki	May-12	Cs137	39.3	Bq/Kg raw	± 9.0	Bq/Kg raw	67.8	Cs137	7.0 Bq/Kg raw
			Cs134	28.5	Bq/Kg raw	± 6.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Japanese bitterling (bones and)	Onahama, Iwaki	May-12	Cs137	30.9	Bq/Kg raw	± 7.4	Bq/Kg raw	53.4	Cs137	7.0 Bq/Kg raw
			Cs134	22.5	Bq/Kg raw	± 5.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Hijiki	Toyoma, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Hijiki (boiled)	Toyoma, Iwaki	Apr-12	Cs137	15.0	Bq/Kg raw	± 4.9	Bq/Kg raw	20.2	Cs137	7.0 Bq/Kg raw
			Cs134	5.2	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Eggs	Kawauchi, Futaba	Unknown	Cs137	2.3	Bq/Kg raw	± 2.1	Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Breast milk	Onahama, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Pickled plums	Yamada, Iwaki	Jun-11	Cs137	49.5	Bq/Kg raw	± 10.9	Bq/Kg raw	96.7	Cs137 7.0 Bq/Kg raw
			Cs134	47.2	Bq/Kg raw	± 9.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Green chill Miso	Shimogawa, Izumi, Iwaki	Sep-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Honey	Ohisa, Ohisa, Fukushima	Nov-11	Cs137	9.3	Bq/Kg raw	± 3.5	Bq/Kg raw	23.1	Cs137 7.0 Bq/Kg raw
			Cs134	13.8	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Honey	Yotsukura, Iwaki	May-12	Cs137	15.1	Bq/Kg raw	± 4.0	Bq/Kg raw	25.1	Cs137 7.0 Bq/Kg raw
			Cs134	10.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Honey	Akai, Taira, Iwaki	May-12	Cs137	3.4	Bq/Kg raw	± 1.7	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 1.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Soybean paste	Minamisoma	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Soybean paste	Kawauchi, Futaba	Dec-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137 7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Soybean paste	Nakayama, Taira, Iwaki	Jan-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
School lunch	Uchigo, Iwaki	May-12	Cs137	1.9	Bq/Kg raw	± 1.8	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
School lunch	Uchigo, Iwaki	May-12	Cs137	2.5	Bq/Kg raw	± 1.9	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
School lunch	Joban, Iwaki	May-12	Cs137	1.7	Bq/Kg raw	± 1.6	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Kikuimo tea	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Underground water	Tabiuto, Tabito, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Ide, Naraha, Futaba	Apr-12	Cs137	1.1	Bq/Kg raw	± 1.0	Bq/Kg raw	1.1	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Furudono, Ishikawa	Apr-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137 5.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Mineral water	Tenryosui, Hita, Oita	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.0	Cs137 5.0 Bq/Kg raw
			Cs134	1.0	Bq/Kg raw	± 0.9	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Mineral water	Kagoshima	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Mineral water	Yakuishima, Kagoshima	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap water	Nishigo, Joban, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.2	Cs137 5.0 Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap water	Kuroda, Tabito, Iwaki	May-12	Cs137	0.9	Bq/Kg raw	± 0.9	Bq/Kg raw	0.9	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Akai, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137 5.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Shiratori, Joban, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Yamada, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Shiitake wood	Funaita, Yamada, Iwaki	May-12	Cs137	52.0	Bq/Kg raw	± 13.0	Bq/Kg raw	76.0	Cs137 7.0 Bq/Kg raw
			Cs134	24.0	Bq/Kg raw	± 8.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Ezoukogi	Kaneyama, Iwaki	May-12	Cs137	5.6	Bq/Kg raw	± 4.6	Bq/Kg raw	10.9	Cs137 7.0 Bq/Kg raw
			Cs134	5.3	Bq/Kg raw	± 4.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Rice husks	Tomitsu, Iwaki	May-12	Cs137	697	Bq/Kg raw	± 87.0	Bq/Kg raw	1,154	Cs137 8.0 Bq/Kg raw
			Cs134	457	Bq/Kg raw	± 54.0	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Rice husks	Numabe, Iwaki	May-12	Cs137	34.0	Bq/Kg raw	± 17.9	Bq/Kg raw	50.6	Cs137 7.0 Bq/Kg raw
			Cs134	16.6	Bq/Kg raw	± 14.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Compost	Kanuma, Tochigi	Unknown	Cs137	46.0	Bq/Kg dry	± 16.0	Bq/Kg dry	81.0	Cs137 8.0 Bq/Kg dry
			Cs134	35.0	Bq/Kg dry	± 10.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Leaf mold	Jiyugaoka, Iwaki	May-12	Cs137	2639	Bq/Kg dry	± 254	Bq/Kg dry	4,433	Cs137 8.0 Bq/Kg dry
			Cs134	1794	Bq/Kg dry	± 166	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Shibahara, Ogawa, Iawki	May-12	Cs137	111	Bq/Kg dry	± 19.0	Bq/Kg dry	190	Cs137 8.0 Bq/Kg dry
			Cs134	79.0	Bq/Kg dry	± 12.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Shimoogoe, Taira, Iwaki	May-12	Cs137	338	Bq/Kg dry	± 43.0	Bq/Kg dry	508	Cs137 8.0 Bq/Kg dry
			Cs134	170	Bq/Kg dry	± 23.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Shimoogoe, Taira, Iwaki	May-12	Cs137	224	Bq/Kg dry	± 33.0	Bq/Kg dry	382	Cs137 8.0 Bq/Kg dry
			Cs134	158	Bq/Kg dry	± 21.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Taira, Iwaki	May-12	Cs137	176	Bq/Kg dry	± 24.0	Bq/Kg dry	287	Cs137 8.0 Bq/Kg dry
			Cs134	111	Bq/Kg dry	± 15.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Taira, Iwaki	May-12	Cs137	161	Bq/Kg dry	± 24.0	Bq/Kg dry	268	Cs137 8.0 Bq/Kg dry
			Cs134	107	Bq/Kg dry	± 15.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Kashima, Iwaki	May-12	Cs137	227	Bq/Kg dry	± 32.0	Bq/Kg dry	393	Cs137 8.0 Bq/Kg dry
			Cs134	166	Bq/Kg dry	± 21.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Ohara, Onahama, Iwaki	May-12	Cs137	406	Bq/Kg dry	± 50.0	Bq/Kg dry	684	Cs137 8.0 Bq/Kg dry
			Cs134	278	Bq/Kg dry	± 32.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Shima, Onahama, Iwaki	May-12	Cs137	103	Bq/Kg dry	± 18.0	Bq/Kg dry	174	Cs137 8.0 Bq/Kg dry
			Cs134	71.0	Bq/Kg dry	± 12.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Takijiri, Izumi, Iwaki	May-12	Cs137	335	Bq/Kg dry	± 42.0	Bq/Kg dry	554	Cs137 8.0 Bq/Kg dry
			Cs134	219	Bq/Kg dry	± 26.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Numabe, Iwaki	May-12	Cs137	148	Bq/Kg dry	± 23.0	Bq/Kg dry	259	Cs137 8.0 Bq/Kg dry
			Cs134	111	Bq/Kg dry	± 15.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Nakadera, Miwa, Iwaki	May-12	Cs137	192	Bq/Kg dry	± 29.0	Bq/Kg dry	316	Cs137 8.0 Bq/Kg dry
			Cs134	124	Bq/Kg dry	± 18.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Oyasaku, Yoshima, Iwaki	May-12	Cs137	163	Bq/Kg dry	± 25.0	Bq/Kg dry	272	Cs137 8.0 Bq/Kg dry
			Cs134	109	Bq/Kg dry	± 16.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Ohisa, Iwaki	May-12	Cs137	72.0	Bq/Kg dry	± 15.0	Bq/Kg dry	121	Cs137 8.0 Bq/Kg dry
			Cs134	49.0	Bq/Kg dry	± 10.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Ohisa, Iwaki	May-12	Cs137	158	Bq/Kg dry	± 24.0	Bq/Kg dry	264	Cs137 8.0 Bq/Kg dry
			Cs134	106	Bq/Kg dry	± 15.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Ohisa, Ohisa, Iwaki	May-12	Cs137	54.0	Bq/Kg dry	± 15.0	Bq/Kg dry	92.0	Cs137 8.0 Bq/Kg dry
			Cs134	38.0	Bq/Kg dry	± 10.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Ohisa, Ohisa, Iwaki	May-12	Cs137	387	Bq/Kg dry	± 48.0	Bq/Kg dry	661	Cs137 8.0 Bq/Kg dry
			Cs134	274	Bq/Kg dry	± 31.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Hirakubo, Taira, Iwaki	Apr-12	Cs137	94.0	Bq/Kg dry	± 17.0	Bq/Kg dry	143	Cs137 8.0 Bq/Kg dry
			Cs134	49.0	Bq/Kg dry	± 11.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Soil	Kitashirado, Taira, Iwaki	May-12	Cs137	44.0	Bq/Kg dry	± 12.0	Bq/Kg dry	67.0	Cs137 8.0 Bq/Kg dry
			Cs134	23.0	Bq/Kg dry	± 7.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Kitashirado, Taira, Iwaki	May-12	Cs137	439	Bq/Kg dry	± 52.0	Bq/Kg dry	757	Cs137 8.0 Bq/Kg dry
			Cs134	318	Bq/Kg dry	± 34.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Kimigatsuka, Onahama, Iwak	May-12	Cs137	123	Bq/Kg dry	± 20.0	Bq/Kg dry	205	Cs137 8.0 Bq/Kg dry
			Cs134	82.0	Bq/Kg dry	± 13.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Mizunoya, Joban, Iwaki	May-12	Cs137	225	Bq/Kg dry	± 32.0	Bq/Kg dry	361	Cs137 8.0 Bq/Kg dry
			Cs134	136	Bq/Kg dry	± 19.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Ena, Iwaki	May-12	Cs137	119	Bq/Kg dry	± 20.0	Bq/Kg dry	216	Cs137 8.0 Bq/Kg dry
			Cs134	97.0	Bq/Kg dry	± 44.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Hananoi, Nishiki, Iwaki	May-12	Cs137	116	Bq/Kg dry	± 21.0	Bq/Kg dry	201	Cs137 8.0 Bq/Kg dry
			Cs134	85.0	Bq/Kg dry	± 14.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shiroyone, Nakoso, Iwaki	Apr-12	Cs137	318	Bq/Kg dry	± 42.0	Bq/Kg dry	536	Cs137 8.0 Bq/Kg dry
			Cs134	218	Bq/Kg dry	± 27.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Sumiyoshi, Onahama, Iwaki	May-12	Cs137	701	Bq/Kg dry	± 76.0	Bq/Kg dry	1,182	Cs137 8.0 Bq/Kg dry
			Cs134	481	Bq/Kg dry	± 49.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry

