



# Radiation Measurement Results of 255 items in June 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

**Radioactive contamination level may differ according to sampling points even within the same address.**

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Niigata	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Brown rice	Yuri, Akita	Nov-11	Cs137	2.8	Bq/Kg raw	± 2.2	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Brown rice	Futaba	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Aomori	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Ishinomaki, Miyagi	Oct-11	Cs137	2.1	Bq/Kg raw	± 1.9	4.4	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.7		Cs134	6.0 Bq/Kg raw
White rice	Tome, Miyagi	Oct-11	Cs137	4.0	Bq/Kg raw	± 2.0	4.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Ichinoseki, Iwate	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	2.4	Cs137	7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
White rice	Ichinoseki, Iwate	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.0	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
White rice	Tokushima	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.3		Cs134	6.0 Bq/Kg raw
White rice	Fukui	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	1.7	Cs137	7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.3		Cs134	6.0 Bq/Kg raw
White rice	Fukui	Oct-11	Cs137	2.2	Bq/Kg raw	± 1.8	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Niigata	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.0	Cs137	7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.7		Cs134	6.0 Bq/Kg raw
White rice	Uonuma, Niigata	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	2.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
White rice	Okayama	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Fukushima	Oct-10	Cs137	2.4	Bq/Kg raw	± 2.3	2.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
White rice	Ootama, Adachi	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	1.7	Cs137	6.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.4		Cs134	4.0 Bq/Kg raw
White rice	Kikuta, Koriyama	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Odaka, Minamisoma	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Oohisa, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.2	Cs137 7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ogawa, Iwaki	Unknown	Cs137	7.5	Bq/Kg raw	± 3.0	Bq/Kg raw	12.4	Cs137 7.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Makunouchi, Taira, Iwaki	Oct-11	Cs137	6.4	Bq/Kg raw	± 2.4	Bq/Kg raw	8.8	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Oogoe, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg raw	± 2.1	Bq/Kg raw	3.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Toyoma, Taira, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yoshima, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Osamago, joban, Iwaki	Oct-10	Cs137	3.5	Bq/Kg raw	± 2.2	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamada. Iwaki	Sep-10	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamada. Iwaki	Oct-11	Cs137	3.6	Bq/Kg raw	± 2.2	Bq/Kg raw	6.9	Cs137 7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Unknown	Unknown	Cs137	3.0	Bq/Kg raw	± 2.2	Bq/Kg raw	5.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Unknown	Unknown	Cs137	3.0	Bq/Kg raw	± 2.2	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sticky rice	Odaka, Minamisoma	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Ogawa, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Shioda, Ogawa, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Tamayama, Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Shimoyoshima, Yoshima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Kitakabeya, Taira, Iwaki	Jun-12	Cs137	2.5	Bq/Kg raw	± 2.3	Bq/Kg raw	2.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Shimoogoe, Taira, Iwaki	Jun-12	Cs137	2.9	Bq/Kg raw	± 2.1	Bq/Kg raw	2.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Shimoogoe, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Potato	Shimoogoe, Taira, Iwaki	Jun-12	Cs137	3.7	Bq/Kg raw	± 1.9	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Shimoyamaguchi , Taira, Iwaki	Jun-12	Cs137	2.3	Bq/Kg raw	± 2.2	Bq/Kg raw	2.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Numanouchi, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Kashima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137 8.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Potato	Onahama, Iwaki	Jun-12	Cs137	3.5	Bq/Kg raw	± 2.2	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Shimokajiro, Onahama, Iwaki	Jun-12	Cs137	4.6	Bq/Kg raw	± 2.8	Bq/Kg raw	4.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Kamikajiro, Onahama, Iwaki	Unknown	Cs137	2.4	Bq/Kg raw	± 2.3	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Kamikajiro, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Obama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Izumi, Iwaki	Jun-12	Cs137	2.2	Bq/Kg raw	± 1.9	Bq/Kg raw	2.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Watanabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Watanabe, Iwaki	Jun-12	Cs137	3.5	Bq/Kg raw	± 2.4	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Nishiki, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Kawabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato (with skin)	Suginami, Taira, Iwaki	Jun-12	Cs137	4.8	Bq/Kg raw	± 2.7	Bq/Kg raw	4.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato (with skin)	Shimoyunagaya, Joban, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onion	Ogawa, Iwaki	Jun-12	Cs137	3.0	Bq/Kg raw	± 2.5	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onion	Shioda, Ogawa, Iwaki	Jun-12	Cs137	5.2	Bq/Kg raw	± 3.1	Bq/Kg raw	5.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onion	Taira, Iwaki	Jun-12	Cs137	3.1	Bq/Kg raw	± 2.0	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onion	Hosoya, Yotsukura, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 2.1	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onion	Yotsukura, Iwaki	Jun-12	Cs137	3.9	Bq/Kg raw	± 2.5	Bq/Kg raw	3.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Onion	Yotsukura, Iwaki	Jun-12	Cs137	5.3	Bq/Kg raw	± 2.8	Bq/Kg raw	5.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Onion	Yotsukura, Iwaki	Jun-12	Cs137	3.4	Bq/Kg raw	± 2.4	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Yotsukura, Iwaki	Jun-12	Cs137	3.4	Bq/Kg raw	± 2.4	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Shimotakaku, Taira, Iwaki	Jun-12	Cs137	2.8	Bq/Kg raw	± 2.1	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Kitakabeya, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Akai, Taira, Iwaki	Jun-12	Cs137	2.2	Bq/Kg raw	± 2.2	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Shimohirakubo, Taira, Iwaki	Jun-12	Cs137	2.9	Bq/Kg raw	± 2.3	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Shimokabeya, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Numanouchi, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.0		Cs134	6.0 Bq/Kg raw
Onion	Numanouchi, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	1.7	Cs137	7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.5		Cs134	6.0 Bq/Kg raw
Onion	Ogoe, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Shimoogoe, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Uchigo, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.3	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
Onion	Joban, Iwaki	Jun-12	Cs137	2.4	Bq/Kg raw	± 2.1	2.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Shimokuramochi, Kashima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Nagasaki, Iwaki	Jun-12	Cs137	2.7	Bq/Kg raw	± 2.5	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Onahama, Iwaki	Jun-12	Cs137	3.8	Bq/Kg raw	± 2.8	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Onahama, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 4.4	5.0	Cs137	8.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 3.6		Cs134	8.0 Bq/Kg raw
Onion	Onahama, Iwaki	Jun-12	Cs137	4.9	Bq/Kg raw	± 2.9	4.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Tamagawa, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 1.7		Cs134	6.0 Bq/Kg raw
Onion	Okaona, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Okaona, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.5	Cs137	7.0 Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.6		Cs134	6.0 Bq/Kg raw
Onion	Kamikajiro, Onahama, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Onion	Tono, Iwaki	Jun-12	Cs137	2.9	Bq/Kg raw	± 2.0	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Kawabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Shimokawa, Izumi, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Baba, Nishiki, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Watanabe, Iwaki	Jun-12	Cs137	6.2	Bq/Kg raw	± 3.1	6.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Shimoyoshima, Yoshima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Sumiyoshi, Onahama, Iwaki	Jun-12	Cs137	4.2	Bq/Kg raw	± 2.1	4.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Sumiyoshi, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Ena, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Carrot	Numanouchi, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Carrot	Shimokuramochi, Kashima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Carrot	Kawabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese shallot	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	1.2	Cs137	8.0 Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 5.0		Cs134	8.0 Bq/Kg raw
Garlic	Tamayama, Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Garlic	Kitakabeya, Taira, Iwaki	Jun-12	Cs137	0.2	Bq/Kg raw	± 3.7	3.1	Cs137	8.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 3.2		Cs134	8.0 Bq/Kg raw
Garlic	Akai, Taira, Iwaki	Jun-12	Cs137	4.2	Bq/Kg raw	± 2.9	4.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Garlic	Shimohirakubo, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Garlic	Shimoogoe, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Garlic	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Garlic	Izumi, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Garlic	Watanabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cabbage	Yotsukura, Iwaki	Jun-12	Cs137	4.7	Bq/Kg raw	± 4.6	4.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Cabbage	Kattsuo, Toyoma, Iwaki	Jun-12	Cs137	3.7	Bq/Kg raw	± 3.4	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cabbage	Shimotakaku, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Cabbage	Kashima, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cabbage	Kamikajiro, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Lettuce	Unknown	Unknown	Cs137	1.6	Bq/Kg raw	± 0.1	3.5	Cs137	8.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 3.7		Cs134	8.0 Bq/Kg raw
Lettuce	Okaona, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Chinese chive	Yoshima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	5.0	Cs137	7.0 Bq/Kg raw
			Cs134	5.0	Bq/Kg raw	± 4.0		Cs134	6.0 Bq/Kg raw
Chinese chive	Tamagawa, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Chinese chive	Kimigatsuka, Onahama, Iwaki	Jun-12	Cs137	5.5	Bq/Kg raw	± 4.8	10.8	Cs137	7.0 Bq/Kg raw
			Cs134	5.3	Bq/Kg raw	± 3.9		Cs134	6.0 Bq/Kg raw
Chinese chive	Shimokawa, Izumi, Iwaki	Jun-12	Cs137	6.3	Bq/Kg raw	± 3.7	6.3	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Yoshima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Cucumber	Jiyugaoka, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 1.7		Cs134	6.0 Bq/Kg raw
Cucumber	Shimohirakubo, Taira, Iwaki	Jun-12	Cs137	2.1	Bq/Kg raw	± 2.0	2.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Suginami, Taira, Iwaki	Jun-12	Cs137	4.5	Bq/Kg raw	± 3.4	4.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Shimoogoe, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Akai, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Onahama, Iwaki	Jun-12	Cs137	2.6	Bq/Kg raw	± 2.2	2.6	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Kamikajiro, Onahama, Iwaki	Jun-12	Cs137	2.1	Bq/Kg raw	± 1.6	2.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Onahama, Iwaki	Jun-12	Cs137	3.4	Bq/Kg raw	± 2.5	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Kashima, Iwaki	Jun-12	Cs137	2.5	Bq/Kg raw	± 2.1	2.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Izumi, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Watanabe, Iwaki	Jun-12	Cs137	4.4	Bq/Kg raw	± 3.0	4.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Cucumber	Watanabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Cucumber	Tanabe, Watanabe, Iwaki	Jun-12	Cs137	4.2	Bq/Kg raw	± 2.5	Bq/Kg raw	4.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Yotsukura, Iwaki	Jun-12	Cs137	6.8	Bq/Kg raw	± 3.0	Bq/Kg raw	6.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Shimotakaku, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Ena, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137 8.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 5.1	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Leek	Kawamae, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Ice plant	Shimokabeya, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.0	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Podded pea	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Snap garden pea	Shimoogoe, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Snap garden pea	Kamiyamaguchi, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broad bean	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Azuki bean	Kawahara, Kawabe, Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Azuki bean	Kamata, Fukushima	Sep-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbage	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Shungiku	Nishiki, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 2.8	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Zucchini	Kitashirado, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Celery	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Stem lettuce	Kamiyamaguchi, Taira, Iwaki	Jun-12	Cs137	7.4	Bq/Kg raw	± 4.4	Bq/Kg raw	7.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Watercress	Kawamae, Iwaki	Jun-12	Cs137	16.2	Bq/Kg raw	± 5.7	Bq/Kg raw	22.1	Cs137 7.0 Bq/Kg raw
			Cs134	5.9	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Pumpkin	Suginami, Taira, Iwaki	Jun-12	Cs137	7.3	Bq/Kg raw	± 3.4	Bq/Kg raw	12.3	Cs137 7.0 Bq/Kg raw
			Cs134	5.0	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Pumpkin	Nishiki, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Turnip	Ogawa, Iwaki	Jun-12	Cs137	4.9	Bq/Kg raw	± 2.5	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sesame leaf	Kawabe, Iwaki	Jun-12	Cs137	2.8	Bq/Kg raw	± 1.8	Bq/Kg raw	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Japanese basil	Akai, Taira, Iwaki	Jun-12	Cs137	5.7	Bq/Kg raw	± 4.9	5.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese basil	Shimoyunagaya, Joban, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.5	Cs137	8.0 Bq/Kg raw
			Cs134	0.5	Bq/Kg raw	± 11.9		Cs134	8.0 Bq/Kg raw
Japanese basil	Kawabe, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 2.1	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese ginger	Kawamae, Iwaki	Jun-12	Cs137	7.0	Bq/Kg raw	± 3.6	7.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese horseradish	Kawamae, Iwaki	Jun-12	Cs137	77.5	Bq/Kg raw	± 17.1	134	Cs137	7.0 Bq/Kg raw
			Cs134	56.3	Bq/Kg raw	± 12.2		Cs134	6.0 Bq/Kg raw
Butterbur	Kawamae, Iwaki	Unknown	Cs137	6.5	Bq/Kg raw	± 3.8	13.1	Cs137	7.0 Bq/Kg raw
			Cs134	6.6	Bq/Kg raw	± 3.3		Cs134	6.0 Bq/Kg raw
Butterbur	Kawamae, Iwaki	Jun-12	Cs137	6.4	Bq/Kg raw	± 4.2	12.2	Cs137	7.0 Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 3.6		Cs134	6.0 Bq/Kg raw
Butterbur	Onahama, Iwaki	Jun-12	Cs137	5.1	Bq/Kg raw	± 3.3	5.4	Cs137	8.0 Bq/Kg raw
			Cs134	0.3	Bq/Kg raw	± 5.6		Cs134	8.0 Bq/Kg raw
Butterbur	Numabe, Iwaki	Jun-12	Cs137	3.6	Bq/Kg raw	± 3.4	7.8	Cs137	7.0 Bq/Kg raw
			Cs134	4.2	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Butterbur (with skin)	Tanakura, Higashishirakawa	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Kamiyamaguchi, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	7.7	Cs137	7.0 Bq/Kg raw
			Cs134	7.7	Bq/Kg raw	± 4.1		Cs134	6.0 Bq/Kg raw
Butterbur (boiled)	Numabe, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Urui	Kawamae, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	7.4	Cs137	7.0 Bq/Kg raw
			Cs134	7.4	Bq/Kg raw	± 4.2		Cs134	6.0 Bq/Kg raw
Shidoki	Kawamae, Iwaki	Jun-12	Cs137	96.5	Bq/Kg raw	± 21.7	162	Cs137	7.0 Bq/Kg raw
			Cs134	65.8	Bq/Kg raw	± 14.7		Cs134	6.0 Bq/Kg raw
Bracken	Furudono, Ishikawa	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	6.2	Cs137	7.0 Bq/Kg raw
			Cs134	6.2	Bq/Kg raw	± 3.5		Cs134	6.0 Bq/Kg raw
Bracken	Kawamae, Iwaki	Unknown	Cs137	94.5	Bq/Kg raw	± 20.4	163	Cs137	7.0 Bq/Kg raw
			Cs134	68.8	Bq/Kg raw	± 14.5		Cs134	6.0 Bq/Kg raw
Udo	Kawamae, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	7.0	Cs137	7.0 Bq/Kg raw
			Cs134	7.0	Bq/Kg raw	± 3.5		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Kawamae, Iwaki	Jun-12	Cs137	256.0	Bq/Kg raw	± 51.0	427	Cs137	7.0 Bq/Kg raw
			Cs134	171.0	Bq/Kg raw	± 34.0		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Arakawa, Taira, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.2		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Shimoyamaguchi, Taira, Iwaki	Jun-12	Cs137	20.6	Bq/Kg raw	± 5.3	39.2	Cs137	7.0 Bq/Kg raw
			Cs134	18.6	Bq/Kg raw	± 4.4		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Shimokajiro, Onahama, Iwaki	Jun-12	Cs137	118	Bq/Kg raw	± 24.0	201	Cs137	7.0 Bq/Kg raw
			Cs134	82.6	Bq/Kg raw	± 16.5		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Ueda, Iwaki	May-12	Cs137	5.3	Bq/Kg raw	± 4.1	17.0	Cs137	7.0 Bq/Kg raw
			Cs134	11.7	Bq/Kg raw	± 4.0		Cs134	6.0 Bq/Kg raw
Bamboo shoot	Joba, Iwaki	Jun-12	Cs137	47.6	Bq/Kg raw	± 10.8	89.6	Cs137	7.0 Bq/Kg raw
			Cs134	42.0	Bq/Kg raw	± 8.9		Cs134	6.0 Bq/Kg raw

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Bamboo shoot (boiled)	Kamiyamaguchi, Taira, Iwaki	Jun-12	Cs137	18.1	Bq/Kg raw	± 5.1	30.8	Cs137	7.0 Bq/Kg raw
			Cs134	12.7	Bq/Kg raw	± 3.7		Cs134	6.0 Bq/Kg raw
Bamboo shoot (boiled)	Shimokajiro, Onahama, Iwaki	Jun-12	Cs137	27.5	Bq/Kg raw	± 7.0	50.1	Cs137	7.0 Bq/Kg raw
			Cs134	22.6	Bq/Kg raw	± 5.4		Cs134	6.0 Bq/Kg raw
Unripe plum	Miwa, Iwaki	Jun-12	Cs137	4.3	Bq/Kg raw	± 2.3	8.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
Unripe plum	Yoshima, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 2.0	6.3	Cs137	7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
Unripe plum	Yotsukura, Iwaki	Jun-12	Cs137	15.8	Bq/Kg raw	± 4.6	26.4	Cs137	7.0 Bq/Kg raw
			Cs134	10.6	Bq/Kg raw	± 3.3		Cs134	6.0 Bq/Kg raw
Unripe plum	Taira, Iwaki	Jun-12	Cs137	20.9	Bq/Kg raw	± 5.6	34.8	Cs137	7.0 Bq/Kg raw
			Cs134	13.9	Bq/Kg raw	± 3.8		Cs134	6.0 Bq/Kg raw
Unripe plum	Hashirikuma, Kashima, Iwaki	Jun-12	Cs137	13.6	Bq/Kg raw	± 4.3	26.0	Cs137	7.0 Bq/Kg raw
			Cs134	12.4	Bq/Kg raw	± 3.6		Cs134	6.0 Bq/Kg raw
Unripe plum	Okaona, Onahama, Iwaki	Jun-12	Cs137	7.8	Bq/Kg raw	± 3.7	15.0	Cs137	7.0 Bq/Kg raw
			Cs134	7.2	Bq/Kg raw	± 3.0		Cs134	6.0 Bq/Kg raw
Unripe plum	Okaona, Onahama, Iwaki	Jun-12	Cs137	5.9	Bq/Kg raw	± 2.9	10.7	Cs137	7.0 Bq/Kg raw
			Cs134	4.8	Bq/Kg raw	± 2.3		Cs134	6.0 Bq/Kg raw
Unripe plum	Shimoyunagaya, Joban, Iwaki	Jun-12	Cs137	14.4	Bq/Kg raw	± 4.0	23.1	Cs137	7.0 Bq/Kg raw
			Cs134	8.7	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Unripe plum	Shimokawa, Izumi, Iwaki	Jun-12	Cs137	9.3	Bq/Kg raw	± 3.2	19.3	Cs137	7.0 Bq/Kg raw
			Cs134	10.0	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Unripe plum	Izumi, Iwaki	Jun-12	Cs137	16.9	Bq/Kg raw	± 4.7	26.4	Cs137	7.0 Bq/Kg raw
			Cs134	9.5	Bq/Kg raw	± 3.1		Cs134	6.0 Bq/Kg raw
Unripe plum	Sawagami, Kawabe, Iwaki	Jun-12	Cs137	24.5	Bq/Kg raw	± 5.9	45.7	Cs137	7.0 Bq/Kg raw
			Cs134	21.2	Bq/Kg raw	± 4.9		Cs134	6.0 Bq/Kg raw
Unripe plum	Kawabe, Iwaki	Jun-12	Cs137	27.1	Bq/Kg raw	± 6.6	46.5	Cs137	7.0 Bq/Kg raw
			Cs134	19.4	Bq/Kg raw	± 4.7		Cs134	6.0 Bq/Kg raw
Unripe plum	Kawabe, Iwaki	Jun-06	Cs137	4.7	Bq/Kg raw	± 2.6	8.7	Cs137	7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.3		Cs134	6.0 Bq/Kg raw
Unripe plum	Iwaki	Jun-12	Cs137	4.3	Bq/Kg raw	± 3.2	8.2	Cs137	7.0 Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Unripe plum (with seed)	Yoshima, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 2.2	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Unripe plum (with seed)	Ena, Iwaki	Jun-12	Cs137	25.5	Bq/Kg raw	± 6.8	43.2	Cs137	7.0 Bq/Kg raw
			Cs134	17.7	Bq/Kg raw	± 4.7		Cs134	6.0 Bq/Kg raw
Unripe plum (with seed)	Onahama, Iwaki	Jun-12	Cs137	3.2	Bq/Kg raw	± 2.9	7.5	Cs137	7.0 Bq/Kg raw
			Cs134	4.3	Bq/Kg raw	± 2.5		Cs134	6.0 Bq/Kg raw
Plum (pickled in salt)	Kawabe, Iwaki	Jun-12	Cs137	4.9	Bq/Kg raw	± 2.3	7.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
Pickled plum	Sawagami, Kawabe, Iwaki	Jun-12	Cs137	12.3	Bq/Kg raw	± 3.8	19.1	Cs137	7.0 Bq/Kg raw
			Cs134	6.8	Bq/Kg raw	± 2.7		Cs134	6.0 Bq/Kg raw
Pickled plum	Kawahara, Kawabe, Iwaki	Jun-11	Cs137	41.0	Bq/Kg raw	± 9.1	69.6	Cs137	7.0 Bq/Kg raw
			Cs134	28.6	Bq/Kg raw	± 6.2		Cs134	6.0 Bq/Kg raw
Pickled plum (without seed)	Hisanoohama, Iwaki	Jun-11	Cs137	167	Bq/Kg raw	± 33.0	299	Cs137	7.0 Bq/Kg raw
			Cs134	132	Bq/Kg raw	± 26.0		Cs134	6.0 Bq/Kg raw

## ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Shiitake mushroom	Shimokajiro, Onahama, Iwaki	Unknown	Cs137	130.0	Bq/Kg raw	± 20.9	218	Cs137	8.0 Bq/Kg raw
			Cs134	87.7	Bq/Kg raw	± 13.1		Cs134	8.0 Bq/Kg raw
Maitake mushroom	Fukushima	Jun-12	Cs137	19.9	Bq/Kg raw	± 14.6	32.4	Cs137	8.0 Bq/Kg raw
			Cs134	12.5	Bq/Kg raw	± 7.2		Cs134	8.0 Bq/Kg raw
Dried Shiitake mushroom	Shimokabeya, Taira, Iwaki	Mar-12	Cs137	1640	Bq/Kg raw	± 330.0	2,850	Cs137	7.0 Bq/Kg raw
			Cs134	1210	Bq/Kg raw	± 240.0		Cs134	6.0 Bq/Kg raw
Dried Shiitake mushroom	Kashima, Iwaki	Apr-12	Cs137	973.0	Bq/Kg raw	± 195.0	1,690	Cs137	7.0 Bq/Kg raw
			Cs134	717.0	Bq/Kg raw	± 143.0		Cs134	6.0 Bq/Kg raw
Tea leaf	Shizuoka	Apr-12	Cs137	137.0	Bq/Kg raw	± 30.0	246	Cs137	7.0 Bq/Kg raw
			Cs134	109.0	Bq/Kg raw	± 23.0		Cs134	6.0 Bq/Kg raw
Tea leaf	Ibaraki	Jun-12	Cs137	146.0	Bq/Kg raw	± 31.0	248	Cs137	7.0 Bq/Kg raw
			Cs134	102.0	Bq/Kg raw	± 21.0		Cs134	6.0 Bq/Kg raw
Fertile egg	Kashima, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.3		Cs134	6.0 Bq/Kg raw
Japanese mitten crab	Natsuigawa, Iwaki	Jun-12	Cs137	21.0	Bq/Kg raw	± 5.3	37.5	Cs137	7.0 Bq/Kg raw
			Cs134	16.5	Bq/Kg raw	± 4.0		Cs134	6.0 Bq/Kg raw
Shellfish	Shimokajiro, Onahama, Iwaki	Jun-12	Cs137	7.0	Bq/Kg raw	± 3.4	13.0	Cs137	7.0 Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Greenling	Pier 7, Onahama, Iwaki	Jun-12	Cs137	77.0	Bq/Kg raw	± 16.1	129	Cs137	7.0 Bq/Kg raw
			Cs134	51.7	Bq/Kg raw	± 10.7		Cs134	6.0 Bq/Kg raw
Japanese plum	Iwaki	Jun-12	Cs137	4.2	Bq/Kg raw	± 3.3	8.0	Cs137	7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 3.1		Cs134	6.0 Bq/Kg raw
Loquat	Shimoyunagaya, Joban, Iwaki	Jun-12	Cs137	15.0	Bq/Kg raw	± 4.2	26.6	Cs137	7.0 Bq/Kg raw
			Cs134	11.6	Bq/Kg raw	± 3.1		Cs134	6.0 Bq/Kg raw
Melon	Ibaraki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Dried persimmon	Tabito, Iwaki	Nov-11	Cs137	102	Bq/Kg raw	± 17.8	162	Cs137	8.0 Bq/Kg raw
			Cs134	59.6	Bq/Kg raw	± 10.5		Cs134	8.0 Bq/Kg raw
Miso	Naraha, Futaba	Apr-10	Cs137	1.9	Bq/Kg raw	± 1.3	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Miso	Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Miso	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Salted rice bran used for pickling	Yamagata	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
School lunch	Uchigo, Iwaki	Jun-12	Cs137	3.1	Bq/Kg raw	± 1.7	3.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
School lunch	Uchigo, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Rose jam	Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	3.9	Cs137	7.0 Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 1.8		Cs134	6.0 Bq/Kg raw
Soup	Shimokajiro, Onahama, Iwaki	Jun-12	Cs137	3.3	Bq/Kg raw	± 2.2	5.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
Sake(Daiginjo)	Takasaka, Uchigo, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Sake(Ginjo)	Takasaka, Uchigo, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sake(Junmai)	Takasaka, Uchigo, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137 6.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.2	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Sake(Genshu)	Takasaka, Uchigo, Iwaki	May-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137 6.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.2	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Sake made from apricot	Jiyugaoka, Iwaki	Unknown	Cs137	12.7	Bq/Kg raw	± 2.9	Bq/Kg raw	21.0	Cs137 5.0 Bq/Kg raw
			Cs134	8.3	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Apricot (in liquor)	Jiyugaoka, Iwaki	Aug-11	Cs137	13.1	Bq/Kg raw	± 3.8	Bq/Kg raw	24.1	Cs137 7.0 Bq/Kg raw
			Cs134	11.0	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Amazake	Taira, Iwaki	Jun-12	Cs137	1.3	Bq/Kg raw	± 0.7	Bq/Kg raw	1.3	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap water	Hanabatake, Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Nakanosaku, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
River water	Tabito, Iwaki	Jun-12	Cs137	0.9	Bq/Kg raw	± 0.9	Bq/Kg raw	0.9	Cs137 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Tea beverage	Ibaraki	Unknown	Cs137	1.1	Bq/Kg raw	± 0.7	Bq/Kg raw	2.1	Cs137 5.0 Bq/Kg raw
			Cs134	1.0	Bq/Kg raw	± 0.6	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Rain water	Kawasaki, Kanagawa	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Hot spring water	Kusakidai, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Bay leaf	Ichikawa, Chiba	Jun-12	Cs137	35.7	Bq/Kg raw	± 146.5	Bq/Kg raw	0.0	Cs137 8.0 Bq/Kg raw
			Cs134	28.7	Bq/Kg raw	± 15.8	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Field horsetail	Nakoso, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.4	Cs137 7.0 Bq/Kg raw
			Cs134	5.4	Bq/Kg raw	± 4.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White clover	Nakoso, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.4	Cs137 7.0 Bq/Kg raw
			Cs134	5.4	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Rosemary	Shimoyunagaya, Joban, Iwaki	Jun-12	Cs137	237.0	Bq/Kg raw	± 54.0	Bq/Kg raw	416	Cs137 7.0 Bq/Kg raw
			Cs134	179.0	Bq/Kg raw	± 40.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Oak leaves	Nishiki, Iwaki	Jun-12	Cs137	15.4	Bq/Kg raw	± 6.6	Bq/Kg raw	21.5	Cs137 7.0 Bq/Kg raw
			Cs134	6.1	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Rice straw	Onahama, Iwaki	Oct-10	Cs137	635	Bq/Kg raw	± 136	Bq/Kg raw	1,125	Cs137 7.0 Bq/Kg raw
			Cs134	490	Bq/Kg raw	± 101	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Compost	Naraha, Futaba	Jun-12	Cs137	8400	Bq/Kg raw	± 750	Bq/Kg raw	13,800	Cs137 8.0 Bq/Kg raw
			Cs134	5400	Bq/Kg raw	± 470	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Humus	Kubo, Kashima, Iwaki	Jun-12	Cs137	4294	Bq/Kg dry	± 403	Bq/Kg dry	7,065	Cs137 8.0 Bq/Kg dry
			Cs134	2771	Bq/Kg dry	± 252	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Sea sand	Shimotakaku, Taira, Iwaki	Oct-10	Cs137	12.0	Bq/Kg dry	± 7.0	Bq/Kg dry	16.0	Cs137 8.0 Bq/Kg dry
			Cs134	4.0	Bq/Kg dry	± 5.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Sandbox	Suginami, Taira, Iwaki	Jun-12	Cs137	95	Bq/Kg dry	± 15.0	Bq/Kg dry	151	Cs137 8.0 Bq/Kg dry
			Cs134	56	Bq/Kg dry	± 9.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Tatemachi, Nagai, Yamagata	Jun-12	Cs137	31.0	Bq/Kg dry	± 10.0	Bq/Kg dry	51.0	Cs137 8.0 Bq/Kg dry
			Cs134	20.0	Bq/Kg dry	± 6.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry

# ★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Field soil	Naraha, Futaba	Jun-12	Cs137	646	Bq/Kg dry	± 71.0	Bq/Kg dry	1,059	Cs137	8.0 Bq/Kg dry
			Cs134	413	Bq/Kg dry	± 43.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Naraha, Futaba	Jun-12	Cs137	671	Bq/Kg dry	± 73.0	Bq/Kg dry	1,082	Cs137	8.0 Bq/Kg dry
			Cs134	411	Bq/Kg dry	± 44.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Naraha, Futaba	Jun-12	Cs137	342	Bq/Kg dry	± 42.0	Bq/Kg dry	564	Cs137	8.0 Bq/Kg dry
			Cs134	222	Bq/Kg dry	± 26.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Midaisakai, Uchigo, Iwaki	Jun-12	Cs137	247	Bq/Kg dry	± 33.0	Bq/Kg dry	416	Cs137	8.0 Bq/Kg dry
			Cs134	169	Bq/Kg dry	± 21.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Kubo, Kashima, Iwaki	Jun-12	Cs137	525	Bq/Kg dry	± 61.0	Bq/Kg dry	840	Cs137	8.0 Bq/Kg dry
			Cs134	315	Bq/Kg dry	± 36.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Izumi, Iwaki	May-12	Cs137	429	Bq/Kg dry	± 50.0	Bq/Kg dry	712	Cs137	8.0 Bq/Kg dry
			Cs134	283	Bq/Kg dry	± 32.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Field soil	Suginami, Taira, Iwaki	Jun-12	Cs137	42	Bq/Kg dry	± 12.0	Bq/Kg dry	73	Cs137	8.0 Bq/Kg dry
			Cs134	31	Bq/Kg dry	± 7.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry
Rice-field soil	Tatemachi, Nagai, Yamagata	Jun-12	Cs137	36.0	Bq/Kg dry	± 12.0	Bq/Kg dry	56.0	Cs137	8.0 Bq/Kg dry
			Cs134	20.0	Bq/Kg dry	± 8.0	Bq/Kg dry		Cs134	8.0 Bq/Kg dry

