



Radiation Measurement Results of 328 items in July 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Akita	Nov-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	± 1.9 Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw
			Cs134	2.2 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Chiba	Sep-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	± 2.0 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw
			Cs134	2.1 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Aizu	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Aizu	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	± 1.8 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw
			Cs134	2.1 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Aizu	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	± 1.9 Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.7 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Furudono, Ishikawa	Oct-11	Cs137	4.2 Bq/Kg raw	± 2.3 Bq/Kg raw	± 2.0 Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.7 Bq/Kg raw	± 2.3 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Kusano, Iwaki	Sep-11	Cs137	1.9 Bq/Kg raw	± 2.3 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 2.3 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Sep-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	± 2.2 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw
			Cs134	1.6 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Unknown	Cs137	4.0 Bq/Kg raw	± 2.2 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 2.2 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Unknown	Cs137	7.8 Bq/Kg raw	± 3.5 Bq/Kg raw	± 2.6 Bq/Kg raw	12.5	Cs137	7.0 Bq/Kg raw
			Cs134	4.7 Bq/Kg raw	± 3.5 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Unknown	Cs137	3.1 Bq/Kg raw	± 2.5 Bq/Kg raw	± 2.2 Bq/Kg raw	5.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.7 Bq/Kg raw	± 2.5 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Unknown	Cs137	6.9 Bq/Kg raw	± 3.0 Bq/Kg raw	± 0.0 Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 3.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Unknown	Cs137	3.5 Bq/Kg raw	± 2.6 Bq/Kg raw	± 2.3 Bq/Kg raw	7.4	Cs137	7.0 Bq/Kg raw
			Cs134	3.9 Bq/Kg raw	± 2.6 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Unknown	Cs137	6.0 Bq/Kg raw	± 3.1 Bq/Kg raw	± 2.4 Bq/Kg raw	10.7	Cs137	7.0 Bq/Kg raw
			Cs134	4.7 Bq/Kg raw	± 3.1 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
White rice	Hokkaido	Unknown	Cs137	3.0 Bq/Kg raw	± 2.1 Bq/Kg raw	± 0.0 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 2.1 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
White rice	Akita	Nov-11	Cs137	3.1 Bq/Kg raw	± 1.9 Bq/Kg raw	± 0.0 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 1.9 Bq/Kg raw			Cs134	6.0 Bq/Kg raw
White rice	Tochigi	Oct-11	Cs137	1.8 Bq/Kg raw	± 1.0 Bq/Kg raw	± 0.8 Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.9 Bq/Kg raw	± 1.0 Bq/Kg raw			Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
White rice	Yamagata	Oct-11	Cs137	3.5 Bq/Kg raw	± 2.1 Bq/Kg raw	3.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Gifu	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Sekimoto, Kitaibaraki	Oct-11	Cs137	3.2 Bq/Kg raw	± 2.3 Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Hitachi, Ibaraki	Oct-11	Cs137	1.7 Bq/Kg raw	± 1.5 Bq/Kg raw	1.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Motomiya	Nov-11	Cs137	2.1 Bq/Kg raw	± 1.5 Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 1.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Yotsukura, Iwaki	Nov-11	Cs137	2.2 Bq/Kg raw	± 1.2 Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw	
			Cs134	1.9 Bq/Kg raw	± 1.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Kusano, Iwaki	Sep-11	Cs137	3.8 Bq/Kg raw	± 2.3 Bq/Kg raw	3.8	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Kamitakaku, Taira, Iwaki	Oct-03	Cs137	3.6 Bq/Kg raw	± 2.0 Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Onahamashima, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Yamada, Iwaki	Oct-11	Cs137	3.9 Bq/Kg raw	± 2.0 Bq/Kg raw	3.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Tabito, Iwaki	Nov-11	Cs137	3.2 Bq/Kg raw	± 2.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Nishiki, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw	
			Cs134	2.2 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Nakoso, Iwaki	Sep-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	1.9 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Nakoso, Iwaki	Unknown	Cs137	2.1 Bq/Kg raw	± 2.0 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Nakoso, Iwaki	Unknown	Cs137	2.2 Bq/Kg raw	± 1.6 Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Nakoso, Iwaki	Unknown	Cs137	2.2 Bq/Kg raw	± 2.1 Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sticky rice	Akita	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kozaki rice	Nakoso, Iwaki	Unknown	Cs137	3.0 Bq/Kg raw	± 2.9 Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw	
			Cs134	3.9 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kozaki rice	Nakoso, Iwaki	Unknown	Cs137	7.7 Bq/Kg raw	± 3.2 Bq/Kg raw	10.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.9 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Fried rice	Sekiba, Ogawa, Iwaki	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw	
			Cs134	2.9 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Fried rice	Nishiki, Iwaki	Oct-11	Cs137	3.3 Bq/Kg raw	± 2.3 Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Nagai, Yamagata	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Kitaibaraki, Ibaraki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Potato	Ohisa, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Ogawa, Iwaki	Jun-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Ogawa, Iwaki	Jul-12	Cs137	3.0 Bq/Kg raw	± 2.4 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Shibahara, Ogawa, Iwaki	Jul-12	Cs137	3.3 Bq/Kg raw	± 2.4 Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Yotsukura, Iwaki	Sep-12	Cs137	2.6 Bq/Kg raw	± 2.5 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Yotsukura, Iwaki	Jul-12	Cs137	4.7 Bq/Kg raw	± 2.4 Bq/Kg raw	6.8	Cs137	7.0 Bq/Kg raw	
			Cs134	2.1 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Kamiyoshima, Yoshima, Iwaki	Jul-12	Cs137	2.0 Bq/Kg raw	± 1.9 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Yoshima, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Nakajima, Yoshima, Iwaki	Jul-12	Cs137	3.1 Bq/Kg raw	± 2.0 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Uchigo, Iwaki	Jun-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Hirakubo, Taira, Iwaki	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	1.9 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Nakayama, Taira, Iwaki	Jun-12	Cs137	2.4 Bq/Kg raw	± 2.1 Bq/Kg raw	2.4	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Nakayama, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Shimoyamaguchi, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Fujima, Taira, Iwaki	Jul-12	Cs137	4.2 Bq/Kg raw	± 2.8 Bq/Kg raw	4.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Shimokatayose, Taira, Iwaki	Jun-12	Cs137	4.7 Bq/Kg raw	± 2.4 Bq/Kg raw	4.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Kitashirado, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Kabeya, Taira, Iwaki	Jun-12	Cs137	2.1 Bq/Kg raw	± 2.1 Bq/Kg raw	4.2	Cs137	7.0 Bq/Kg raw	
			Cs134	2.1 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Nakashio, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Shimokabeya, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Kamihirakubo, Taira, Iwaki	Jul-12	Cs137	2.1 Bq/Kg raw	± 3.2 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Potato	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Potato	Chuodai, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Kashima, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Kashima, Iwaki	Jul-12	Cs137	2.5	Bq/Kg raw	± 2.2	Bq/Kg raw	2.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Kashima, Iwaki	Jul-12	Cs137	4.2	Bq/Kg raw	± 3.0	Bq/Kg raw	4.2	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Ena, Iwaki	Jul-12	Cs137	2.3	Bq/Kg raw	± 2.6	Bq/Kg raw	2.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Onahama, Iwaki	Jul-12	Cs137	3.1	Bq/Kg raw	± 2.4	Bq/Kg raw	3.1	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137	7.0	Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Kamikajiro, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Ohara, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Sumiyoshi, Onahama, Iwaki	Jul-12	Cs137	3.0	Bq/Kg raw	± 2.1	Bq/Kg raw	3.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Onahama, Iwaki	Jul-12	Cs137	4.3	Bq/Kg raw	± 3.6	Bq/Kg raw	4.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Shimoyunagaya, Joban, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Misawa, Joban, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Nishigo, Joban, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Tamatsuyu, Izumi, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Izumi, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Miyamada, Tono, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Negishi, Tono, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Miyamada, Tono, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Iritono, Tono, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Potato	Yamada, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Namesawa, Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Soeno, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Yoshinoya, Iwaki	Jul-12	Cs137	2.6	Bq/Kg raw	± 2.3	Bq/Kg raw	2.6	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Tabito, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Kawabe, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Nakoso, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato (with skin)	Shimotakaku, Taira, Iwaki	Jul-12	Cs137	3.8	Bq/Kg raw	± 2.7	Bq/Kg raw	3.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato (with skin)	Shimoyoshima, Yoshima, Iwaki	Jul-12	Cs137	4.4	Bq/Kg raw	± 7.0	Bq/Kg raw	7.4	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 6.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato (with skin)	Kashima, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.7	Cs137	7.0	Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Ogawa, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Komagome, Yotsukura, Iwaki	Jul-12	Cs137	4.0	Bq/Kg raw	± 2.8	Bq/Kg raw	4.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Kamihirakubo, Taira, Iwaki	Jul-12	Cs137	2.8	Bq/Kg raw	± 5.0	Bq/Kg raw	3.8	Cs137	8.0	Bq/Kg raw
			Cs134	1.0	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Pumpkin	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Kitashirado, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Shimotakaku, Taira, Iwaki	Jul-12	Cs137	3.4	Bq/Kg raw	± 3.4	Bq/Kg raw	3.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Shimotakaku, Taira, Iwaki	Jul-12	Cs137	3.9	Bq/Kg raw	± 3.2	Bq/Kg raw	3.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Numanouchi, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Izumi, Iwaki	Jul-12	Cs137	4.3	Bq/Kg raw	± 3.2	Bq/Kg raw	4.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pumpkin	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.2	Cs137	7.0	Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 3.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Nishiogawa, Ogawa, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Japanese radish	Akai, Taira, Iwaki	Jul-12	Cs137	2.7 Bq/Kg raw	± 2.7 Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Japanese radish	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Japanese radish	Fujima, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Japanese radish	Tono, Iwaki	Jul-12	Cs137	4.7 Bq/Kg raw	± 2.5 Bq/Kg raw	4.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Komagome, Yotsukura, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Kabeya, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Ena, Iwaki	Unknown	Cs137	3.5 Bq/Kg raw	± 3.2 Bq/Kg raw	3.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Okaona, Onahama, Iwaki	Jul-12	Cs137	4.4 Bq/Kg raw	± 2.6 Bq/Kg raw	4.4	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Tanabe, Watanabe, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Yotsukura, Iwaki	Jul-12	Cs137	2.8 Bq/Kg raw	± 2.5 Bq/Kg raw	2.8	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Nakashio, Taira, Iwaki	Jun-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Numanouchi, Taira, Iwaki	Jul-12	Cs137	2.9 Bq/Kg raw	± 1.9 Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Shimokatayose, Taira, Iwaki	Jun-12	Cs137	2.7 Bq/Kg raw	± 1.8 Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Kamihirakubo, Taira, Iwaki	Jul-12	Cs137	4.8 Bq/Kg raw	± 4.5 Bq/Kg raw	6.7	Cs137	8.0 Bq/Kg raw	
			Cs134	1.9 Bq/Kg raw	± 3.6 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Onion	Chuodai, Iwaki	Jun-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.4	Cs137	5.0 Bq/Kg raw	
			Cs134	1.4 Bq/Kg raw	± 0.8 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Onion	Kashima, Iwaki	Jun-12	Cs137	2.9 Bq/Kg raw	± 2.7 Bq/Kg raw	5.4	Cs137	7.0 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Sumiyoshi, Onahama, Iwaki	Jun-12	Cs137	3.7 Bq/Kg raw	± 2.5 Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Onion	Yamada, Iwaki	Jun-12	Cs137	1.9 Bq/Kg raw	± 1.8 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Yamada, Iwaki	Jul-12	Cs137	2.7 Bq/Kg raw	± 2.7 Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Shallot	Shimoyunagaya, Joban, Iwaki	Jul-12	Cs137	3.2 Bq/Kg raw	± 2.1 Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Garlic	Yamada, Iwaki	Jul-12	Cs137	3.7 Bq/Kg raw	± 3.2 Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Cucumber	Naraha, Futaba	Jul-12	Cs137	9.9 Bq/Kg raw	± 3.7 Bq/Kg raw	14.4	Cs137	7.0 Bq/Kg raw	
			Cs134	4.5 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Ohisa, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Miwa, Iwaki	Jul-12	Cs137	2.7 Bq/Kg raw	± 2.2 Bq/Kg raw	2.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Nishiogawa, Ogawa, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Shibahara, Ogawa, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw	
			Cs134	3.1 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Ogawa, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.8	Cs137	7.0 Bq/Kg raw	
			Cs134	2.8 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Ogawa, Iwaki	Jul-12	Cs137	2.3 Bq/Kg raw	± 2.3 Bq/Kg raw	2.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Komagome, Yotsukura, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Yotsukura, Iwaki	Jul-12	Cs137	3.0 Bq/Kg raw	± 2.2 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Yoshima, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw	
			Cs134	2.1 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Shimoarakawa, Taira, Iwaki	Jul-12	Cs137	2.5 Bq/Kg raw	± 1.9 Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Fujima, Taira, Iwaki	Jul-12	Cs137	3.7 Bq/Kg raw	± 2.6 Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Yagawase, Taira, Iwaki	Jul-12	Cs137	2.0 Bq/Kg raw	± 2.3 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Kitakabeya, Taira, Iwaki	Unknown	Cs137	2.9 Bq/Kg raw	± 2.3 Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Nakayama, Taira, Iwaki	Jul-12	Cs137	5.4 Bq/Kg raw	± 2.5 Bq/Kg raw	8.2	Cs137	7.0 Bq/Kg raw	
			Cs134	2.8 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Numanouchi, Taira, Iwaki	Jul-12	Cs137	2.2 Bq/Kg raw	± 2.9 Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Kamiyunagaya, Joban, Iwaki	Jul-12	Cs137	5.8 Bq/Kg raw	± 3.0 Bq/Kg raw	5.8	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Shimoyunagaya, Joban, Iwaki	Jul-12	Cs137	2.6 Bq/Kg raw	± 2.5 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Sakuragaoka, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Yoshinoya, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Kubo, Kashima, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Cucumber	Kashima, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Kashima, Iwaki	Jul-12	Cs137	3.8 Bq/Kg raw	± 2.3 Bq/Kg raw	3.8	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Kashima, Iwaki	Jul-12	Cs137	2.3 Bq/Kg raw	± 2.5 Bq/Kg raw	2.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw	
			Cs134	3.0 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Onahama, Iwaki	Jul-12	Cs137	2.6 Bq/Kg raw	± 2.4 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Onahama, Iwaki	Jul-12	Cs137	3.4 Bq/Kg raw	± 2.1 Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Tamatsuyu, Izumi, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Numabe, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Miyamada, Tono, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Yamada, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Yamada, Iwaki	Jul-12	Cs137	4.2 Bq/Kg raw	± 3.7 Bq/Kg raw	7.3	Cs137	7.0 Bq/Kg raw	
			Cs134	3.1 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cucumber	Nakoso, Iwaki	Jul-12	Cs137	2.3 Bq/Kg raw	± 1.9 Bq/Kg raw	2.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Miwa, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Ogawa, Iwaki	Jul-12	Cs137	4.6 Bq/Kg raw	± 3.7 Bq/Kg raw	4.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Yoshima, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Kitsunozuka, Yotsukura, Iwaki	Jul-12	Cs137	4.3 Bq/Kg raw	± 3.0 Bq/Kg raw	4.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Akai, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Yagawase, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw	
			Cs134	1.5 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Egg plant	Hirakubo, Taira, Iwaki	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Egg plant	Kitakabeya, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Suginami, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.5	Cs137	7.0	Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Sakuragaoka, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Shimoyunagaya, Joban, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.2	Cs137	7.0	Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Okaona, Onahama, Iwaki	Jul-12	Cs137	3.2	Bq/Kg raw	± 2.4	Bq/Kg raw	3.2	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Tono, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Iritono, Tono, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	6.9	Cs137	7.0	Bq/Kg raw
			Cs134	6.9	Bq/Kg raw	± 5.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Egg plant	Yamada, Iwaki	Jul-12	Cs137	3.0	Bq/Kg raw	± 2.4	Bq/Kg raw	3.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Cabbage	Nagai, Yamagata	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.9	Cs137	7.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Cabbage	Kitaibaraki, Ibaraki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.1	Cs137	7.0	Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Cabbage	Onahama, Iwaki	Jul-12	Cs137	3.8	Bq/Kg raw	± 2.4	Bq/Kg raw	3.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Cabbage	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Cherry tomato	Kashima, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Akai, Taira, Iwaki	Jul-12	Cs137	1.9	Bq/Kg raw	± 1.8	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Hirakubo, Taira, Iwaki	Unknown	Cs137	1.4	Bq/Kg raw	± 2.1	Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Shimoogoe, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Suginami, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Satogaoka, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tomato	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Tomato	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Tomato	Moegidai, Izumi, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Tomato	Kaneyama, Iwaki	Jul-12	Cs137	2.3 Bq/Kg raw	± 2.0 Bq/Kg raw	2.3	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Japanese mustard spinach	Shimohirakubo, Taira, Iwaki	Jul-12	Cs137	3.1 Bq/Kg raw	± 2.4 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Japanese mustard spinach (salted)	Shimohirakubo, Taira, Iwaki	Jul-12	Cs137	3.7 Bq/Kg raw	± 1.9 Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Japanese mustard spinach (boiled)	Shimohirakubo, Taira, Iwaki	Jul-12	Cs137	1.9 Bq/Kg raw	± 1.8 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Malabar spinach	Yamagata	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Chinese chive	Sakuragaoka, Iwaki	Jul-12	Cs137	6.4 Bq/Kg raw	± 5.3 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
New Zealand Spinach	Yamada, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Jew's mallow	Yamagata	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Jew's mallow	Numanouchi, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw	
			Cs134	4.1 Bq/Kg raw	± 3.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Jew's mallow	Sakuragaoka, Iwaki	Jul-12	Cs137	8.9 Bq/Kg raw	± 8.0 Bq/Kg raw	8.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Corn	Numanouchi, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Corn	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Corn	Tamatsuyu, Izumi, Iwaki	Jul-12	Cs137	2.0 Bq/Kg raw	± 2.6 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Corn	Moegidai, Izumi, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Corn	Tanabe, Watanabe, Iwaki	Jul-12	Cs137	6.5 Bq/Kg raw	± 3.9 Bq/Kg raw	6.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Corn	Kaneyama, Iwaki	Jul-12	Cs137	1.5 Bq/Kg raw	± 2.4 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Romano bean	Shimotakaku, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Romano bean	Hirakubo, Taira, Iwaki	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Romano bean	Sakuragaoka, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Romano bean	Ena, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Romano bean	Namesawa, Yamada, Iwaki	Jul-12	Cs137	2.6 Bq/Kg raw	± 2.1 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Romano bean	Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Manzunashi (bean)	Watanabe, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pea	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kidney bean	Shimokajiro, Onahama, Iwaki	Jul-12	Cs137	5.3	Bq/Kg raw	± 3.6	Bq/Kg raw	5.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kidney bean	Kaneyama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kidney bean	Unknown	Unknown	Cs137	3.4	Bq/Kg raw	± 4.4	Bq/Kg raw	3.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro stem	Makunouchi, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Onahama, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Zucchini	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Bitter melon	Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Green pepper	Kitakabeya, Taira, Iwaki	Jul-12	Cs137	3.8	Bq/Kg raw	± 3.0	Bq/Kg raw	3.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Green pepper	Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Winter melon	Kubo, Kashima, Iwaki	Jul-12	Cs137	2.4	Bq/Kg raw	± 2.1	Bq/Kg raw	4.8	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Konjak	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137	7.0	Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Naraha, Futaba	Jul-12	Cs137	35.6	Bq/Kg raw	± 8.5	Bq/Kg raw	58.6	Cs137	7.0	Bq/Kg raw
			Cs134	23.0	Bq/Kg raw	± 5.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Ide, Naraha, Futaba	Unknown	Cs137	55.7	Bq/Kg raw	± 12.0	Bq/Kg raw	95.5	Cs137	7.0	Bq/Kg raw
			Cs134	39.8	Bq/Kg raw	± 8.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Ide, Naraha, Futaba	Unknown	Cs137	190	Bq/Kg raw	± 38.0	Bq/Kg raw	323	Cs137	7.0	Bq/Kg raw
			Cs134	133	Bq/Kg raw	± 27.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Yotsukura, Iwaki	Jul-12	Cs137	13.8	Bq/Kg raw	± 3.7	Bq/Kg raw	19.8	Cs137	7.0	Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Nakashio, Taira, Iwaki	Jul-12	Cs137	4.5	Bq/Kg raw	± 2.3	Bq/Kg raw	7.2	Cs137	7.0	Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Miya, Uchigo, Iwaki	Jun-12	Cs137	5.0	Bq/Kg raw	± 2.5	Bq/Kg raw	10.8	Cs137	7.0	Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Onahama, Iwaki	Jul-12	Cs137	3.6	Bq/Kg raw	± 2.7	Bq/Kg raw	7.2	Cs137	7.0	Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Ena, Iwaki	Jul-12	Cs137	21.8	Bq/Kg raw	± 5.4	Bq/Kg raw	39.7	Cs137	7.0	Bq/Kg raw
			Cs134	17.9	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Unripe plum	Kaneyama, Iwaki	Jun-12	Cs137	3.6	Bq/Kg raw	± 2.8	Bq/Kg raw	6.5	Cs137	7.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Unripe plum	Kawabe, Iwaki	Jul-12	Cs137	4.3 Bq/Kg raw	± 2.2 Bq/Kg raw	7.2	Cs137	7.0 Bq/Kg raw	
			Cs134	2.9 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Unripe plum (with seeds)	Shimokabeya, Taira, Iwaki	Jul-12	Cs137	6.7 Bq/Kg raw	± 2.8 Bq/Kg raw	9.2	Cs137	7.0 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Unripe plum (with seeds)	Izumi, Iwaki	Jul-12	Cs137	5.8 Bq/Kg raw	± 3.1 Bq/Kg raw	9.5	Cs137	7.0 Bq/Kg raw	
			Cs134	3.7 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Plum jam (Made in 2011)	Nakahirakubo, Taira, Iwaki	Jul-11	Cs137	4.5 Bq/Kg raw	± 2.3 Bq/Kg raw	6.8	Cs137	7.0 Bq/Kg raw	
			Cs134	2.3 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Plum jam (Made in 2012)	Nakahirakubo, Taira, Iwaki	Jul-12	Cs137	11.6 Bq/Kg raw	± 3.7 Bq/Kg raw	20.0	Cs137	7.0 Bq/Kg raw	
			Cs134	8.4 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Plum	Shibahara, Ogawa, Iwaki	Jun-12	Cs137	3.6 Bq/Kg raw	± 2.2 Bq/Kg raw	6.9	Cs137	7.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Plum seed	Shibahara, Ogawa, Iwaki	Jun-12	Cs137	6.7 Bq/Kg raw	± 4.3 Bq/Kg raw	6.9	Cs137	8.0 Bq/Kg raw	
			Cs134	0.2 Bq/Kg raw	± 3.3 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Dried fern	Niigata	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Bracken	Kawamae, Iwaki	Jul-12	Cs137	6.0 Bq/Kg raw	± 3.4 Bq/Kg raw	10.4	Cs137	7.0 Bq/Kg raw	
			Cs134	4.4 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Loquat	Numabe, Iwaki	Jul-12	Cs137	14.2 Bq/Kg raw	± 7.9 Bq/Kg raw	22.4	Cs137	8.0 Bq/Kg raw	
			Cs134	8.2 Bq/Kg raw	± 4.6 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Butterbur	Naraha, Futaba	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw	
			Cs134	1.8 Bq/Kg raw	± 1.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Butterbur	Naraha, Futaba	Jul-12	Cs137	31.8 Bq/Kg raw	± 8.2 Bq/Kg raw	49.9	Cs137	7.0 Bq/Kg raw	
			Cs134	18.1 Bq/Kg raw	± 5.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Bamboo shoot (Raw)	Naraha, Futaba	Jul-12	Cs137	47.7 Bq/Kg raw	± 11.0 Bq/Kg raw	80.9	Cs137	7.0 Bq/Kg raw	
			Cs134	33.2 Bq/Kg raw	± 7.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Bamboo shoot (boiled)	Naraha, Futaba	Jul-12	Cs137	7.7 Bq/Kg raw	± 2.7 Bq/Kg raw	10.3	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Soldum plum	Yamada, Iwaki	Jul-12	Cs137	4.6 Bq/Kg raw	± 2.3 Bq/Kg raw	4.6	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Soldum plum	Numabe, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.5	Cs137	7.0 Bq/Kg raw	
			Cs134	3.5 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Plum	Numabe, Iwaki	Jul-12	Cs137	5.2 Bq/Kg raw	± 2.4 Bq/Kg raw	8.7	Cs137	7.0 Bq/Kg raw	
			Cs134	3.5 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Amanatsu orange (with skin)	Ito, Shizuoka	May-12	Cs137	9.2 Bq/Kg raw	± 3.1 Bq/Kg raw	13.0	Cs137	7.0 Bq/Kg raw	
			Cs134	3.8 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blackberry	Onahama, Iwaki	Jul-12	Cs137	1.6 Bq/Kg raw	± 2.3 Bq/Kg raw	2.2	Cs137	8.0 Bq/Kg raw	
			Cs134	0.6 Bq/Kg raw	± 3.3 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Blackberry	Onahama, Iwaki	Jul-12	Cs137	2.9 Bq/Kg raw	± 2.1 Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blackberry	Yamada, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry	Yotsukura, Iwaki	Jul-12	Cs137	2.3 Bq/Kg raw	± 2.6 Bq/Kg raw	3.1	Cs137	8.0 Bq/Kg raw	
			Cs134	0.8 Bq/Kg raw	± 3.9 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Blueberry	Kitsunozuka, Yotsukura, Iwaki	Jul-12	Cs137	10.0 Bq/Kg raw	± 3.0 Bq/Kg raw	14.0	Cs137	7.0 Bq/Kg raw	
			Cs134	4.0 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Blueberry	Nakashio, Taira, Iwaki	Jul-12	Cs137	7.1 Bq/Kg raw	± 3.4 Bq/Kg raw	11.5	Cs137	7.0 Bq/Kg raw	
			Cs134	4.4 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry	Yumoto, Joban, Iwaki	Jul-12	Cs137	1.7 Bq/Kg raw	± 1.4 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw	
			Cs134	1.3 Bq/Kg raw	± 1.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry	Onahama, Iwaki	Jul-12	Cs137	5.2 Bq/Kg raw	± 3.0 Bq/Kg raw	8.2	Cs137	7.0 Bq/Kg raw	
			Cs134	3.0 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry	Watanabe, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry	Yamada, Iwaki	Jul-12	Cs137	4.1 Bq/Kg raw	± 3.6 Bq/Kg raw	6.2	Cs137	7.0 Bq/Kg raw	
			Cs134	2.1 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry	Soeno, Iwaki	Jul-12	Cs137	4.7 Bq/Kg raw	± 2.5 Bq/Kg raw	7.4	Cs137	7.0 Bq/Kg raw	
			Cs134	2.7 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Blueberry jam	Soeno, Iwaki	Jul-12	Cs137	1.6 Bq/Kg raw	± 1.1 Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw	
			Cs134	1.7 Bq/Kg raw	± 1.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Loquat (with skin)	Shimoyamaguchi, Taira, Iwaki	Jul-12	Cs137	31.1 Bq/Kg raw	± 11.4 Bq/Kg raw	50.3	Cs137	8.0 Bq/Kg raw	
			Cs134	19.2 Bq/Kg raw	± 6.2 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Loquat (without skin)	Shimoyamaguchi, Taira, Iwaki	Jul-12	Cs137	27.3 Bq/Kg raw	± 9.8 Bq/Kg raw	44.2	Cs137	8.0 Bq/Kg raw	
			Cs134	16.9 Bq/Kg raw	± 5.9 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Water melon	Kashima, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Water melon	Shimokabeya, Taira, Iwaki	Jul-12	Cs137	3.4 Bq/Kg raw	± 2.0 Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Water melon (without seeds)	Numanouchi, Taira, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Fig	Unknown	Unknown	Cs137	2.7 Bq/Kg raw	± 2.5 Bq/Kg raw	3.8	Cs137	8.0 Bq/Kg raw	
			Cs134	1.1 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Tea leaf	Shizuoka	May-12	Cs137	10.8 Bq/Kg raw	± 8.0 Bq/Kg raw	10.8	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Oak leaf	Akai, Taira, Iwaki	Jul-12	Cs137	21.9 Bq/Kg raw	± 7.6 Bq/Kg raw	32.0	Cs137	7.0 Bq/Kg raw	
			Cs134	10.1 Bq/Kg raw	± 5.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Dokudami	Taira, Iwaki	Jul-12	Cs137	8.4 Bq/Kg raw	± 4.1 Bq/Kg raw	16.8	Cs137	7.0 Bq/Kg raw	
			Cs134	8.4 Bq/Kg raw	± 3.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Mulberry	Ishimori, Iwaki	Jul-12	Cs137	4.2 Bq/Kg raw	± 3.5 Bq/Kg raw	8.3	Cs137	7.0 Bq/Kg raw	
			Cs134	4.1 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Dokudami	Onahama, Iwaki	Jul-12	Cs137	17.0 Bq/Kg raw	± 11.9 Bq/Kg raw	17.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Wild boar meat	Watanabe, Iwaki	Jul-12	Cs137	91.5 Bq/Kg raw	± 19.3 Bq/Kg raw	176	Cs137	7.0 Bq/Kg raw	
			Cs134	84.1 Bq/Kg raw	± 16.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sea bass	Onahama, Iwaki	Jul-12	Cs137	17.7 Bq/Kg raw	± 4.7 Bq/Kg raw	34.1	Cs137	7.0 Bq/Kg raw	
			Cs134	16.4 Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sea bass	Onahama, Iwaki	Jul-12	Cs137	17.2 Bq/Kg raw	± 4.5 Bq/Kg raw	25.7	Cs137	7.0 Bq/Kg raw	
			Cs134	8.5 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Mackerel	Onahama Pier 3, Iwaki	Jul-12	Cs137	3.8 Bq/Kg raw	± 2.5 Bq/Kg raw	6.1	Cs137	7.0 Bq/Kg raw	
			Cs134	2.3 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sweetfish	Asami River, Hirono, Futaba	Jul-12	Cs137	50.2 Bq/Kg raw	± 12.9 Bq/Kg raw	78.5	Cs137	8.0 Bq/Kg raw	
			Cs134	28.3 Bq/Kg raw	± 7.2 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Sweetfish	Hirono, Iwaki	Jul-12	Cs137	29.4 Bq/Kg raw	± 7.2 Bq/Kg raw	58.5	Cs137	7.0 Bq/Kg raw	
			Cs134	29.1 Bq/Kg raw	± 6.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sweetfish	Yoshima, Iwaki	Jul-12	Cs137	139.0 Bq/Kg raw	± 28.0 Bq/Kg raw	234	Cs137	7.0 Bq/Kg raw	
			Cs134	95.2 Bq/Kg raw	± 19.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Honey	Yotsukura, Iwaki	Jul-12	Cs137	16.3 Bq/Kg raw	± 4.2 Bq/Kg raw	25.5	Cs137	7.0 Bq/Kg raw	
			Cs134	9.2 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Miso	Kamiyoshima, Yoshima, Iwaki	May-12	Cs137	2.9 Bq/Kg raw	± 2.0 Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Baby formula	Unknown	Unknown	Cs137	2.9 Bq/Kg raw	± 2.8 Bq/Kg raw	2.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
School lunch	Tsuzura, Uchigo, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
School lunch	Tsuzura, Uchigo, Iwaki	Jul-12	Cs137	2.2 Bq/Kg raw	± 1.9 Bq/Kg raw	2.2	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
School lunch	Tsuzura, Uchigo, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Nikko maple	Onahama, Iwaki	Jun-12	Cs137	22.4 Bq/Kg raw	± 13.0 Bq/Kg raw	33.3	Cs137	8.0 Bq/Kg raw	
			Cs134	10.9 Bq/Kg raw	± 13.5 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Sea water (Condensed to 2 times)	Onahama port, Iwaki	Jun-12	Cs137	1.8 Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	5.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Sea water (Condensed to 2 times)	Onahama port, Iwaki	Jun-12	Cs137	2.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	5.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Well water	Koya, Uchigo, Iwaki	Jul-12	Cs137	1.2 Bq/Kg raw	± 1.0 Bq/Kg raw	1.2	Cs137	5.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Spring water	Okaona, Onahama, Iwaki	Jul-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	5.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Pasture grass	Miwa, Iwaki	Jul-12	Cs137	61.0 Bq/Kg raw	± 16.0 Bq/Kg raw	95.0	Cs137	7.0 Bq/Kg raw	
			Cs134	34.0 Bq/Kg raw	± 11.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Pasture grass	Miwa, Iwaki	Jul-12	Cs137	28.0 Bq/Kg raw	± 10.0 Bq/Kg raw	40.0	Cs137	7.0 Bq/Kg raw	
			Cs134	12.0 Bq/Kg raw	± 7.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Dried grass	Kusakidai, Iwaki	Jul-12	Cs137	328 Bq/Kg dry	± 77.0 Bq/Kg dry	648	Cs137	7.0 Bq/Kg dry	
			Cs134	320 Bq/Kg dry	± 69.0 Bq/Kg dry		Cs134	6.0 Bq/Kg dry	
Soil	Sakuradai, Morioka, Iwate	Jun-12	Cs137	12.0 Bq/Kg dry	± 10.0 Bq/Kg dry	20.0	Cs137	8.0 Bq/Kg dry	
			Cs134	8.0 Bq/Kg dry	± 5.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Soil	Naraha, Futaba	Jul-12	Cs137	583 Bq/Kg dry	± 63.0 Bq/Kg dry	923	Cs137	8.0 Bq/Kg dry	
			Cs134	340 Bq/Kg dry	± 37.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Soil	Yotsukura, Iwaki	Jul-12	Cs137	299 Bq/Kg dry	± 37.0 Bq/Kg dry	463	Cs137	8.0 Bq/Kg dry	
			Cs134	164 Bq/Kg dry	± 21.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Soil	Kamihirakubo, Taira, Iwaki	Jul-12	Cs137	408 Bq/Kg dry	± 49.0 Bq/Kg dry	635	Cs137	8.0 Bq/Kg dry	
			Cs134	227 Bq/Kg dry	± 28.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Soil	Fujima, Taira, Iwaki	Jul-12	Cs137	652 Bq/Kg dry	± 71.0 Bq/Kg dry	1,046	Cs137	8.0 Bq/Kg dry	
			Cs134	394 Bq/Kg dry	± 42.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Soil	Satogaoka, Iwaki	Jul-12	Cs137	945 Bq/Kg dry	± 98.0 Bq/Kg dry	1,474	Cs137	8.0 Bq/Kg dry	
			Cs134	529 Bq/Kg dry	± 55.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Soil	Sakuragaoka, Iwaki	Jul-12	Cs137	439 Bq/Kg dry	± 51.0 Bq/Kg dry	677	Cs137	8.0 Bq/Kg dry	
			Cs134	238 Bq/Kg dry	± 28.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Soil	Mizunoya, Joban, Iwaki	Jul-12	Cs137	357	Bq/Kg dry	± 42.0	Bq/Kg dry	570	Cs137	8.0	Bq/Kg dry
			Cs134	213	Bq/Kg dry	± 25.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Takizawa, Iwate	Jun-12	Cs137	30.0	Bq/Kg dry	± 13.0	Bq/Kg dry	50.0	Cs137	8.0	Bq/Kg dry
			Cs134	20.0	Bq/Kg dry	± 7.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Takizawa, Iwate	Jun-12	Cs137	102	Bq/Kg dry	± 20.0	Bq/Kg dry	169	Cs137	8.0	Bq/Kg dry
			Cs134	67.0	Bq/Kg dry	± 12.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Ohisa, Iwaki	Jul-12	Cs137	222	Bq/Kg dry	± 29.0	Bq/Kg dry	342	Cs137	8.0	Bq/Kg dry
			Cs134	120	Bq/Kg dry	± 17.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Yoshima, Iwaki	Jul-12	Cs137	218	Bq/Kg dry	± 31.0	Bq/Kg dry	351	Cs137	8.0	Bq/Kg dry
			Cs134	133	Bq/Kg dry	± 19.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Yoshima, Iwaki	Jul-12	Cs137	205	Bq/Kg dry	± 29.0	Bq/Kg dry	302	Cs137	8.0	Bq/Kg dry
			Cs134	97.0	Bq/Kg dry	± 15.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Kamihirakubo, Taira, Iwaki	Jul-12	Cs137	238	Bq/Kg dry	± 33.0	Bq/Kg dry	371	Cs137	8.0	Bq/Kg dry
			Cs134	133	Bq/Kg dry	± 19.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Field soil	Miya, Uchigo, Iwaki	Jul-12	Cs137	246	Bq/Kg dry	± 33.0	Bq/Kg dry	384	Cs137	8.0	Bq/Kg dry
			Cs134	138	Bq/Kg dry	± 19.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Schoolyard soil	Mizunoya, Joban, Iwaki	Jul-12	Cs137	177	Bq/Kg dry	± 224	Bq/Kg dry	293	Cs137	8.0	Bq/Kg dry
			Cs134	116	Bq/Kg dry	± 16.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Sandbox	Kusakidai, Iwaki	Jul-12	Cs137	137	Bq/Kg dry	± 20.0	Bq/Kg dry	201	Cs137	8.0	Bq/Kg dry
			Cs134	64.0	Bq/Kg dry	± 11.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry
Sandbox	Sakuradai, Morioka, Iwate	Jun-12	Cs137	16.0	Bq/Kg dry	± 10.0	Bq/Kg dry	26.0	Cs137	8.0	Bq/Kg dry
			Cs134	10.0	Bq/Kg dry	± 5.0	Bq/Kg dry		Cs134	8.0	Bq/Kg dry