



Radiation Measurement Results of 117 items in August 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Niigata	Unknown	Cs137	1.8	Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kawamata, Date	Oct-11	Cs137	1.8	Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kawamae, Iwaki	Oct-11	Cs137	1.9	Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Shimohirakubo, Taira, Iwaki	Oct-11	Cs137	3.3	Bq/Kg raw	5.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Onahama, Iwaki	Oct-11	Cs137	2.2	Bq/Kg raw	6.4	Cs137	7.0 Bq/Kg raw
			Cs134	4.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Niigata	Nov-11	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Nishigo, Nishishirakawa	Oct-11	Cs137	2.2	Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Kikuta, Koriyama	Oct-11	Cs137	3.1	Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	2.8	Bq/Kg raw	5.0	Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	2.3	Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Sep-11	Cs137	2.1	Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Sep-11	Cs137	0.0	Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Kashima, Iwaki	Sep-11	Cs137	3.8	Bq/Kg raw	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Onahama, Iwaki	Oct-11	Cs137	1.5	Bq/Kg raw	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Potato	Yabuki, Nishishirakawa	Jul-12	Cs137	1.9	Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Potato	Higashikaneko, Shirakawa	Aug-12	Cs137	0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Potato	Furudono, Ishikawa	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potato	Yotsukura, Iwaki	Jul-12	Cs137	11.5	Bq/Kg raw	± 3.7	16.8	Cs137	7.0 Bq/Kg raw
			Cs134	5.3	Bq/Kg raw	± 2.5		Cs134	6.0 Bq/Kg raw
Potato	Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potato	Onahama, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potato	Izumi, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Potato	Tabito, Iwaki	Jul-12	Cs137	0.1	Bq/Kg raw	± 0.9	0.1	Cs137	8.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	8.0 Bq/Kg raw
Pumpkin	Ena, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pumpkin	Satogaoka, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pumpkin	Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.1		Cs134	6.0 Bq/Kg raw
Pumpkin	Ohisa, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pumpkin	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	4.0 Bq/Kg raw
Pumpkin	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	1.5	Cs137	7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 3.1		Cs134	6.0 Bq/Kg raw
Pumpkin	Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pumpkin	Yamada, Iwaki	Unknown	Cs137	1.3	Bq/Kg raw	± 2.2	1.3	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pumpkin	Onahama, Iwaki	Aug-12	Cs137	2.1	Bq/Kg raw	± 3.7	2.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pumpkin	Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Onion	Yagawase, Taira, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Leek	Kashima, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Japanese radish	Yotsukura, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.6	Cs137	7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.8		Cs134	6.0 Bq/Kg raw
Cabbage	Kashima, Iwaki	Aug-12	Cs137	2.7	Bq/Kg raw	± 3.5	2.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Tomato	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Tomato	Satogaoka, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Tomato	Joban, Iwaki	Aug-12	Cs137	1.9	Bq/Kg raw	± 2.2	1.9	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Tomato	Onahamajima, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.3	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tomato	Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.8	Cs137 7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tomato	Iritono, Tono, Iwaki	Aug-12	Cs137	1.5	Bq/Kg raw	± 2.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tomato	Yamada, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137 7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cherry tomato	Yoshima, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cherry tomato	Ena, Iwaki	Aug-12	Cs137	1.3	Bq/Kg raw	± 2.3	Bq/Kg raw	1.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cherry tomato	Yamada, Iwaki	Aug-12	Cs137	1.6	Bq/Kg raw	± 2.4	Bq/Kg raw	1.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Yotukura, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Uchigo, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Okaona, Onahama, Iwaki	Aug-12	Cs137	2.1	Bq/Kg raw	± 2.6	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Onahamajima, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Onahama, Iwaki	Aug-12	Cs137	6.6	Bq/Kg raw	± 3.4	Bq/Kg raw	6.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cucumber	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Egg plant	Ohisa, Iwaki	Aug-12	Cs137	2.6	Bq/Kg raw	± 2.1	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Egg plant	Kashima, Iwaki	Aug-12	Cs137	3.2	Bq/Kg raw	± 4.6	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Egg plant	Okaona, Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Egg plant	Onahama, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Egg plant	Soeno, Iwaki	Aug-12	Cs137	3.0	Bq/Kg raw	± 3.8	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese ginger	Naraha, Futaba	Jul-12	Cs137	69.3	Bq/Kg raw	± 15.9	Bq/Kg raw	127	Cs137 7.0 Bq/Kg raw
			Cs134	57.9	Bq/Kg raw	± 12.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese ginger	Shimokabeya, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.7	Cs137 7.0 Bq/Kg raw
			Cs134	5.7	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese ginger	Onahama, Iwaki	Unknown	Cs137	9.0	Bq/Kg raw	± 4.5	Bq/Kg raw	9.0	Cs137 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Japanese ginger	Onahama, Iwaki	Aug-12	Cs137	57.7	Bq/Kg raw	± 13.9	Bq/Kg raw	112	Cs137 7.0 Bq/Kg raw
			Cs134	54.0	Bq/Kg raw	± 12.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Japanese ginger	Tono, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese ginger	Tabito, Iwaki	Unknown	Cs137	2.8	Bq/Kg raw	± 3.9	Bq/Kg raw	5.3	Cs137 7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Corn	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	1.3	Bq/Kg raw	± 2.1	Bq/Kg raw	1.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Corn	Shimohirakubo, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Corn	Onahama, Iwaki	Aug-12	Cs137	1.5	Bq/Kg raw	± 0.1	Bq/Kg raw	1.5	Cs137 8.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 8.0 Bq/Kg raw
Corn	Tono, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Corn	Yamada, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broth left over after boiling corn	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137 5.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Okra	Shimoogoe, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
Okra	Sakuragaoka, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bitter melon	Ogawa, Iwaki	Unknown	Cs137	1.8	Bq/Kg raw	± 3.2	Bq/Kg raw	1.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bitter melon	Joban, Iwaki	Aug-12	Cs137	1.2	Bq/Kg raw	± 2.2	Bq/Kg raw	1.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bitter melon	Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Zucchini	Joban, Iwaki	Aug-12	Cs137	2.4	Bq/Kg raw	± 2.6	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bean	Taira, Iwaki	Aug-12	Cs137	4.4	Bq/Kg raw	± 4.5	Bq/Kg raw	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bean	Shimohirakubo, Taira, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Bean	Yamada, Iwaki	Unknown	Cs137	1.7	Bq/Kg raw	± 2.8	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Soybean	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Soybean	Watanabe, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Green pepper	Yamada, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Plum	Numabe, Iwaki	Aug-12	Cs137	2.0	Bq/Kg raw	± 2.3	Bq/Kg raw	5.4	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Plum	Numabe, Iwaki	Aug-12	Cs137	7.5	Bq/Kg raw	± 2.8	Bq/Kg raw	13.0	Cs137 7.0 Bq/Kg raw
			Cs134	5.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Dried Shiitake mushroom	Kawauchi, Futaba	Unknown	Cs137	140	Bq/Kg raw	± 34.0	Bq/Kg raw	225	Cs137 7.0 Bq/Kg raw
			Cs134	85.3	Bq/Kg raw	± 21.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Dried Maitake mushroom	Kawauchi, Futaba	Unknown	Cs137	235	Bq/Kg raw	± 56.0	441	Cs137	7.0 Bq/Kg raw
			Cs134	206	Bq/Kg raw	± 46.0		Cs134	6.0 Bq/Kg raw
Egg	Ogawa, Iwaki	Aug-12	Cs137	2.6	Bq/Kg raw	± 1.3	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 1.1		Cs134	6.0 Bq/Kg raw
Japanese Char	Shitoki River, Iwaki(upstream)	Aug-12	Cs137	129	Bq/Kg raw	± 27.0	227	Cs137	7.0 Bq/Kg raw
			Cs134	98.3	Bq/Kg raw	± 19.7		Cs134	6.0 Bq/Kg raw
Watermelon	Ohisa, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.8	Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.2		Cs134	6.0 Bq/Kg raw
Watermelon	Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	4.0 Bq/Kg raw
Watermelon	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Watermelon	Shimoogoe, Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Watermelon	Okaona, Onahama, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Watermelon	Watanabe, Iwaki	Aug-12	Cs137	1.1	Bq/Kg raw	± 1.3	1.1	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Melon	Ohisa, Iwaki	Aug-12	Cs137	1.5	Bq/Kg raw	± 1.6	1.5	Cs137	6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	4.0 Bq/Kg raw
Blueberry	Shimohirakubo, Taira, Iwaki	Jul-12	Cs137	1.8	Bq/Kg raw	± 2.0	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Blueberry	Shimohirakubo, Taira, Iwaki	Aug-12	Cs137	3.1	Bq/Kg raw	± 2.1	4.2	Cs137	7.0 Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 1.2		Cs134	6.0 Bq/Kg raw
Blueberry	Taira, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	2.8	Cs137	6.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 3.4		Cs134	4.0 Bq/Kg raw
Blueberry	Taira, Iwaki	Aug-12	Cs137	10.6	Bq/Kg raw	± 4.0	16.6	Cs137	7.0 Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 3.0		Cs134	6.0 Bq/Kg raw
Blueberry	Sakuragaoka, Iwaki	Aug-12	Cs137	6.7	Bq/Kg raw	± 2.7	6.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Blueberry	Satogaoka, Iwaki	Aug-12	Cs137	7.3	Bq/Kg raw	± 3.3	13.2	Cs137	7.0 Bq/Kg raw
			Cs134	5.9	Bq/Kg raw	± 2.6		Cs134	6.0 Bq/Kg raw
Blueberry	Ishimori, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	4.0 Bq/Kg raw
Blueberry	Joban, Iwaki	Unknown	Cs137	6.3	Bq/Kg raw	± 3.9	6.3	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Pear	Izumi, Iwaki	Jul-12	Cs137	3.7	Bq/Kg raw	± 2.2	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Blueberry jam	Ohara, Onahama, Iwaki	Aug-12	Cs137	6.4	Bq/Kg raw	± 2.5	8.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.9		Cs134	6.0 Bq/Kg raw
Basil sauce	Shimohirakubo, Taira, Iwaki	Unknown	Cs137	2.3	Bq/Kg raw	± 1.2	2.3	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Tomato sauce	Shimohirakubo, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0 Bq/Kg raw
Plum (in sugar)	Izumi, Iwaki	Jun-11	Cs137	14.0	Bq/Kg raw	± 4.0	26.2	Cs137	7.0 Bq/Kg raw
			Cs134	12.2	Bq/Kg raw	± 3.2		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Box lunch	Uchigo, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Box lunch	Uchigo, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Box lunch	Uchigo, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Wire plant	Joban, Iwaki	Unknown	Cs137	20.4	Bq/Kg raw	± 19.3	Bq/Kg raw	51.6	Cs137 7.0 Bq/Kg raw
			Cs134	31.2	Bq/Kg raw	± 16.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Tap water	Izumi, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Tabito, Iwaki	Aug-12	Cs137	0.9	Bq/Kg raw	± 1.2	Bq/Kg raw	0.9	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Tabito, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Spring water	Miwa, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw

