



Radiation Measurement Results of 386 items in October 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Hokkaido	Unknown	Cs137	1.5 Bq/Kg raw	± 2.4 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kumamoto	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Toyama	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Joetsu, Niigata	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Uonuma, Niigata	Oct-12	Cs137	1.7 Bq/Kg raw	± 2.2 Bq/Kg raw	1.7	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Uonuma, Niigata	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Takahagi, Ibaraki	Sep-11	Cs137	9.2 Bq/Kg raw	± 3.5 Bq/Kg raw	17.6	Cs137	7.0 Bq/Kg raw
			Cs134	8.4 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Kitaibaraki, Ibaraki	Oct-12	Cs137	3.3 Bq/Kg raw	± 1.9 Bq/Kg raw	5.9	Cs137	7.0 Bq/Kg raw
			Cs134	2.6 Bq/Kg raw	± 1.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Uchihara, Mito, Ibaraki	Oct-11	Cs137	10.3 Bq/Kg raw	± 3.5 Bq/Kg raw	16.4	Cs137	7.0 Bq/Kg raw
			Cs134	6.1 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Nihonmatsu	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw
			Cs134	1.5 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Ogawa, Iwaki	Oct-12	Cs137	3.8 Bq/Kg raw	± 2.3 Bq/Kg raw	3.8	Cs137	7.0 Bq/Kg raw
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Yotsukura, Iwaki	Oct-12	Cs137	1.7 Bq/Kg raw	± 2.0 Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw
			Cs134	1.5 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Yotsukura, Iwaki	Sep-12	Cs137	2.1 Bq/Kg raw	± 2.2 Bq/Kg raw	3.7	Cs137	7.0 Bq/Kg raw
			Cs134	1.6 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Shimokabeya, Taira, Iwaki	Oct-12	Cs137	2.5 Bq/Kg raw	± 2.8 Bq/Kg raw	4.7	Cs137	7.0 Bq/Kg raw
			Cs134	2.2 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Shimokabeya, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.8 Bq/Kg raw	± 2.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	1.5 Bq/Kg raw	± 2.4 Bq/Kg raw	3.4	Cs137	7.0 Bq/Kg raw
			Cs134	1.9 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Brown rice	Akai, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.2	Cs137	7.0 Bq/Kg raw
			Cs134	1.2 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Brown rice	Takaku, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Komoda, Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 1.9	Bq/Kg raw	
Brown rice	Komoda, Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Kamikuramochi, Kasima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Shimokuramochi , Kasima, Iwaki	Oct-12	Cs137	2.3	Bq/Kg raw	± 2.4	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.1	Bq/Kg raw	
Brown rice	Kashima, Iwaki	Oct-12	Cs137	4.3	Bq/Kg raw	± 2.5	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Kashima, Iwaki	Oct-12	Cs137	1.8	Bq/Kg raw	± 2.3	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Kanari, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.1	Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.0	Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Sep-12	Cs137	11.2	Bq/Kg raw	± 3.9	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	12.2	Bq/Kg raw	± 3.4	Bq/Kg raw	
Brown rice	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	1.4	Bq/Kg raw	± 2.4	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Kamikajiro, Onahama, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw	
Brown rice	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	3.1	Bq/Kg raw	± 2.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Shimokajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.9	Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	20.2	Bq/Kg raw	± 5.2	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	11.6	Bq/Kg raw	± 3.4	Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	2.3	Bq/Kg raw	± 2.6	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw	
Brown rice	Fujiwara, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.1	Bq/Kg raw	
Brown rice	Kadono, Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.2	Bq/Kg raw	
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.9	Bq/Kg raw	
Brown rice	Tono, Iwaki	Sep-12	Cs137	4.5	Bq/Kg raw	± 2.3	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
Brown rice	Tono, Iwaki	Sep-12	Cs137	3.6	Bq/Kg raw	± 2.6	Bq/Kg raw	Cs137 7.0 Bq/Kg raw Cs134 6.0 Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 2.2	Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Tono, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Oct-12	Cs137	4.7	Bq/Kg raw	± 2.5	Bq/Kg raw	7.8	Cs137 7.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Oct-12	Cs137	5.3	Bq/Kg raw	± 2.2	Bq/Kg raw	9.0	Cs137 7.0 Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Sep-12	Cs137	4.1	Bq/Kg raw	± 2.6	Bq/Kg raw	7.5	Cs137 7.0 Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Sep-12	Cs137	3.2	Bq/Kg raw	± 1.4	Bq/Kg raw	4.7	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Sep-12	Cs137	4.0	Bq/Kg raw	± 2.7	Bq/Kg raw	6.7	Cs137 7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Oct-10	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Miyamada, Tono, Iwaki	Oct-12	Cs137	3.6	Bq/Kg raw	± 2.3	Bq/Kg raw	5.4	Cs137 7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Miyamada, Tono, Iwaki	Oct-12	Cs137	2.0	Bq/Kg raw	± 2.4	Bq/Kg raw	2.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Taki, Tono, Iwaki	Oct-12	Cs137	5.4	Bq/Kg raw	± 2.8	Bq/Kg raw	7.1	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Taki, Tono, Iwaki	Oct-12	Cs137	2.0	Bq/Kg raw	± 2.5	Bq/Kg raw	3.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Odaira, Tono, Iwaki	Oct-12	Cs137	4.2	Bq/Kg raw	± 2.5	Bq/Kg raw	7.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Numabe, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Kawabe, Iwaki	Sep-10	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Kawabe, Iwaki	Sep-11	Cs137	3.8	Bq/Kg raw	± 2.7	Bq/Kg raw	9.6	Cs137 7.0 Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Kawabe, Iwaki	Sep-12	Cs137	8.9	Bq/Kg raw	± 3.5	Bq/Kg raw	18.0	Cs137 7.0 Bq/Kg raw
			Cs134	9.1	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Ayanouchi, Nishiki, Iwaki	Oct-12	Cs137	1.5	Bq/Kg raw	± 2.3	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Magarida, Nishiki, Iwaki	Oct-12	Cs137	2.2	Bq/Kg raw	± 2.5	Bq/Kg raw	2.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Nakoso, Iwaki	Sep-12	Cs137	1.4	Bq/Kg raw	± 2.5	Bq/Kg raw	3.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Osawa, Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Soeno, Iwaki	Oct-12	Cs137	3.9	Bq/Kg raw	± 2.2	Bq/Kg raw	6.2	Cs137 7.0 Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Soeno, Iwaki	Sep-12	Cs137	4.4	Bq/Kg raw	± 2.3	Bq/Kg raw	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137 7.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Soeno, Iwaki	Sep-12	Cs137	4.5	Bq/Kg raw	± 2.4	Bq/Kg raw	4.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamadama, Iwaki	Oct-12	Cs137	1.4	Bq/Kg raw	± 2.6	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamadama, Iwaki	Sep-12	Cs137	1.9	Bq/Kg raw	± 2.4	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Seto, Iwaki	Oct-12	Cs137	3.8	Bq/Kg raw	± 2.4	Bq/Kg raw	3.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Seto, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamada, Iwaki	Oct-12	Cs137	1.6	Bq/Kg raw	± 2.2	Bq/Kg raw	1.6	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamada, Iwaki	Oct-12	Cs137	2.0	Bq/Kg raw	± 2.2	Bq/Kg raw	2.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamada, Iwaki	Oct-12	Cs137	2.2	Bq/Kg raw	± 2.5	Bq/Kg raw	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Brown rice	Yamada, Iwaki	Sep-12	Cs137	3.1	Bq/Kg raw	± 2.0	Bq/Kg raw	4.1	Cs137 7.0 Bq/Kg raw
			Cs134	1.0	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Hokkaido	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Hokkaido	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kochi	Oct-11	Cs137	1.5	Bq/Kg raw	± 2.4	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kumamoto	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Miyagi	Oct-11	Cs137	3.8	Bq/Kg raw	± 1.9	Bq/Kg raw	5.9	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 1.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tochigi	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tochigi	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamagata	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.9	Bq/Kg raw	± 1.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamagata	Oct-11	Cs137	1.3	Bq/Kg raw	± 2.0	Bq/Kg raw	1.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
White rice	Yamagata	Sep-12	Cs137	1.7	Bq/Kg raw	± 2.2 Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Niigata	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Uonuma, Niigata	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nagaoka, Niigata	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	2.1	Cs137	7.0	Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Chiba	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.3	Cs137	7.0	Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Yonehara, Shiga	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Akita	Oct-12	Cs137	1.8	Bq/Kg raw	± 2.2 Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Watari, Miyagi	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ibaraki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Takahagi, Ibaraki	Sep-12	Cs137	4.1	Bq/Kg raw	± 2.2 Bq/Kg raw	4.1	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Uchihara, Mito, Ibaraki	Oct-11	Cs137	3.8	Bq/Kg raw	± 1.9 Bq/Kg raw	5.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Hitachi, Ibaraki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nasu, Nasu, Tochigi	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Aizuwakamatsu	Oct-11	Cs137	1.3	Bq/Kg raw	± 2.2 Bq/Kg raw	1.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Aizuwakamatsu	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Aizuwakamatsu	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Hirata, Ishikawa	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	2.1	Cs137	7.0	Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Koriyama	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Koriyama	Oct-12	Cs137	4.5	Bq/Kg raw	± 2.4 Bq/Kg raw	4.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Tamura	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ogawa, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ogawa, Iwaki	Oct-12	Cs137	1.4	Bq/Kg raw	± 2.3 Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Yotsukura, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yotsukura, Iwaki	Sep-12	Cs137	1.8	Bq/Kg raw	± 2.2	Bq/Kg raw	1.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Komagome, Yotsukura, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yoshima, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kamiogoe, Taira, Iwaki	Oct-10	Cs137	2.1	Bq/Kg raw	± 2.2	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kamiogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kamiogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Numanouchi, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Shimotakaku, Taira, Iwaki	Oct-11	Cs137	1.7	Bq/Kg raw	± 1.8	Bq/Kg raw	4.8	Cs137 6.0 Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 1.6	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
White rice	Shimokabeya, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Takaku, Taira, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.4	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137 7.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	4.9	Bq/Kg raw	± 2.2	Bq/Kg raw	4.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	6.4	Bq/Kg raw	± 3.0	Bq/Kg raw	12.3	Cs137 7.0 Bq/Kg raw
			Cs134	5.9	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Okaona, Onahama, Iwaki	Sep-12	Cs137	3.1	Bq/Kg raw	± 2.1	Bq/Kg raw	7.1	Cs137 7.0 Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Ohara, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Misawa, Joban, Iwaki	Sep-12	Cs137	1.9	Bq/Kg raw	± 2.2	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
White rice	Misawa, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.8	Bq/Kg raw	
White rice	Shirakawa, Joban, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Fujiwara, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Watanabe, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Watanabe, Iwaki	Unknown	Cs137	1.2	Bq/Kg raw	± 1.7	Bq/Kg raw	Cs137 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Matsugoya, Watanabe, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-10	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.2	Bq/Kg raw	
White rice	Tono, Iwaki	Sep-12	Cs137	3.5	Bq/Kg raw	± 2.6	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	3.8	Bq/Kg raw	± 2.2	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.1	Bq/Kg raw	
White rice	Tono, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 6.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.0	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.1	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Tono, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.9	Bq/Kg raw	
White rice	Odaira, Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Taki, Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.9	Bq/Kg raw	
White rice	Taki, Tono, Iwaki	Sep-12	Cs137	4.3	Bq/Kg raw	± 2.4	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Taki, Tono, Iwaki	Oct-12	Cs137	2.1	Bq/Kg raw	± 2.5	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Miyamada, Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Nakaoka, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	
White rice	Kawabe, Iwaki	Oct-09	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.1	Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
White rice	Kawabe, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kawabe, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137 7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Tabito, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137 7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Oct-12	Cs137	1.4	Bq/Kg raw	± 2.1	Bq/Kg raw	1.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137 7.0 Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Soeno, Iwaki	Sep-12	Cs137	1.2	Bq/Kg raw	± 1.6	Bq/Kg raw	1.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Osawa, Soeno, Iwaki	Oct-12	Cs137	1.0	Bq/Kg raw	± 1.1	Bq/Kg raw	1.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.8	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamadama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamadama, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Komoda, Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamadama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.8	Cs137 6.0 Bq/Kg raw
			Cs134	0.8	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134 4.0 Bq/Kg raw
White rice	Yamadama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamadama, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137 7.0 Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Yamadama, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nakoso, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Kubota, Nakoso, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nakoso, Iwaki	Oct-11	Cs137	4.8	Bq/Kg raw	± 2.4	Bq/Kg raw	4.8	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Nishiki, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.2	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Unknown	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137 7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
White rice	Unknown	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Brown sticky rice	Shimokuramochi, Kasima, Iwaki	Oct-12	Cs137	2.0	Bq/Kg raw	± 2.3 Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown sticky rice	Shimokajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown sticky rice	Numabe, Iwaki	Unknown	Cs137	1.5	Bq/Kg raw	± 2.4 Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown sticky rice	Yamadama, Iwaki	Sep-12	Cs137	13.2	Bq/Kg raw	± 3.6 Bq/Kg raw	23.4	Cs137	7.0	Bq/Kg raw
			Cs134	10.2	Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Toyama	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Joban, Iwaki	Oct-11	Cs137	5.3	Bq/Kg raw	± 2.6 Bq/Kg raw	5.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Tono, Iwaki	Sep-12	Cs137	2.0	Bq/Kg raw	± 2.1 Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Akasaka, Kawabe, Iwaki	Aug-12	Cs137	4.5	Bq/Kg raw	± 2.0 Bq/Kg raw	7.9	Cs137	7.0	Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Black rice	Date	Oct-11	Cs137	12.4	Bq/Kg raw	± 4.0 Bq/Kg raw	23.1	Cs137	7.0	Bq/Kg raw
			Cs134	10.7	Bq/Kg raw	± 3.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Steamed rice	Akai, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Rice bran	Unknown	Unknown	Cs137	7.0	Bq/Kg raw	± 5.0 Bq/Kg raw	13.0	Cs137	7.0	Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato (with skin)	Yamada, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Taira, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.9 Bq/Kg raw	3.4	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	2.3	Cs137	7.0	Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	2.3	Bq/Kg raw	± 2.3 Bq/Kg raw	3.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	2.2	Cs137	7.0	Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.3	Cs137	7.0	Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Satogaoka, Taira, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Sweet potato	Komoda, Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potato	Okaona, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potato	Kawabe, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potato	Kawabe, Iwaki	Oct-12	Cs137	4.5	Bq/Kg raw	± 2.7	Bq/Kg raw	4.5	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potato	Yamada, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potato	Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.1	Cs137 7.0 Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Sweet potato	Ushiroda, Ueda, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Yacon	Shimohirakubo, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Yam	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.6	Cs137 7.0 Bq/Kg raw
			Cs134	2.6	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese yam	Yoshima, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.2	Bq/Kg raw	1.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Numabe, Iwaki	Jul-12	Cs137	2.4	Bq/Kg raw	± 3.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Ueda, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Potato	Yamada, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Taro	Kitsunezuka, Yotsukura, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Taro	Shimotakaku, Taira, Iwaki	Unknown	Cs137	3.1	Bq/Kg raw	± 3.5	Bq/Kg raw	3.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Taro	Shimoogoe, Taira, Iwaki	Sep-12	Cs137	6.2	Bq/Kg raw	± 3.1	Bq/Kg raw	6.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Taro	Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Pumpkin	Kitsunezuka, Yotsukura, Iwaki	Sep-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Pumpkin	Midaisakai, Uchigo, Iwaki	Sep-12	Cs137	3.9	Bq/Kg raw	± 2.1	Bq/Kg raw	3.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Pumpkin	Nakoso, Iwaki	Aug-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137 7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Carrot	Nagai, Yamagata	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese radish	Ueda, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Japanese radish leaf	Ena, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Turnip	Kashima, Iwaki	Oct-12	Cs137	3.2	Bq/Kg raw	± 5.6	Bq/Kg raw	3.2	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leek	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leek	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leek	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	3.0	Bq/Kg raw	± 3.1	Bq/Kg raw	3.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leek	Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leek	Yamada, Iwaki	Oct-12	Cs137	2.1	Bq/Kg raw	± 2.8	Bq/Kg raw	2.1	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Leek	Nishiki, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese chive	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	2.9	Bq/Kg raw	± 4.4	Bq/Kg raw	2.9	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese chive	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	10.7	Bq/Kg raw	± 5.6	Bq/Kg raw	16.8	Cs137 7.0 Bq/Kg raw
			Cs134	6.1	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese basil seed and leaf	Sakuragaoka, Iwaki	Sep-12	Cs137	15.1	Bq/Kg raw	± 8.0	Bq/Kg raw	20.5	Cs137 7.0 Bq/Kg raw
			Cs134	5.4	Bq/Kg raw	± 6.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Lettuce	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.4	Cs137 7.0 Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 5.1	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Broccoli	Nagai, Yamagata	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Cauliflower	Okaona, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese spinach	Nishigo, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.2	Cs137 7.0 Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbage	Yoshima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137 7.0 Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Chinese cabbage	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.1	Cs137 7.0 Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 1.5	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Japanese mustard spinach	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	2.3	Bq/Kg raw	± 2.6	Bq/Kg raw	2.3	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Egg plant	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Green pepper	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Scallion	Kamiyunagaya, Joban, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Peanut	Shimotakaku, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw
Peanut	Shimohirakubo, Taira, Iwaki	Oct-12	Cs137	6.7	Bq/Kg raw	± 3.4	Bq/Kg raw	6.7	Cs137 7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Peanut	Hirakubo, Taira, Iwaki	Oct-12	Cs137	0.3	Bq/Kg raw	± 0.1 Bq/Kg raw	0.4	Cs137	8.0	Bq/Kg raw
			Cs134	0.1	Bq/Kg raw	± 0.7 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Tea leaf	Ibaraki	Jun-12	Cs137	83.4	Bq/Kg raw	± 19.1 Bq/Kg raw	138	Cs137	7.0	Bq/Kg raw
			Cs134	54.2	Bq/Kg raw	± 12.6 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tea leaf	Shizuoka	May-11	Cs137	171	Bq/Kg raw	± 36.0 Bq/Kg raw	292	Cs137	7.0	Bq/Kg raw
			Cs134	121	Bq/Kg raw	± 25.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Tea leaf	Kagoshima	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Bay leaf	Sakuragaoka, Iwaki	Sep-12	Cs137	5.5	Bq/Kg raw	± 7.5 Bq/Kg raw	5.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Buckwheat seed	Shimoyamaguchi , Taira, Iwaki	Oct-12	Cs137	10.1	Bq/Kg raw	± 4.6 Bq/Kg raw	17.4	Cs137	7.0	Bq/Kg raw
			Cs134	7.3	Bq/Kg raw	± 3.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Oldham Blueberry	Wakabadai, Iwaki	Oct-12	Cs137	17.2	Bq/Kg raw	± 5.1 Bq/Kg raw	27.1	Cs137	7.0	Bq/Kg raw
			Cs134	9.9	Bq/Kg raw	± 3.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Ginkgo	Tabito, Iwaki	Oct-12	Cs137	10.6	Bq/Kg raw	± 3.7 Bq/Kg raw	21.9	Cs137	7.0	Bq/Kg raw
			Cs134	11.3	Bq/Kg raw	± 3.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Citron	Suginami, Taira, Iwaki	Oct-12	Cs137	21.6	Bq/Kg raw	± 5.7 Bq/Kg raw	39.8	Cs137	7.0	Bq/Kg raw
			Cs134	18.2	Bq/Kg raw	± 4.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Citron	Komoda, Kashima, Iwaki	Oct-12	Cs137	43.4	Bq/Kg raw	± 9.7 Bq/Kg raw	69.2	Cs137	7.0	Bq/Kg raw
			Cs134	25.8	Bq/Kg raw	± 5.9 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Citron	Tono, Iwaki	Oct-12	Cs137	30.2	Bq/Kg raw	± 7.4 Bq/Kg raw	64.0	Cs137	7.0	Bq/Kg raw
			Cs134	33.8	Bq/Kg raw	± 7.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Akebi (pulp)	Soeno, Iwaki	Oct-12	Cs137	4.0	Bq/Kg raw	± 1.9 Bq/Kg raw	7.7	Cs137	7.0	Bq/Kg raw
			Cs134	3.7	Bq/Kg raw	± 1.6 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Noda, Chiba	Oct-12	Cs137	4.9	Bq/Kg raw	± 3.3 Bq/Kg raw	11.8	Cs137	7.0	Bq/Kg raw
			Cs134	6.9	Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Naraha, Futaba	Oct-12	Cs137	398	Bq/Kg raw	± 80.0 Bq/Kg raw	670	Cs137	7.0	Bq/Kg raw
			Cs134	272	Bq/Kg raw	± 54.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Akai, Taira, Iwaki	Oct-12	Cs137	10.8	Bq/Kg raw	± 5.0 Bq/Kg raw	18.8	Cs137	7.0	Bq/Kg raw
			Cs134	8.0	Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Shimokabeya, Taira, Iwaki	Oct-12	Cs137	11.3	Bq/Kg raw	± 4.9 Bq/Kg raw	21.3	Cs137	7.0	Bq/Kg raw
			Cs134	10.0	Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Kashima, Iwaki	Oct-12	Cs137	11.1	Bq/Kg raw	± 4.6 Bq/Kg raw	20.1	Cs137	7.0	Bq/Kg raw
			Cs134	9.0	Bq/Kg raw	± 3.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Tabito, Iwaki	Oct-12	Cs137	72.9	Bq/Kg raw	± 15.7 Bq/Kg raw	101	Cs137	8.0	Bq/Kg raw
			Cs134	28.5	Bq/Kg raw	± 8.8 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Chestnut	Tabito, Iwaki	Sep-12	Cs137	94.9	Bq/Kg raw	± 17.7 Bq/Kg raw	143	Cs137	8.0	Bq/Kg raw
			Cs134	48.5	Bq/Kg raw	± 10.1 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Chestnut	Joban, Iwaki	Oct-12	Cs137	11.1	Bq/Kg raw	± 5.3 Bq/Kg raw	22.8	Cs137	7.0	Bq/Kg raw
			Cs134	11.7	Bq/Kg raw	± 4.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Izumi, Iwaki	Oct-12	Cs137	7.6	Bq/Kg raw	± 4.0 Bq/Kg raw	11.2	Cs137	7.0	Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut	Izumi, Iwaki	Oct-12	Cs137	14.3	Bq/Kg raw	± 5.2 Bq/Kg raw	27.8	Cs137	7.0	Bq/Kg raw
			Cs134	13.5	Bq/Kg raw	± 4.4 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chestnut(boiled)	Numabe, Iwaki	Sep-12	Cs137	19.8	Bq/Kg raw	± 5.3 Bq/Kg raw	35.7	Cs137	7.0	Bq/Kg raw
			Cs134	15.9	Bq/Kg raw	± 4.2 Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Chestnut(boiled)	Yamada, Iwaki	Sep-12	Cs137	14.5	Bq/Kg raw	± 4.7	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
				10.6	Bq/Kg raw	± 3.6	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Apple	Ohara, Onahama, Iwaki	Oct-12	Cs137	5.8	Bq/Kg raw	± 11.7	Bq/Kg raw		Cs137	8.0 Bq/Kg raw
			Cs134	3.2	Bq/Kg raw	± 5.5	Bq/Kg raw			Cs134 8.0 Bq/Kg raw
Persimmon	Hobara, Date	Oct-12	Cs137	11.0	Bq/Kg raw	± 4.1	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	11.4	Bq/Kg raw	± 3.5	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Akai, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Hashirikuma, Kashima, Iwaki	Oct-12	Cs137	4.3	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Hashirikuma, Kashima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.9	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Nishigo, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.1	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Okaona, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.7	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	6.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Shimokajiro, Onahama, Iwaki	Oct-12	Cs137	2.7	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.5	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Hayashinoue, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 3.4	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Numabe, Iwaki	Oct-12	Cs137	5.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 2.2	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Yamada, Iwaki	Oct-12	Cs137	3.7	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Persimmon	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Fig	Shimokabeya, Taira, Iwaki	Oct-12	Cs137	7.9	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 2.5	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Kiwi fruit	Ohara, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs137	8.0 Bq/Kg raw
			Cs134	5.3	Bq/Kg raw	± 5.1	Bq/Kg raw			Cs134 8.0 Bq/Kg raw
Kiwi fruit	Komoda, Kashima, Iwaki	Oct-12	Cs137	6.3	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	6.7	Bq/Kg raw	± 2.6	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Kiwi fruit	Kawabe, Iwaki	Oct-12	Cs137	1.9	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 2.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Blueberry	Shimoyoshima, Yoshima, Iwaki	Sep-12	Cs137	8.7	Bq/Kg raw	± 7.0	Bq/Kg raw		Cs137	8.0 Bq/Kg raw
			Cs134	0.1	Bq/Kg raw	± 0.2	Bq/Kg raw			Cs134 8.0 Bq/Kg raw
Blueberry	Chuodai, Taira, Iwaki	Aug-12	Cs137	9.7	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	4.5	Bq/Kg raw	± 2.3	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Blueberry	Okaona, Onahama, Iwaki	Sep-12	Cs137	2.9	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 2.2	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Blueberry	Izumigaoka, Iwaki	Oct-12	Cs137	10.0	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	4.5	Bq/Kg raw	± 2.2	Bq/Kg raw			Cs134 6.0 Bq/Kg raw
Fig	Shimohirakubo, Taira, Iwaki	Oct-12	Cs137	1.8	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw			Cs134 6.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Fig (with skin)	Nishiki, Iwaki	Oct-12	Cs137	1.2	Bq/Kg raw	± 2.2 Bq/Kg raw	1.2	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shimeji mushroom	Akai, Taira, Iwaki	Oct-12	Cs137	19.3	Bq/Kg raw	± 5.8 Bq/Kg raw	31.2	Cs137	7.0	Bq/Kg raw
			Cs134	11.9	Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Hatake shimeji mushroom	Miwa, Iwaki	Oct-12	Cs137	161	Bq/Kg raw	± 25.6 Bq/Kg raw	232	Cs137	8.0	Bq/Kg raw
			Cs134	70.9	Bq/Kg raw	± 12.8 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Ipponshimeji mushroom	Tabito, Iwaki	Oct-12	Cs137	136	Bq/Kg raw	± 28.0 Bq/Kg raw	218	Cs137	7.0	Bq/Kg raw
			Cs134	81.5	Bq/Kg raw	± 16.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Ipponshimeji mushroom	Ogawa, Iwaki	Oct-12	Cs137	3708	Bq/Kg raw	± 338 Bq/Kg raw	5,529	Cs137	8.0	Bq/Kg raw
			Cs134	1821	Bq/Kg raw	± 169 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Ipponshimeji mushroom	Obama, Iwaki	Oct-12	Cs137	33.0	Bq/Kg raw	± 10.4 Bq/Kg raw	43.4	Cs137	8.0	Bq/Kg raw
			Cs134	10.4	Bq/Kg raw	± 6.7 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
shakashimeji mushroom	Onigajo, Iwaki	Oct-12	Cs137	106	Bq/Kg raw	± 22.0 Bq/Kg raw	179	Cs137	7.0	Bq/Kg raw
			Cs134	72.9	Bq/Kg raw	± 14.6 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
shakashimeji mushroom (boiled)	Onigajo, Iwaki	Oct-12	Cs137	11.8	Bq/Kg raw	± 4.0 Bq/Kg raw	21.9	Cs137	7.0	Bq/Kg raw
			Cs134	10.1	Bq/Kg raw	± 3.2 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sakurashimeji mushroom	Taira, Iwaki	Oct-12	Cs137	74.5	Bq/Kg raw	± 15.7 Bq/Kg raw	128	Cs137	7.0	Bq/Kg raw
			Cs134	53.8	Bq/Kg raw	± 11.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sakurashimeji mushroom	Onigajo, Iwaki	Oct-12	Cs137	481	Bq/Kg raw	± 55.2 Bq/Kg raw	715	Cs137	8.0	Bq/Kg raw
			Cs134	234	Bq/Kg raw	± 28.2 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Murasakishimeji mushroom	Kawauchi, Futaba	Oct-12	Cs137	2,340	Bq/Kg raw	± 470 Bq/Kg raw	3,950	Cs137	7.0	Bq/Kg raw
			Cs134	1,610	Bq/Kg raw	± 320 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Honshimeji mushroom	Ogawa, Iwaki	Oct-12	Cs137	15.5	Bq/Kg raw	± 6.1 Bq/Kg raw	22.8	Cs137	7.0	Bq/Kg raw
			Cs134	7.3	Bq/Kg raw	± 4.5 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Hoteishimeji mushroom	Onigajo, Iwaki	Oct-12	Cs137	29.0	Bq/Kg raw	± 8.0 Bq/Kg raw	62.7	Cs137	7.0	Bq/Kg raw
			Cs134	33.7	Bq/Kg raw	± 7.9 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Urabenihoteishimeji mushroom (boiled)	Uchigo, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw	3.4	Cs137	7.0	Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Urabenihoteishimeji mushroom	Uchigo, Iwaki	Oct-12	Cs137	7.0	Bq/Kg raw	± 6.0 Bq/Kg raw	7.0	Cs137	8.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Urabenihoteishimeji mushroom	Uchigo, Iwaki	Oct-12	Cs137	61.7	Bq/Kg raw	± 13.6 Bq/Kg raw	101	Cs137	7.0	Bq/Kg raw
			Cs134	39.6	Bq/Kg raw	± 8.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Urabenihoteishimeji mushroom	Tabito, Iwaki	Oct-12	Cs137	15.9	Bq/Kg raw	± 4.3 Bq/Kg raw	25.8	Cs137	7.0	Bq/Kg raw
			Cs134	9.9	Bq/Kg raw	± 3.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Urabenihoteishimeji mushroom	Tabito, Iwaki	Oct-12	Cs137	1510	Bq/Kg raw	± 300 Bq/Kg raw	3,110	Cs137	7.0	Bq/Kg raw
			Cs134	1600	Bq/Kg raw	± 320 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Maitake mushroom	Kawauchi, Futaba	Oct-12	Cs137	62.1	Bq/Kg raw	± 13.6 Bq/Kg raw	109	Cs137	7.0	Bq/Kg raw
			Cs134	46.8	Bq/Kg raw	± 9.9 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Maitake mushroom	Ogawa, Iwaki	Unknown	Cs137	13.7	Bq/Kg raw	± 6.4 Bq/Kg raw	19.5	Cs137	8.0	Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 5.5 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Maitake mushroom	Ogawa, Iwaki	Oct-12	Cs137	5.2	Bq/Kg raw	± 3.7 Bq/Kg raw	10.6	Cs137	7.0	Bq/Kg raw
			Cs134	5.4	Bq/Kg raw	± 3.2 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Maitake mushroom	Yamadama, Iwaki	Oct-12	Cs137	42.2	Bq/Kg raw	± 13.7 Bq/Kg raw	47.0	Cs137	8.0	Bq/Kg raw
			Cs134	4.8	Bq/Kg raw	± 6.6 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Maitake mushroom	Yokokawa, Tabito, Iwaki	Oct-12	Cs137	9.7	Bq/Kg raw	± 6.0 Bq/Kg raw	16.1	Cs137	8.0	Bq/Kg raw
			Cs134	6.4	Bq/Kg raw	± 4.9 Bq/Kg raw		Cs134	8.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Matsutake mushroom	Ogawa, Iwaki	Oct-12	Cs137	63.4	Bq/Kg raw ± 11.9 Bq/Kg raw	80.2	Cs137	8.0	Bq/Kg raw
			Cs134	16.8	Bq/Kg raw ± 5.8 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Matsutake mushroom	Kawauchi, Iwaki	Oct-12	Cs137	75.4	Bq/Kg raw ± 15.8 Bq/Kg raw	118	Cs137	7.0	Bq/Kg raw
			Cs134	42.7	Bq/Kg raw ± 9.1 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Matsutake mushroom	Naraha, Futaba	Oct-12	Cs137	2,495	Bq/Kg raw ± 239 Bq/Kg raw	3,693	Cs137	8.0	Bq/Kg raw
			Cs134	1,198	Bq/Kg raw ± 118 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Matsutake mushroom	Takaku, Taira, Iwaki	Unknown	Cs137	73.9	Bq/Kg raw ± 22.0 Bq/Kg raw	105	Cs137	8.0	Bq/Kg raw
			Cs134	30.9	Bq/Kg raw ± 13.7 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Matsutake mushroom	Hashirikuma, Kashima, Iwaki	Oct-12	Cs137	153	Bq/Kg raw ± 31.0 Bq/Kg raw	255	Cs137	7.0	Bq/Kg raw
			Cs134	102	Bq/Kg raw ± 20.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Matsutake mushroom	Nyujou, Tono, Iwaki	Oct-12	Cs137	125	Bq/Kg raw ± 30.0 Bq/Kg raw	233	Cs137	7.0	Bq/Kg raw
			Cs134	108	Bq/Kg raw ± 24.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Matsutake mushroom	Taki, Tono, Iwaki	Oct-12	Cs137	249	Bq/Kg raw ± 31.6 Bq/Kg raw	373	Cs137	8.0	Bq/Kg raw
			Cs134	124	Bq/Kg raw ± 16.5 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Matsutake mushroom	Tono, Iwaki	Oct-12	Cs137	84.7	Bq/Kg raw ± 17.8 Bq/Kg raw	145	Cs137	7.0	Bq/Kg raw
			Cs134	60.5	Bq/Kg raw ± 12.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Matsutake mushroom	Tabito, Iwaki	Oct-12	Cs137	64.2	Bq/Kg raw ± 14.6 Bq/Kg raw	104	Cs137	7.0	Bq/Kg raw
			Cs134	40.0	Bq/Kg raw ± 9.2 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Log (for Shiitake)	Komoda, Kashima, Iwaki	Nov-12	Cs137	346.0	Bq/Kg dry ± 69.0 Bq/Kg dry	572	Cs137	7.0	Bq/Kg dry
			Cs134	226.0	Bq/Kg dry ± 45.0 Bq/Kg dry		Cs134	6.0	Bq/Kg dry
Jersey cow mushroom (washed)	Samegawa, Higashishirakawa	Oct-12	Cs137	231	Bq/Kg raw ± 46.0 Bq/Kg raw	474	Cs137	7.0	Bq/Kg raw
			Cs134	243	Bq/Kg raw ± 49.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jersey cow mushroom	Hanawa, Higashishirakawa	Oct-12	Cs137	41.4	Bq/Kg raw ± 9.5 Bq/Kg raw	70.9	Cs137	7.0	Bq/Kg raw
			Cs134	29.5	Bq/Kg raw ± 6.7 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jersey cow mushroom	Samegawa, Higashishirakawa	Oct-12	Cs137	622	Bq/Kg raw ± 124 Bq/Kg raw	1,259	Cs137	7.0	Bq/Kg raw
			Cs134	637	Bq/Kg raw ± 127 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jersey cow mushroom	Onigajo, Iwaki	Oct-12	Cs137	388	Bq/Kg raw ± 78.0 Bq/Kg raw	790	Cs137	7.0	Bq/Kg raw
			Cs134	402	Bq/Kg raw ± 80.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jersey cow mushroom	Misaka, Miwa, Iwaki	Oct-12	Cs137	210	Bq/Kg raw ± 42.0 Bq/Kg raw	358	Cs137	7.0	Bq/Kg raw
			Cs134	148	Bq/Kg raw ± 30.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jersey cow mushroom	Koya, Uchigo, Iwaki	Oct-12	Cs137	139	Bq/Kg raw ± 28.0 Bq/Kg raw	275	Cs137	7.0	Bq/Kg raw
			Cs134	136	Bq/Kg raw ± 27.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jersey cow mushroom (washed)	Onigajo, Iwaki	Oct-12	Cs137	26.0	Bq/Kg raw ± 6.4 Bq/Kg raw	48.7	Cs137	7.0	Bq/Kg raw
			Cs134	22.7	Bq/Kg raw ± 5.3 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kuritake mushroom	Iwade, Onahama, Iwaki	Oct-12	Cs137	149	Bq/Kg raw ± 30.0 Bq/Kg raw	253	Cs137	7.0	Bq/Kg raw
			Cs134	104	Bq/Kg raw ± 21.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shishitake mushroom	Ogawa, Iwaki	Oct-12	Cs137	140	Bq/Kg raw ± 21.3 Bq/Kg raw	180	Cs137	8.0	Bq/Kg raw
			Cs134	40.1	Bq/Kg raw ± 9.0 Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Koganetake mushroom	Minamiodaira, Tabito, Iwaki	Oct-12	Cs137	5.6	Bq/Kg raw ± 2.9 Bq/Kg raw	7.0	Cs137	7.0	Bq/Kg raw
			Cs134	1.4	Bq/Kg raw ± 2.4 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Hatsutake mushroom	Koya, Uchigo, Iwaki	Oct-12	Cs137	460	Bq/Kg raw ± 92.0 Bq/Kg raw	760	Cs137	7.0	Bq/Kg raw
			Cs134	300	Bq/Kg raw ± 60.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yamatoritakemodoki mushroom	Chuodai, Taira, Iwaki	Oct-12	Cs137	39.9	Bq/Kg raw ± 9.8 Bq/Kg raw	77.5	Cs137	7.0	Bq/Kg raw
			Cs134	37.6	Bq/Kg raw ± 8.6 Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Koutake mushroom	Miwa, Iwaki	Oct-12	Cs137	172	Bq/Kg raw ± 35.0 Bq/Kg raw	245	Cs137	7.0	Bq/Kg raw
			Cs134	73.4	Bq/Kg raw ± 16.0 Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection		
Koutake(boiled)	Onigajo, Iwaki	Oct-12	Cs137	7.4	Bq/Kg raw	± 30.6	Bq/Kg raw	9.1	Cs137	8.0 Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Koutake(boiled)	Miwa, Iwaki	Oct-12	Cs137	17.5	Bq/Kg raw	± 4.7	Bq/Kg raw	29.4	Cs137	7.0 Bq/Kg raw
			Cs134	11.9	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Koutake(boiled)	Unknown	Unknown	Cs137	3.3	Bq/Kg raw	± 3.5	Bq/Kg raw	3.6	Cs137	8.0 Bq/Kg raw
			Cs134	0.3	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Koutake (boiled twice in salt water)	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Kurihusentake mushroom	Miwa, Iwaki	Oct-12	Cs137	109	Bq/Kg raw	± 23.0	Bq/Kg raw	173	Cs137	7.0 Bq/Kg raw
			Cs134	63.7	Bq/Kg raw	± 13.5	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Kusaurabenitake mushroom	Miwa, Iwaki	Oct-12	Cs137	4570	Bq/Kg raw	± 910	Bq/Kg raw	7,660	Cs137	7.0 Bq/Kg raw
			Cs134	3090	Bq/Kg raw	± 620	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Nameko mushroom	Iwai, Onahama, Iwaki	Oct-12	Cs137	528	Bq/Kg raw	± 106	Bq/Kg raw	878	Cs137	7.0 Bq/Kg raw
			Cs134	350	Bq/Kg raw	± 70.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Nameko mushroom	Tabito, Iwaki	Unknown	Cs137	205	Bq/Kg raw	± 29.0	Bq/Kg raw	303	Cs137	8.0 Bq/Kg raw
			Cs134	98.0	Bq/Kg raw	± 15.6	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Honey	Ohisa, Iwaki	Oct-12	Cs137	86.4	Bq/Kg raw	± 18.7	Bq/Kg raw	186	Cs137	7.0 Bq/Kg raw
			Cs134	100	Bq/Kg raw	± 20.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Honey	Yotsukura, Iwaki	Oct-12	Cs137	54.2	Bq/Kg raw	± 11.8	Bq/Kg raw	94.4	Cs137	7.0 Bq/Kg raw
			Cs134	40.2	Bq/Kg raw	± 8.4	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Honey	Tabito, Iwaki	Oct-12	Cs137	130	Bq/Kg raw	± 26.0	Bq/Kg raw	221	Cs137	7.0 Bq/Kg raw
			Cs134	90.7	Bq/Kg raw	± 18.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Miso	Tochigi	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Miso	Tochigi	Feb-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.3	Cs137	7.0 Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Miso	Iwaki	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
School lunch	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
School lunch	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Salmon	Tomioka, Futaba	Oct-12	Cs137	4.9	Bq/Kg raw	± 2.5	Bq/Kg raw	8.8	Cs137	7.0 Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Marlin	Ena, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Young tuna	Onahama, Iwaki	Oct-12	Cs137	1.4	Bq/Kg raw	± 2.2	Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Mahi mahi	Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0 Bq/Kg raw
Mussel	Onahama, Iwaki	Oct-12	Cs137	1.0	Bq/Kg raw	± 4.3	Bq/Kg raw	3.4	Cs137	8.0 Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 4.2	Bq/Kg raw		Cs134	8.0 Bq/Kg raw
Gourd water	Shimohirakubo, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	3.0 Bq/Kg raw
Mineral water	Kitaazumi, Nagano	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	3.0 Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Mineral water	Hokkaido	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap water	Yumoto, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Tap water	Nakoso, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.5	Cs137 5.0 Bq/Kg raw
			Cs134	0.5	Bq/Kg raw	± 0.8	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Well water	Kabeya, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137 5.0 Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134 3.0 Bq/Kg raw
Japanese torreya	Honya, Izumi, Iwaki	Oct-12	Cs137	12.0	Bq/Kg dry	± 10.0	Bq/Kg dry	42	Cs137 7.0 Bq/Kg dry
			Cs134	30.0	Bq/Kg dry	± 10.0	Bq/Kg dry		Cs134 6.0 Bq/Kg dry
Ash	Honya, Izumi, Iwaki	Nov-12	Cs137	972.0	Bq/Kg dry	± 194.0	Bq/Kg dry	1,643	Cs137 7.0 Bq/Kg dry
			Cs134	671.0	Bq/Kg dry	± 134.0	Bq/Kg dry		Cs134 6.0 Bq/Kg dry
Bamboo	Honya, Izumi, Iwaki	Oct-12	Cs137	43.0	Bq/Kg dry	± 12.0	Bq/Kg dry	67	Cs137 7.0 Bq/Kg dry
			Cs134	24.0	Bq/Kg dry	± 8.0	Bq/Kg dry		Cs134 6.0 Bq/Kg dry
Pottery	Ohisa, Iwaki	Jul-12	Cs137	27.0	Bq/Kg dry	± 8.0	Bq/Kg dry	89	Cs137 7.0 Bq/Kg dry
			Cs134	62.0	Bq/Kg dry	± 13.0	Bq/Kg dry		Cs134 6.0 Bq/Kg dry
Pottery	Ohisa, Iwaki	Jul-12	Cs137	47.0	Bq/Kg dry	± 13.0	Bq/Kg dry	57	Cs137 8.0 Bq/Kg dry
			Cs134	10.0	Bq/Kg dry	± 7.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Leaf mold	Nishikimigatsu ka, Onahama, Iwaki	Oct-12	Cs137	20.0	Bq/Kg dry	± 9.0	Bq/Kg dry	23.0	Cs137 8.0 Bq/Kg dry
			Cs134	3.0	Bq/Kg dry	± 6.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Mihara, Hiroshima	Oct-12	Cs137	30.0	Bq/Kg dry	± 10.0	Bq/Kg dry	32.0	Cs137 8.0 Bq/Kg dry
			Cs134	2.0	Bq/Kg dry	± 5.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Mihara, Hiroshima	Oct-12	Cs137	28.0	Bq/Kg dry	± 9.0	Bq/Kg dry	28.0	Cs137 8.0 Bq/Kg dry
			Cs134	0.0	Bq/Kg dry	± 0.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Shimokawa, Izumi, Iwaki	Oct-12	Cs137	501	Bq/Kg dry	± 57.0	Bq/Kg dry	766	Cs137 8.0 Bq/Kg dry
			Cs134	265	Bq/Kg dry	± 31.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Field soil	Shimokawa, Izumi, Iwaki	Oct-12	Cs137	664	Bq/Kg dry	± 72.0	Bq/Kg dry	1,005	Cs137 8.0 Bq/Kg dry
			Cs134	341	Bq/Kg dry	± 38.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Kashima, Chuodai, Iwaki	Unknown	Cs137	708	Bq/Kg dry	± 76.0	Bq/Kg dry	1,080	Cs137 8.0 Bq/Kg dry
			Cs134	372	Bq/Kg dry	± 41.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Izumi, Iwaki	Oct-12	Cs137	682	Bq/Kg dry	± 74.0	Bq/Kg dry	1,032	Cs137 8.0 Bq/Kg dry
			Cs134	350	Bq/Kg dry	± 40.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Izumi, Iwaki	Oct-12	Cs137	707	Bq/Kg dry	± 77.0	Bq/Kg dry	1,086	Cs137 8.0 Bq/Kg dry
			Cs134	379	Bq/Kg dry	± 42.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shimokawa, Izumi, Iwaki	Oct-12	Cs137	312	Bq/Kg dry	± 39.0	Bq/Kg dry	484	Cs137 8.0 Bq/Kg dry
			Cs134	172	Bq/Kg dry	± 22.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shimokawa, Izumi, Iwaki	Oct-12	Cs137	1098	Bq/Kg dry	± 112	Bq/Kg dry	1,671	Cs137 8.0 Bq/Kg dry
			Cs134	573	Bq/Kg dry	± 60.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shimokawa, Izumi, Iwaki	Oct-12	Cs137	1290	Bq/Kg dry	± 134	Bq/Kg dry	1,969	Cs137 8.0 Bq/Kg dry
			Cs134	679	Bq/Kg dry	± 71.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Shimokawa, Izumi, Iwaki	Oct-12	Cs137	661	Bq/Kg dry	± 72.0	Bq/Kg dry	1,018	Cs137 8.0 Bq/Kg dry
			Cs134	357	Bq/Kg dry	± 40.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry
Soil	Honya, Izumi, Iwaki	Oct-12	Cs137	237.0	Bq/Kg dry	± 29.0	Bq/Kg dry	358	Cs137 8.0 Bq/Kg dry
			Cs134	121.0	Bq/Kg dry	± 16.0	Bq/Kg dry		Cs134 8.0 Bq/Kg dry

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Soil	Kubota, Nakoso, Iwaki	Oct-12	Cs137	706	Bq/Kg dry	± 74.0	Bq/Kg dry	Cs137 Cs134
			Cs134	393	Bq/Kg dry	± 41.0	Bq/Kg dry	

