



Radiation Measurement Results of 233 items in November 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Aizuwakamatsu	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw	
			Cs134	1.4 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Aizuwakamatsu	Oct-12	Cs137	1.8 Bq/Kg raw	± 2.5 Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Aizuwakamatsu	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Inawashiro, Yama	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Funahiki, Tamura, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Ogawa, Iwaki	Oct-12	Cs137	6.2 Bq/Kg raw	± 3.1 Bq/Kg raw	12.3	Cs137	7.0 Bq/Kg raw	
			Cs134	6.1 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Yotsukura, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Kamihirakubo, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw	
			Cs134	1.6 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Shimokajiro, Onahama, Iwaki	Oct-12	Cs137	7.3 Bq/Kg raw	± 2.3 Bq/Kg raw	12.3	Cs137	7.0 Bq/Kg raw	
			Cs134	5.0 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.2	Cs137	6.0 Bq/Kg raw	
			Cs134	1.2 Bq/Kg raw	± 1.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Brown rice	Nakoso, Iwaki	Oct-12	Cs137	1.3 Bq/Kg raw	± 1.9 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw	
			Cs134	1.8 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Akita	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Yamagata	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Ishinomaki, Miyagi	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Miyagi	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.3	Cs137	7.0 Bq/Kg raw	
			Cs134	2.3 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
White rice	Shiraishi, Miyagi	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	1.9 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
White rice	Nanyou, Yamagata	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nihonmatsu	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Inawashiro, Yama	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ogawa, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kamiogawa, Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Fukuoka, Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Uwadaira, Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Yotsukura, Iwaki	Nov-12	Cs137	1.5	Bq/Kg raw	± 2.3	Bq/Kg raw	2.8	Cs137	7.0	Bq/Kg raw
			Cs134	1.3	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-11	Cs137	4.0	Bq/Kg raw	± 2.5	Bq/Kg raw	5.9	Cs137	7.0	Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nagatomo, Yotsukura, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Shimoyoshima, Yoshima, Iwaki	Oct-12	Cs137	5.4	Bq/Kg raw	± 2.4	Bq/Kg raw	10.2	Cs137	7.0	Bq/Kg raw
			Cs134	4.8	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Shimoyoshima, Yoshima, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Akai, Taira, Iwaki	Oct-12	Cs137	2.0	Bq/Kg raw	± 2.2	Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nakashio, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Shimotakaku, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kamioogoe, Taira, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Toyoma, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Shimoyamaguchi, Taira, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.4	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kitakabeya, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
White rice	Kuramochi, Kashima, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kamikajiro, Onahama, Iwaki	Oct-12	Cs137	2.1	Bq/Kg raw	± 2.4	Bq/Kg raw	2.1	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kanari, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Sumiyoshi, Onahama, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Misawa, Joban, Iwaki	Oct-12	Cs137	1.6	Bq/Kg raw	± 2.2	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Misawa, Joban, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137	7.0	Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Izumi, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Tono, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kadono, Tono, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kadono, Tono, Iwaki	Oct-12	Cs137	1.9	Bq/Kg raw	± 2.1	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ueda, Iwaki	Oct-11	Cs137	1.4	Bq/Kg raw	± 2.2	Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Numabe, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kawabe, Iwaki	Oct-12	Cs137	3.4	Bq/Kg raw	± 1.4	Bq/Kg raw	5.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 1.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kawabe, Iwaki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Soeno, Iwaki	Oct-12	Cs137	1.6	Bq/Kg raw	± 2.4	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nakoso, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Unknown	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pre-washed white rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	3.4	Bq/Kg raw	± 2.2	Bq/Kg raw	6.3	Cs137	7.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Fried rice	Shimokabeya, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	6.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Sticky brown rice	Kasama, Ibaraki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky brown rice	Nakahirakubo, Taira, Iwaki	Oct-12	Cs137	2.0	Bq/Kg raw	± 2.3	Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky brown rice	Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Sticky brown rice	Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky brown rice	Unknown	Nov-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Nishiura, Niigata, Niigata	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Ogawa, Iwaki	Unknown	Cs137	3.8	Bq/Kg raw	± 2.3	Bq/Kg raw	3.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Ogawa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Soeno, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Rice bran	Unknown	Unknown	Cs137	7.0	Bq/Kg raw	± 5.0	Bq/Kg raw	13.0	Cs137	7.0	Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 4.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Nagai, Yamagata	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Nakashio, Taira, Iwaki	Nov-12	Cs137	1.9	Bq/Kg raw	± 2.6	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Nakahirakubo, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.4	Cs137	8.0	Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 6.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Taro	Numanouchi, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Unknown	Unknown	Cs137	1.5	Bq/Kg raw	± 2.4	Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Iritono, Tono, Iwaki	Nov-12	Cs137	1.4	Bq/Kg raw	± 2.6	Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Watanabe, Iwaki	Nov-12	Cs137	2.0	Bq/Kg raw	± 3.0	Bq/Kg raw	2.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Jerusalem artichoke	Taira, Iwaki	Nov-12	Cs137	2.7	Bq/Kg raw	± 3.1	Bq/Kg raw	5.6	Cs137	7.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Dried stems of a taro	Nakahirakubo, Taira, Iwaki	Nov-12	Cs137	0.5	Bq/Kg raw	± 13.3	Bq/Kg raw	2.9	Cs137	8.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 6.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Japanese radish	Miwa, Iwaki	Nov-12	Cs137	1.7	Bq/Kg raw	± 2.4	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Ogawa, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Kitsunozuka, Yotsukura, Iwaki	Oct-12	Cs137	5.2	Bq/Kg raw	± 2.7	Bq/Kg raw	8.1	Cs137	6.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Japanese radish	Yotsukura, Iwaki	Nov-12	Cs137	1.6	Bq/Kg raw	± 2.5	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Hosoya, Yotsukura, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Suginami, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Akai, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Japanese radish	Nakashio, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Kamikuramochi, Kasima, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.2	Cs137	7.0	Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Takaku, Chuodai, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Ohara, Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Okaona, Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.3	Cs137	7.0	Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Watanabe, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.2	Cs137	7.0	Bq/Kg raw
			Cs134	2.2	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Taki, Tono, Iwaki	Nov-12	Cs137	1.7	Bq/Kg raw	± 2.4	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yamada, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Purple Japanese radish	Watanabe, Iwaki	Nov-12	Cs137	2.5	Bq/Kg raw	± 3.4	Bq/Kg raw	4.3	Cs137	7.0	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Turnip	Ogawa, Iwaki	Nov-12	Cs137	2.1	Bq/Kg raw	± 2.9	Bq/Kg raw	2.1	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Turnip leaf	Ogawa, Iwaki	Nov-12	Cs137	2.7	Bq/Kg raw	± 3.6	Bq/Kg raw	2.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Miwa, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Komagome, Yotsukura, Iwaki	Nov-12	Cs137	4.6	Bq/Kg raw	± 2.9	Bq/Kg raw	6.6	Cs137	7.0	Bq/Kg raw
			Cs134	2.0	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Akai, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Takaku, Chuodai, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Kuramochi, Kasima, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Okaona, Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Ohara, Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Iritono, Tono, Iwaki	Nov-12	Cs137	1.5	Bq/Kg raw	± 2.1	Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Taki, Tono, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Watanabe, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Yotsukura, Iwaki	Nov-12	Cs137	20.0	Bq/Kg raw	± 5.9	Bq/Kg raw	28.4	Cs137	7.0	Bq/Kg raw
			Cs134	8.4	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Sweet potato	Nakashio, Taira, Iwaki	Oct-12	Cs137	2.5 Bq/Kg raw	± 2.7 Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sweet potato (with skin)	Kitaibaraki, Ibaraki	Oct-12	Cs137	4.0 Bq/Kg raw	± 2.2 Bq/Kg raw	5.5	Cs137	7.0 Bq/Kg raw	
			Cs134	1.5 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Pumpkin	Azumino, Nagano	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Kitsunozuka, Yotsukura, Iwaki	Oct-12	Cs137	2.4 Bq/Kg raw	± 3.1 Bq/Kg raw	4.4	Cs137	7.0 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Midaisakai, Uchigo, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Carrot	Kashima, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Komagome, Yotsukura, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Yamada, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Akai, Taira, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Ogawa, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Takaku, Taira, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Nagai, Yamagata	Nov-12	Cs137	1.9 Bq/Kg raw	± 2.3 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Akai, Taira, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Leek	Ohara, Onahama, Iwaki	Nov-12	Cs137	3.1 Bq/Kg raw	± 3.1 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Chinese chive	Numanouchi, Taira, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.6	Cs137	7.0 Bq/Kg raw	
			Cs134	4.6 Bq/Kg raw	± 5.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Chinese chive	Shimoogoe, Taira, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.0	Cs137	7.0 Bq/Kg raw	
			Cs134	3.0 Bq/Kg raw	± 3.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Spinach	Suginami, Taira, Iwaki	Nov-12	Cs137	2.7 Bq/Kg raw	± 3.0 Bq/Kg raw	4.3	Cs137	7.0 Bq/Kg raw	
			Cs134	1.6 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Spinach	Ogawa, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Spinach	Okaona, Onahama, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw	
			Cs134	2.1 Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Spinach	Yamada, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Cabbage	Okaona, Onahama, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Crown daisy	Wakabadai, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Bok choy	Akai, Taira, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Kyou-mizuna	Akai, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese mustard spinach	Akai, Taira, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese mustard spinach	Kamikuramochi, Kasima, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Iritono, Tono, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Wild sesame	Fujima, Taira, Iwaki	Oct-12	Cs137	25.3	Bq/Kg raw	± 7.5	Bq/Kg raw	42.9	Cs137	7.0	Bq/Kg raw
			Cs134	17.6	Bq/Kg raw	± 5.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yacon	Hosoya, Yotsukura, Iwaki	Nov-12	Cs137	1.4	Bq/Kg raw	± 2.7	Bq/Kg raw	3.3	Cs137	7.0	Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Green chill	Izumi, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Blue soybean	Niigata	Nov-11	Cs137	14.4	Bq/Kg raw	± 4.6	Bq/Kg raw	20.2	Cs137	7.0	Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Tono, Iwaki	Nov-12	Cs137	139	Bq/Kg raw	± 28.0	Bq/Kg raw	232	Cs137	7.0	Bq/Kg raw
			Cs134	92.6	Bq/Kg raw	± 18.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Izumi, Iwaki	Nov-12	Cs137	147	Bq/Kg raw	± 24.6	Bq/Kg raw	232	Cs137	8.0	Bq/Kg raw
			Cs134	84.5	Bq/Kg raw	± 13.4	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shiitake mushroom	Yotsukura, Iwaki	Oct-12	Cs137	241	Bq/Kg raw	± 48.0	Bq/Kg raw	403	Cs137	7.0	Bq/Kg raw
			Cs134	162	Bq/Kg raw	± 32.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Shimokawa, Izumi, Iwaki	Nov-12	Cs137	83.6	Bq/Kg raw	± 14.5	Bq/Kg raw	120	Cs137	8.0	Bq/Kg raw
			Cs134	36.8	Bq/Kg raw	± 7.7	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shiitake mushroom	Shimokawa, Izumi, Iwaki	Nov-12	Cs137	83.6	Bq/Kg raw	± 14.5	Bq/Kg raw	120	Cs137	8.0	Bq/Kg raw
			Cs134	36.8	Bq/Kg raw	± 7.7	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shiitake mushroom	Yamada, Iwaki	Nov-12	Cs137	99.0	Bq/Kg raw	± 21.7	Bq/Kg raw	179	Cs137	7.0	Bq/Kg raw
			Cs134	80.1	Bq/Kg raw	± 16.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Kashima, Iwaki	Nov-12	Cs137	266.0	Bq/Kg raw	± 53.0	Bq/Kg raw	443	Cs137	7.0	Bq/Kg raw
			Cs134	177.0	Bq/Kg raw	± 35.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Taira, Iwaki	Nov-12	Cs137	224	Bq/Kg raw	± 46.0	Bq/Kg raw	383	Cs137	7.0	Bq/Kg raw
			Cs134	159	Bq/Kg raw	± 32.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Himesakura shimeji mushroom	Akai, Taira, Iwaki	Nov-12	Cs137	456	Bq/Kg raw	± 52.1	Bq/Kg raw	689	Cs137	8.0	Bq/Kg raw
			Cs134	233	Bq/Kg raw	± 27.6	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Hatake shimeji mushroom	Ishimori, Iwaki	Nov-12	Cs137	10.2	Bq/Kg raw	± 5.8	Bq/Kg raw	13.3	Cs137	8.0	Bq/Kg raw
			Cs134	3.1	Bq/Kg raw	± 5.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Urabenihotei shimeji	Ishimori, Iwaki	Oct-12	Cs137	598	Bq/Kg raw	± 66.5	Bq/Kg raw	874	Cs137	8.0	Bq/Kg raw
			Cs134	276	Bq/Kg raw	± 32.9	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shimofuri shimeji mushroom	Tabito, Iwaki	Nov-12	Cs137	41.7	Bq/Kg raw	± 12.1	Bq/Kg raw	61.1	Cs137	8.0	Bq/Kg raw
			Cs134	19.4	Bq/Kg raw	± 6.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shaka shimeji mushroom (boiled)	Ogawa, Iwaki	Oct-12	Cs137	8.8	Bq/Kg raw	± 5.1	Bq/Kg raw	8.8	Cs137	8.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Koutake mushroom	Miwa, Iwaki	Nov-12	Cs137	351	Bq/Kg raw	± 70.0	Bq/Kg raw	523	Cs137	7.0	Bq/Kg raw
			Cs134	172	Bq/Kg raw	± 35.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Koutake mushroom	Ogawa, Iwaki	Oct-12	Cs137	2.9	Bq/Kg raw	± 3.1	Bq/Kg raw	2.9	Cs137	8.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Koutake mushroom (boiled)	Miwa, Iwaki	Nov-12	Cs137	0.3	Bq/Kg raw	± 5.6	Bq/Kg raw	2.8	Cs137	8.0	Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 4.4	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Koutake mushroom (boiled)	Taira, Iwaki	Oct-12	Cs137	9.5	Bq/Kg raw	± 9.0	Bq/Kg raw	20.1	Cs137	8.0	Bq/Kg raw
			Cs134	10.6	Bq/Kg raw	± 9.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Koganetake mushroom	Ishimori, Iwaki	Nov-12	Cs137	16.9	Bq/Kg raw	± 12.2	Bq/Kg raw	26.5	Cs137	8.0	Bq/Kg raw
			Cs134	9.6	Bq/Kg raw	± 7.5	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shishitake mushroom	Naraha, Futaba	Nov-12	Cs137	1570	Bq/Kg raw	± 153	Bq/Kg raw	2,299	Cs137	8.0	Bq/Kg raw
			Cs134	729	Bq/Kg raw	± 73.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shishitake mushroom	Kawamae, Iwaki	Nov-12	Cs137	1.5	Bq/Kg raw	± 0.1	Bq/Kg raw	2.5	Cs137	8.0	Bq/Kg raw
			Cs134	1.0	Bq/Kg raw	± 3.8	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Akamomitake mushroom	Uchigo, Iwaki	Nov-12	Cs137	452	Bq/Kg raw	± 52.2	Bq/Kg raw	680	Cs137	8.0	Bq/Kg raw
			Cs134	228	Bq/Kg raw	± 27.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Kihatsutake mushroom	Mizuishiyama, Iwaki	Nov-12	Cs137	1200	Bq/Kg raw	± 240	Bq/Kg raw	2,004	Cs137	7.0	Bq/Kg raw
			Cs134	804	Bq/Kg raw	± 161	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kuritake mushroom(boiled)	Tabito, Iwaki	Nov-12	Cs137	56.8	Bq/Kg raw	± 12.5	Bq/Kg raw	94.1	Cs137	8.0	Bq/Kg raw
			Cs134	37.3	Bq/Kg raw	± 8.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Hanaiguchi mushroom(boiled)	Nishigo, Joban, Iwaki	Oct-12	Cs137	43.8	Bq/Kg raw	± 9.9	Bq/Kg raw	75.9	Cs137	7.0	Bq/Kg raw
			Cs134	32.1	Bq/Kg raw	± 7.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mandarin orange	Shimokajiro, Taira, Iwaki	Nov-12	Cs137	13.9	Bq/Kg raw	± 4.4	Bq/Kg raw	23.1	Cs137	7.0	Bq/Kg raw
			Cs134	9.2	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mandarin orange	Satogaoka, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mandarin orange	Shimoyamaguchi , Taira, Iwaki	Nov-12	Cs137	12.4	Bq/Kg raw	± 4.3	Bq/Kg raw	21.8	Cs137	7.0	Bq/Kg raw
			Cs134	9.4	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mandarin orange	Midaisakai, Uchigo, Iwaki	Nov-12	Cs137	6.7	Bq/Kg raw	± 2.6	Bq/Kg raw	9.7	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mandarin orange	Youkoudai, Iwaki	Nov-12	Cs137	15.1	Bq/Kg raw	± 4.1	Bq/Kg raw	22.5	Cs137	7.0	Bq/Kg raw
			Cs134	7.4	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mandarin orange	Soeno, Iwaki	Nov-12	Cs137	11.0	Bq/Kg raw	± 5.2	Bq/Kg raw	11.9	Cs137	8.0	Bq/Kg raw
			Cs134	0.9	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Yuzu	Kitaibaraki, Ibaraki	Nov-12	Cs137	26.1	Bq/Kg raw	± 6.3	Bq/Kg raw	41.6	Cs137	7.0	Bq/Kg raw
			Cs134	15.5	Bq/Kg raw	± 4.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Yotsukura, Iwaki	Oct-12	Cs137	83.5	Bq/Kg raw	± 17.6	Bq/Kg raw	144	Cs137	7.0	Bq/Kg raw
			Cs134	60.3	Bq/Kg raw	± 12.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Nakahirakubo, Taira, Iwaki	Nov-12	Cs137	36.4	Bq/Kg raw	± 12.6	Bq/Kg raw	55.9	Cs137	8.0	Bq/Kg raw
			Cs134	19.5	Bq/Kg raw	± 6.2	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Yuzu	Nakakabeya, Taira, Iwaki	Nov-12	Cs137	19.7	Bq/Kg raw	± 5.0	Bq/Kg raw	34.2	Cs137	7.0	Bq/Kg raw
			Cs134	14.5	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Kamikajiro, Onahama, Iwaki	Nov-12	Cs137	35.2	Bq/Kg raw	± 8.2	Bq/Kg raw	57.8	Cs137	7.0	Bq/Kg raw
			Cs134	22.6	Bq/Kg raw	± 5.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Kode, Onahama, Iwaki	Nov-12	Cs137	26.0	Bq/Kg raw	± 6.6	Bq/Kg raw	49.3	Cs137	7.0	Bq/Kg raw
			Cs134	23.3	Bq/Kg raw	± 5.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Tono, Iwaki	Nov-12	Cs137	23.0	Bq/Kg raw	± 6.0	Bq/Kg raw	42.2	Cs137	7.0	Bq/Kg raw
			Cs134	19.2	Bq/Kg raw	± 4.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Watanabe, Iwaki	Nov-12	Cs137	10.8	Bq/Kg raw	± 4.0	Bq/Kg raw	16.4	Cs137	7.0	Bq/Kg raw
			Cs134	5.6	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Yuzu	Numabe, Iwaki	Nov-12	Cs137	22.9 Bq/Kg raw	± 6.1 Bq/Kg raw	40.0	Cs137	7.0 Bq/Kg raw	
			Cs134	17.1 Bq/Kg raw	± 4.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Yuzu	Yamada, Iwaki	Nov-12	Cs137	14.8 Bq/Kg raw	± 4.3 Bq/Kg raw	21.6	Cs137	7.0 Bq/Kg raw	
			Cs134	6.8 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Ginkgo	Toriinish, Nishiki, Iwaki	Nov-12	Cs137	2.9 Bq/Kg raw	± 2.3 Bq/Kg raw	6.5	Cs137	8.0 Bq/Kg raw	
			Cs134	3.6 Bq/Kg raw	± 3.7 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Persimmon	Kamata, Fukushima	Nov-12	Cs137	13.3 Bq/Kg raw	± 4.6 Bq/Kg raw	27.8	Cs137	7.0 Bq/Kg raw	
			Cs134	14.5 Bq/Kg raw	± 4.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Ootama, Adachi	Nov-12	Cs137	8.9 Bq/Kg raw	± 3.1 Bq/Kg raw	11.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Naraha, Iwaki	Nov-12	Cs137	26.2 Bq/Kg raw	± 6.8 Bq/Kg raw	47.1	Cs137	7.0 Bq/Kg raw	
			Cs134	20.9 Bq/Kg raw	± 5.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Shimoogawa, Ogawa, Iwaki	Oct-12	Cs137	1.9 Bq/Kg raw	± 2.2 Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw	
			Cs134	1.4 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Shimoogoe, Taira, Iwaki	Nov-12	Cs137	2.0 Bq/Kg raw	± 2.6 Bq/Kg raw	3.9	Cs137	7.0 Bq/Kg raw	
			Cs134	1.9 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Kitakabeya, taira, Iwaki	Nov-12	Cs137	6.6 Bq/Kg raw	± 2.8 Bq/Kg raw	9.9	Cs137	7.0 Bq/Kg raw	
			Cs134	3.3 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Shimoyamaguchi, Taira, Iwaki	Nov-12	Cs137	7.6 Bq/Kg raw	± 2.7 Bq/Kg raw	11.8	Cs137	7.0 Bq/Kg raw	
			Cs134	4.2 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Shimokabeya, Iwaki	Nov-12	Cs137	8.0 Bq/Kg raw	± 3.2 Bq/Kg raw	10.7	Cs137	7.0 Bq/Kg raw	
			Cs134	2.7 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Kamikuramochi, Kasima, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw	
			Cs134	1.8 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Izumizaki, Taira, Iwaki	Nov-12	Cs137	10.4 Bq/Kg raw	± 3.9 Bq/Kg raw	14.7	Cs137	7.0 Bq/Kg raw	
			Cs134	4.3 Bq/Kg raw	± 2.8 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Shimokawa, Izumi, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Unknown	Unknown	Cs137	1.8 Bq/Kg raw	± 2.3 Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw	
			Cs134	1.8 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Persimmon	Watanabe, Iwaki	Nov-12	Cs137	3.7 Bq/Kg raw	± 2.5 Bq/Kg raw	6.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.8 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Hosoya, Yotsukura, Iwaki	Oct-12	Cs137	7.8 Bq/Kg raw	± 2.9 Bq/Kg raw	12.9	Cs137	7.0 Bq/Kg raw	
			Cs134	5.1 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Shimoogawa, Ogawa, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw	
			Cs134	3.2 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Ogawa, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw	
			Cs134	1.8 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Shimokanbeya, Iwaki	Nov-12	Cs137	13.0 Bq/Kg raw	± 4.4 Bq/Kg raw	25.7	Cs137	7.0 Bq/Kg raw	
			Cs134	12.7 Bq/Kg raw	± 3.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Kashima, Iwaki	Oct-12	Cs137	12.6 Bq/Kg raw	± 3.5 Bq/Kg raw	19.8	Cs137	7.0 Bq/Kg raw	
			Cs134	7.2 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Kubo, Kashima, Iwaki	Nov-12	Cs137	12.1 Bq/Kg raw	± 3.7 Bq/Kg raw	19.4	Cs137	7.0 Bq/Kg raw	
			Cs134	7.3 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Kiwi fruit	Yoneda, Kashima, Iwaki	Nov-12	Cs137	14.9 Bq/Kg raw	± 4.4 Bq/Kg raw	17.7	Cs137	7.0 Bq/Kg raw	
			Cs134	2.8 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Kiwi fruit	Kamikuramochi, Kasima, Iwaki	Nov-12	Cs137	9.6 Bq/Kg raw	± 3.3 Bq/Kg raw	15.4	Cs137	7.0 Bq/Kg raw	
			Cs134	5.8 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Gold kiwi	Kashima, Iwaki	Nov-12	Cs137	1.3 Bq/Kg raw	± 2.3 Bq/Kg raw	3.1	Cs137	7.0 Bq/Kg raw	
			Cs134	1.8 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Apple kiwi	Ogawa, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw	
			Cs134	4.1 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Chestnut	Numanouchi, Iwaki	Oct-12	Cs137	22.6 Bq/Kg raw	± 5.6 Bq/Kg raw	36.1	Cs137	7.0 Bq/Kg raw	
			Cs134	13.5 Bq/Kg raw	± 3.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Chestnut	Tabito, Iwaki	Oct-12	Cs137	32.8 Bq/Kg raw	± 9.7 Bq/Kg raw	44.4	Cs137	8.0 Bq/Kg raw	
			Cs134	11.6 Bq/Kg raw	± 5.5 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Apple	Fukushima	Nov-12	Cs137	4.3 Bq/Kg raw	± 2.4 Bq/Kg raw	8.9	Cs137	7.0 Bq/Kg raw	
			Cs134	4.6 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Pear	Niigata, Niigata	Sep-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Fig	Kitashirado, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.1	Cs137	7.0 Bq/Kg raw	
			Cs134	1.1 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Sea cucumber	Onahama, Iwaki	Nov-12	Cs137	1.5 Bq/Kg raw	± 2.0 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Salmon	Tomioka, Futaba	Nov-12	Cs137	1.9 Bq/Kg raw	± 2.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Eel (boiled)	Unknown	Unknown	Cs137	162 Bq/Kg raw	± 32.0 Bq/Kg raw	269	Cs137	7.0 Bq/Kg raw	
			Cs134	107 Bq/Kg raw	± 21.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Pheasant meat	Miwa, Iwaki	Nov-12	Cs137	19.1 Bq/Kg raw	± 5.2 Bq/Kg raw	34.8	Cs137	7.0 Bq/Kg raw	
			Cs134	15.7 Bq/Kg raw	± 4.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Rice Grasshopper (dried)	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	6.7	Cs137	7.0 Bq/Kg raw	
			Cs134	6.7 Bq/Kg raw	± 8.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Rice Grasshopper (boiled)	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.2	Cs137	7.0 Bq/Kg raw	
			Cs134	3.2 Bq/Kg raw	± 3.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Copper pheasant	Miwa, Iwaki	Nov-12	Cs137	43.2 Bq/Kg raw	± Bq/Kg raw	73.4	Cs137	7.0 Bq/Kg raw	
			Cs134	30.2 Bq/Kg raw	± 6.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Milk	Hokkaido	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	5.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Milk	Hyogo	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	5.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
School lunch	Unknown	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw	
			Cs134	1.4 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
School lunch	Unknown	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
School lunch	Unknown	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Fig jam	Shimoogawa, Ogawa, Iwaki	Oct-12	Cs137	1.4 Bq/Kg raw	± 2.1 Bq/Kg raw	2.6	Cs137	7.0 Bq/Kg raw	
			Cs134	1.2 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Tomato jam	Shimoogawa, Ogawa, Iwaki	Sep-12	Cs137	2.3 Bq/Kg raw	± 2.9 Bq/Kg raw	2.3	Cs137	8.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	8.0 Bq/Kg raw	
Honey	Kamitakaku, Taira, Iwaki	Unknown	Cs137	108 Bq/Kg raw	± 22.0 Bq/Kg raw	185	Cs137	7.0 Bq/Kg raw	
			Cs134	76.5 Bq/Kg raw	± 15.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Tap water	Kitsunozuka, Yotsukura, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	6.0
Bee's nest	Kamitakaku, Taira, Iwaki	Unknown	Cs137	176.0	Bq/Kg raw	± 35.0	293	Cs137	7.0	Bq/Kg raw
			Cs134	117.0	Bq/Kg raw	± 23.0		Bq/Kg raw	Cs134	6.0
Raw wood (Shiitake)	Komoda, Kashima, Iwaki	Nov-12	Cs137	346	Bq/Kg dry	± 69.0	572	Cs137	7.0	Bq/Kg dry
			Cs134	226	Bq/Kg dry	± 45.0		Bq/Kg dry	Cs134	6.0
Pottery	Ohisa, Iwaki	Jul-12	Cs137	27.0	Bq/Kg dry	± 8.0	89.0	Cs137	7.0	Bq/Kg dry
			Cs134	62.0	Bq/Kg dry	± 13.0		Bq/Kg dry	Cs134	6.0
Pottery	Ohisa, Iwaki	Jul-12	Cs137	47.0	Bq/Kg dry	± 13.0	57.0	Cs137	8.0	Bq/Kg dry
			Cs134	10.0	Bq/Kg dry	± 7.0		Bq/Kg dry	Cs134	8.0
Thatch	Honya, Izumi, Iwaki	Oct-12	Cs137	12.0	Bq/Kg dry	± 10.0	42.0	Cs137	7.0	Bq/Kg dry
			Cs134	30.0	Bq/Kg dry	± 10.0		Bq/Kg dry	Cs134	6.0
Bamboo	Honcho, Izumi, Iwaki	Oct-12	Cs137	43.0	Bq/Kg dry	± 12.0	67.0	Cs137	7.0	Bq/Kg dry
			Cs134	24.0	Bq/Kg dry	± 8.0		Bq/Kg dry	Cs134	6.0
Ash (bamboo, thatch)	Honya, Izumi, Iwaki	Nov-12	Cs137	972	Bq/Kg dry	± 194	1,643	Cs137	7.0	Bq/Kg dry
			Cs134	671	Bq/Kg dry	± 134		Bq/Kg dry	Cs134	6.0
Rice field soil	Honya, Izumi, Iwaki	Oct-12	Cs137	237	Bq/Kg dry	± 29.0	358	Cs137	8.0	Bq/Kg dry
			Cs134	121	Bq/Kg dry	± 16.0		Bq/Kg dry	Cs134	8.0