



Radiation Measurement Results of 231 items in December 2012



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Kitaibaraki, Ibaraki	Oct-12	Cs137	9.9 Bq/Kg raw	± 3.4 Bq/Kg raw	12.5	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Nakago, Kitaibaraki	Oct-12	Cs137	5.7 Bq/Kg raw	± 2.5 Bq/Kg raw	5.7	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Sekimoto, Kitaibaraki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Taira, Iwaki	Oct-11	Cs137	11.2 Bq/Kg raw	± 3.7 Bq/Kg raw	16.7	Cs137	7.0 Bq/Kg raw	
			Cs134	5.5 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	4.7 Bq/Kg raw	± 2.9 Bq/Kg raw	10.4	Cs137	7.0 Bq/Kg raw	
			Cs134	5.7 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	7.8 Bq/Kg raw	± 3.1 Bq/Kg raw	11.2	Cs137	7.0 Bq/Kg raw	
			Cs134	3.4 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	6.0 Bq/Kg raw	± 2.1 Bq/Kg raw	8.6	Cs137	7.0 Bq/Kg raw	
			Cs134	2.6 Bq/Kg raw	± 1.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Nakahirakubo, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw	
			Cs134	1.4 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Tono, Iwaki	Oct-12	Cs137	1.5 Bq/Kg raw	± 2.3 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	11.0 Bq/Kg raw	± 3.9 Bq/Kg raw	20.4	Cs137	7.0 Bq/Kg raw	
			Cs134	9.4 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	14.0 Bq/Kg raw	± 4.1 Bq/Kg raw	23.0	Cs137	7.0 Bq/Kg raw	
			Cs134	9.0 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	15.8 Bq/Kg raw	± 4.3 Bq/Kg raw	22.5	Cs137	7.0 Bq/Kg raw	
			Cs134	6.7 Bq/Kg raw	± 2.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	12.3 Bq/Kg raw	± 3.9 Bq/Kg raw	21.0	Cs137	7.0 Bq/Kg raw	
			Cs134	8.7 Bq/Kg raw	± 2.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	16.3 Bq/Kg raw	± 4.6 Bq/Kg raw	26.8	Cs137	7.0 Bq/Kg raw	
			Cs134	10.5 Bq/Kg raw	± 3.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	17.1	Bq/Kg raw	± 4.5	Bq/Kg raw	24.0	Cs137	7.0	Bq/Kg raw
			Cs134	6.9	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	12.8	Bq/Kg raw	± 4.0	Bq/Kg raw	24.7	Cs137	7.0	Bq/Kg raw
			Cs134	11.9	Bq/Kg raw	± 3.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	14.8	Bq/Kg raw	± 4.3	Bq/Kg raw	25.3	Cs137	7.0	Bq/Kg raw
			Cs134	10.5	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	9.4	Bq/Kg raw	± 3.6	Bq/Kg raw	19.2	Cs137	7.0	Bq/Kg raw
			Cs134	9.8	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown rice	Okaona, Onahama, Iwaki	Oct-12	Cs137	9.2	Bq/Kg raw	± 3.6	Bq/Kg raw	21.0	Cs137	7.0	Bq/Kg raw
			Cs134	11.8	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Brown rice	Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Hokkaido	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Hiroshima	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Yonezawa, Yamagata	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Miyagi	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Akita	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ibaraki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ibaraki	Oct-11	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Tadami, Minamiaizu	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Fukagawa, Date	Oct-12	Cs137	1.8	Bq/Kg raw	± 2.1	Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Funahiki, Tamura	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ono, Tamura	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Ogawa, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-12	Cs137	1.6	Bq/Kg raw	± 2.4	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Yotsukura, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nagatomo, Yotsukura, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Kamitakaku, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
White rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	1.7 Bq/Kg raw	± 2.3 Bq/Kg raw	3.6	Cs137	7.0 Bq/Kg raw		
			Cs134	1.9 Bq/Kg raw	± 2.1 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Shimotakaku, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.7	Cs137	7.0 Bq/Kg raw		
			Cs134	1.7 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Nakahirakubo, Taira, Iwaki	Oct-12	Cs137	2.0 Bq/Kg raw	± 2.5 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Nakahirakubo, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Shimoogoe, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Kamioogoe, Taira, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw		
			Cs134	1.6 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Makunouchi, Taira, Iwaki	Oct-12	Cs137	1.4 Bq/Kg raw	± 2.1 Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Izumisaki, Taira, Iwaki	Unknown	Cs137	1.6 Bq/Kg raw	± 1.9 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Shimokajiro, Onahama, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw		
			Cs134	1.5 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.0	Cs137	7.0 Bq/Kg raw		
			Cs134	1.0 Bq/Kg raw	± 1.5 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Watanabe, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Yamada, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Yamada, Iwaki	Oct-12	Cs137	2.0 Bq/Kg raw	± 2.4 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Yamada, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw		
			Cs134	1.6 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Yamada, Iwaki	Oct-12	Cs137	1.4 Bq/Kg raw	± 2.3 Bq/Kg raw	1.4	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
White rice	Nakoso, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Sticky rice (brown rice)	Yamadama, Iwaki	Sep-12	Cs137	3.3 Bq/Kg raw	± 1.9 Bq/Kg raw	3.3	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Sticky rice	Sekimoto, Kitaibaraki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw		
Sticky rice	Nihonmatsu	Oct-12	Cs137	7.2 Bq/Kg raw	± 3.2 Bq/Kg raw	13.5	Cs137	7.0 Bq/Kg raw		
			Cs134	6.3 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Sticky rice	Kamioogoe, Taira, Iwaki	Oct-12	Cs137	1.2 Bq/Kg raw	± 2.0 Bq/Kg raw	1.2	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Sticky rice	Tono, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Sticky rice	Yamada, Iwaki	Unknown	Cs137	1.6 Bq/Kg raw	± 2.2 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Sticky rice	Numabe, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Nakoso, Iwaki	Oct-12	Cs137	1.7	Bq/Kg raw	± 2.3	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Fried rice	Miwa, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Fried rice	Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Red Japanese radish	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yotsukura, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yoshima, Iwaki	Dec-12	Cs137	2.8	Bq/Kg raw	± 3.5	Bq/Kg raw	6.3	Cs137	7.0	Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Shimoogoe, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yoshinoya, Taira, Iwaki	Dec-12	Cs137	1.7	Bq/Kg raw	± 2.3	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Hashirikuma, Kashima, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Shimokajiro, Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.6	Cs137	7.0	Bq/Kg raw
			Cs134	3.6	Bq/Kg raw	± 2.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Shimoyunagaya, Joban, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Misawa, Joban, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Tono, Iwaki	Dec-12	Cs137	1.4	Bq/Kg raw	± 2.1	Bq/Kg raw	2.9	Cs137	7.0	Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yamada, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yamada, Iwaki	Nov-12	Cs137	2.8	Bq/Kg raw	± 3.6	Bq/Kg raw	2.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish leaf	Yoshima, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish leaf	Kashima, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137	7.0	Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 3.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Ogawa, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Taro	Yamada, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yam	Fukushima	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Jerusalem artichoke	Gunma	2012年	Cs137	2.8	Bq/Kg raw	± 3.0	Bq/Kg raw	2.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Yotsukura, Iwaki	Jun-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Potato	Shimohirakubo, Iwaki	Jul-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweet potato	Yamada, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Burdock	Akai, Taira, Iwaki	Dec-12	Cs137	3.3	Bq/Kg raw	± 3.1	Bq/Kg raw	3.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Carrot	Shimoyunagaya, Joban, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Carrot	Inoue, Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Carrot	Namesawa, Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Carrot (with skin)	Fukushima	Dec-12	Cs137	5.6	Bq/Kg raw	± 9.7	Bq/Kg raw	6.3	Cs137	8.0	Bq/Kg raw
			Cs134	0.7	Bq/Kg raw	± 7.4	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Lotus root	Ibaraki	Nov-12	Cs137	7.8	Bq/Kg raw	± 2.7	Bq/Kg raw	10.8	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yacon	Nakahirakubo, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.9	Cs137	7.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yacon	Nakahirakubo, Taira, Iwaki	Dec-12	Cs137	3.3	Bq/Kg raw	± 3.0	Bq/Kg raw	3.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Higashifukaniida, Shirakawa	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Tanakura, Higashishirakawa	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Hosoya, Yotsukura, Iwaki	Dec-12	Cs137	1.9	Bq/Kg raw	± 3.0	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Shimoyoshima, Yoshima, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Kamiyoshima, Yoshima, Iwaki	Dec-12	Cs137	2.5	Bq/Kg raw	± 2.6	Bq/Kg raw	2.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Kamioogoe, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.9	Cs137	7.0	Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Shimoogoe, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Yoshinoya, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Takaku, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Kashima, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Kashima, Iwaki	Nov-12	Cs137	3.8	Bq/Kg raw	± 4.6	Bq/Kg raw	10.1	Cs137	7.0	Bq/Kg raw
			Cs134	6.3	Bq/Kg raw	± 3.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Chinese cabbage	Hashirikuma, Kashima, Iwaki	Dec-12	Cs137	6.7 Bq/Kg raw	± 3.3 Bq/Kg raw	9.0	Cs137	7.0 Bq/Kg raw		
			Cs134	2.3 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Chinese cabbage	Kamikajiro, Onahama, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Chinese cabbage	Shimokajiro, Onahama, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Chinese cabbage	Misawa, Joban, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Chinese cabbage	Obama, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Chinese cabbage	Yamada, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	6.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.0 Bq/Kg raw		
Chinese cabbage	Yamada, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw		
			Cs134	1.9 Bq/Kg raw	± 2.4 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Fukushima	Dec-12	Cs137	3.1 Bq/Kg raw	± 6.8 Bq/Kg raw	3.1	Cs137	8.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.3 Bq/Kg raw		Cs134	8.0 Bq/Kg raw		
Leek	Higashifukaniida, Shirakawa	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Tanakura, Higashishirakawa	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Hosoya, Yotsukura, Iwaki	Dec-12	Cs137	1.6 Bq/Kg raw	± 2.4 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Akai, Taira, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Shimoogoe, Taira, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw		
			Cs134	1.9 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Takaku, Taira, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Miya, Uchigo, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Kashima, Iwaki	Dec-12	Cs137	5.1 Bq/Kg raw	± 3.3 Bq/Kg raw	5.1	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Kamikajiro, Onahama, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Obama, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Yamada, Iwaki	Nov-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Leek	Yamada, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Purple leek	Shimohirakubo, Taira, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw		
			Cs134	1.5 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Tomato	Nagasaki, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.8	Cs137	7.0 Bq/Kg raw		
			Cs134	1.8 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		
Cabbage	Kamikajiro, Onahama, Iwaki	Dec-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.0 Bq/Kg raw		

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Cabbage	Onahama, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Onahama, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.8	Cs137	7.0	Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 4.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Spinach	Kashima, Iwaki	Dec-12	Cs137	4.5	Bq/Kg raw	± 3.4	Bq/Kg raw	7.8	Cs137	7.0	Bq/Kg raw
			Cs134	3.3	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Bok choy	Shimohirakubo, Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Bok choy	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese chive	Taira, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Greens	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	6.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Wild sesame	Ohisa, Iwaki	Nov-12	Cs137	9.6	Bq/Kg raw	± 5.5	Bq/Kg raw	26.9	Cs137	7.0	Bq/Kg raw
			Cs134	17.3	Bq/Kg raw	± 5.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Wild sesame leaf and stem	Ohisa, Iwaki	Nov-12	Cs137	50.0	Bq/Kg raw	± 32.3	Bq/Kg raw	120	Cs137	7.0	Bq/Kg raw
			Cs134	69.7	Bq/Kg raw	± 30.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shungiku	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese mustard spinach	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	4.6	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.1	Bq/Kg raw
Sword bean	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.1	Cs137	7.0	Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Azuki bean	Tabito, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	2.1	Cs137	7.0	Bq/Kg raw
			Cs134	2.1	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Soybean	Kamata, Fukushima	Nov-12	Cs137	2.0	Bq/Kg raw	± 2.6	Bq/Kg raw	3.1	Cs137	7.0	Bq/Kg raw
			Cs134	1.1	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Soybean	Uchimachi, Uchigo, Iwaki	Oct-11	Cs137	16.3	Bq/Kg raw	± 5.0	Bq/Kg raw	27.9	Cs137	7.0	Bq/Kg raw
			Cs134	11.6	Bq/Kg raw	± 4.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Soybean	Uchimachi, Uchigo, Iwaki	Oct-12	Cs137	17.0	Bq/Kg raw	± 4.8	Bq/Kg raw	27.6	Cs137	7.0	Bq/Kg raw
			Cs134	10.6	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Fuyuyamatake mushroom	Kouya, Uchigo, Iwaki	Dec-12	Cs137	154	Bq/Kg raw	± 32.0	Bq/Kg raw	256	Cs137	7.0	Bq/Kg raw
			Cs134	102	Bq/Kg raw	± 21.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Nameko mushroom	Watanabe, Iwaki	Dec-12	Cs137	4.2	Bq/Kg raw	± 2.5	Bq/Kg raw	9.9	Cs137	7.0	Bq/Kg raw
			Cs134	5.7	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Shibahara, Ogawa, Iwaki	Dec-12	Cs137	189	Bq/Kg raw	± 29.0	Bq/Kg raw	289	Cs137	8.0	Bq/Kg raw
			Cs134	99.9	Bq/Kg raw	± 16.5	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shiitake mushroom	Yamaguchi, Taira, Iwaki	Nov-12	Cs137	1350	Bq/Kg raw	± 134	Bq/Kg raw	2,005	Cs137	8.0	Bq/Kg raw
			Cs134	655	Bq/Kg raw	± 66.8	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Shiitake mushroom	Kamikajiro, Onahama, Iwaki	Nov-12	Cs137	280	Bq/Kg raw	± 56.0	Bq/Kg raw	483	Cs137	7.0	Bq/Kg raw
			Cs134	203	Bq/Kg raw	± 41.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Shiitake mushroom	Watanabe, Iwaki	Dec-12	Cs137	51.3	Bq/Kg raw	± 12.2	Bq/Kg raw	86.1	Cs137	7.0	Bq/Kg raw
			Cs134	34.8	Bq/Kg raw	± 8.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Amitake mushroom (washed)	Saiso, Miwa, Iwaki	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Yuzu	Ohisa, Iwaki	Dec-12	Cs137	89.1	Bq/Kg raw	± 18.6	150	Cs137	7.0	Bq/Kg raw
			Cs134	61.3	Bq/Kg raw	± 12.5		Cs134	6.0	Bq/Kg raw
Yuzu	Shibahara, Ogawa, Iwaki	Dec-12	Cs137	24.1	Bq/Kg raw	± 5.8	39.2	Cs137	7.0	Bq/Kg raw
			Cs134	15.1	Bq/Kg raw	± 3.8		Cs134	6.0	Bq/Kg raw
Yuzu	Yotsukura, Iwaki	Nov-12	Cs137	54.4	Bq/Kg raw	± 12.5	91.5	Cs137	7.0	Bq/Kg raw
			Cs134	37.1	Bq/Kg raw	± 8.5		Cs134	6.0	Bq/Kg raw
Yuzu	Taira, Iwaki	Dec-12	Cs137	14.5	Bq/Kg raw	± 4.2	21.8	Cs137	7.0	Bq/Kg raw
			Cs134	7.3	Bq/Kg raw	± 2.7		Cs134	6.0	Bq/Kg raw
Yuzu	Hashirikuma, Kashima, Iwaki	Dec-12	Cs137	26.7	Bq/Kg raw	± 6.6	48.5	Cs137	7.0	Bq/Kg raw
			Cs134	21.8	Bq/Kg raw	± 5.1		Cs134	6.0	Bq/Kg raw
Yuzu	Nagasaki, Iwaki	Dec-12	Cs137	74.9	Bq/Kg raw	± 16.2	118	Cs137	7.0	Bq/Kg raw
			Cs134	42.8	Bq/Kg raw	± 9.5		Cs134	6.0	Bq/Kg raw
Yuzu	Ohara, Onahama, Iwaki	Dec-12	Cs137	16.8	Bq/Kg raw	± 4.8	29.7	Cs137	7.0	Bq/Kg raw
			Cs134	12.9	Bq/Kg raw	± 3.7		Cs134	6.0	Bq/Kg raw
Yuzu	Shimokajiro, Onahama, Iwaki	Dec-12	Cs137	18.4	Bq/Kg raw	± 5.1	30.6	Cs137	7.0	Bq/Kg raw
			Cs134	12.2	Bq/Kg raw	± 3.6		Cs134	6.0	Bq/Kg raw
Yuzu	Tono, Iwaki	Dec-12	Cs137	24.9	Bq/Kg raw	± 6.5	40.9	Cs137	7.0	Bq/Kg raw
			Cs134	16.0	Bq/Kg raw	± 4.5		Cs134	6.0	Bq/Kg raw
Yuzu	Yamada, Iwaki	Nov-12	Cs137	7.6	Bq/Kg raw	± 4.0	18.2	Cs137	7.0	Bq/Kg raw
			Cs134	10.6	Bq/Kg raw	± 3.7		Cs134	6.0	Bq/Kg raw
Yuzu	Nakoso, Iwaki	Nov-12	Cs137	24.4	Bq/Kg raw	± 6.0	45.8	Cs137	7.0	Bq/Kg raw
			Cs134	21.4	Bq/Kg raw	± 5.0		Cs134	6.0	Bq/Kg raw
Mandarin orange	Kamiyoshima, Yoshima, Iwaki	Dec-12	Cs137	7.6	Bq/Kg raw	± 3.1	13.2	Cs137	7.0	Bq/Kg raw
			Cs134	5.6	Bq/Kg raw	± 2.4		Cs134	6.0	Bq/Kg raw
Mandarin orange	Shimotakaku, Taira, Iwaki	Dec-12	Cs137	9.5	Bq/Kg raw	± 7.3	17.6	Cs137	8.0	Bq/Kg raw
			Cs134	8.1	Bq/Kg raw	± 4.6		Cs134	8.0	Bq/Kg raw
Mandarin orange	Wakabadai, Iwaki	Nov-12	Cs137	7.8	Bq/Kg raw	± 3.3	15.5	Cs137	7.0	Bq/Kg raw
			Cs134	7.7	Bq/Kg raw	± 2.9		Cs134	6.0	Bq/Kg raw
Mandarin orange	Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	4.9	Cs137	7.0	Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.4		Cs134	6.0	Bq/Kg raw
Mandarin orange	Ohara, Onahama, Iwaki	Unknown	Cs137	5.5	Bq/Kg raw	± 2.6	9.5	Cs137	7.0	Bq/Kg raw
			Cs134	4.0	Bq/Kg raw	± 2.0		Cs134	6.0	Bq/Kg raw
Mandarin orange	Ohara, Onahama, Iwaki	Nov-12	Cs137	9.1	Bq/Kg raw	± 3.1	13.5	Cs137	7.0	Bq/Kg raw
			Cs134	4.4	Bq/Kg raw	± 2.2		Cs134	6.0	Bq/Kg raw
Mandarin orange	Ohara, Onahama, Iwaki	Dec-12	Cs137	1.3	Bq/Kg raw	± 1.6	2.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.4		Cs134	6.0	Bq/Kg raw
Mandarin orange	Onahama, Iwaki	Dec-12	Cs137	5.2	Bq/Kg raw	± 2.6	10.1	Cs137	7.0	Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.2		Cs134	6.0	Bq/Kg raw
Mandarin orange	Tamagawa, Onahama, Iwaki	Dec-12	Cs137	6.2	Bq/Kg raw	± 3.3	11.1	Cs137	7.0	Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.6		Cs134	6.0	Bq/Kg raw
Mandarin orange	Shonandai, Iwaki	Oct-12	Cs137	8.7	Bq/Kg raw	± 4.6	8.9	Cs137	8.0	Bq/Kg raw
			Cs134	0.2	Bq/Kg raw	± 2.1		Cs134	8.0	Bq/Kg raw
Mandarin orange	Izumigaoka, Iwaki	Dec-12	Cs137	6.9	Bq/Kg raw	± 2.5	6.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	6.0	Bq/Kg raw
Mandarin orange	Numabe, Iwaki	Nov-12	Cs137	11.3	Bq/Kg raw	± 3.5	17.1	Cs137	7.0	Bq/Kg raw
			Cs134	5.8	Bq/Kg raw	± 2.4		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Natsumikan	Numabe, Iwaki	Dec-12	Cs137	7.9	Bq/Kg raw	± 3.3	Bq/Kg raw	14.7	Cs137	7.0	Bq/Kg raw
			Cs134	6.8	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Apple	Nagano	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.7	Cs137	7.0	Bq/Kg raw
			Cs134	1.7	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Apple	Fukushima	Dec-12	Cs137	13.2	Bq/Kg raw	± 4.0	Bq/Kg raw	19.2	Cs137	7.0	Bq/Kg raw
			Cs134	6.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Apple	Fukushima	Dec-12	Cs137	4.7	Bq/Kg raw	± 6.3	Bq/Kg raw	8.1	Cs137	8.0	Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 4.0	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Apple	Ishikawa	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Apple	Date	Nov-12	Cs137	6.6	Bq/Kg raw	± 5.2	Bq/Kg raw	9.5	Cs137	8.0	Bq/Kg raw
			Cs134	2.9	Bq/Kg raw	± 4.2	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Apple	Kamihobara, Date	Nov-12	Cs137	10.5	Bq/Kg raw	± 4.3	Bq/Kg raw	14.4	Cs137	7.0	Bq/Kg raw
			Cs134	3.9	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Apple	Aizuwakamatsu	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Apple	Onahama, Iwaki	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Ohisa, Iwaki	Dec-12	Cs137	14.0	Bq/Kg raw	± 4.0	Bq/Kg raw	21.0	Cs137	7.0	Bq/Kg raw
			Cs134	7.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Yotsukura, Iwaki	Dec-12	Cs137	8.1	Bq/Kg raw	± 3.0	Bq/Kg raw	10.4	Cs137	7.0	Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Kamiyoshima, Yoshima, Iwaki	Nov-12	Cs137	4.1	Bq/Kg raw	± 2.2	Bq/Kg raw	4.1	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Shio, Taira, Iwaki	Dec-12	Cs137	8.3	Bq/Kg raw	± 3.1	Bq/Kg raw	12.4	Cs137	7.0	Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Hashirikuma, Kashima, Iwaki	Dec-12	Cs137	4.3	Bq/Kg raw	± 2.5	Bq/Kg raw	8.6	Cs137	7.0	Bq/Kg raw
			Cs134	4.3	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Kashima, Iwaki	Nov-12	Cs137	8.3	Bq/Kg raw	± 2.9	Bq/Kg raw	13.8	Cs137	7.0	Bq/Kg raw
			Cs134	5.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Shonandai, Iwaki	Oct-12	Cs137	6.5	Bq/Kg raw	± 5.9	Bq/Kg raw	8.1	Cs137	8.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 4.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Kiwi fruit	Tono, Iwaki	Dec-12	Cs137	8.9	Bq/Kg raw	± 3.1	Bq/Kg raw	13.1	Cs137	7.0	Bq/Kg raw
			Cs134	4.2	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Dried persimmon	Shimoyamaguchi, Taira, Iwaki	Dec-12	Cs137	23.9	Bq/Kg raw	± 5.7	Bq/Kg raw	36.9	Cs137	7.0	Bq/Kg raw
			Cs134	13.0	Bq/Kg raw	± 3.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Dried persimmon	Watanabe, Iwaki	Dec-12	Cs137	8.2	Bq/Kg raw	± 2.6	Bq/Kg raw	10.1	Cs137	7.0	Bq/Kg raw
			Cs134	1.9	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Persimmon	Yotsukura, Iwaki	Nov-12	Cs137	6.5	Bq/Kg raw	± 3.0	Bq/Kg raw	16.5	Cs137	7.0	Bq/Kg raw
			Cs134	10.0	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Persimmon	Chuodai, Iwaki	Dec-12	Cs137	3.2	Bq/Kg raw	± 2.3	Bq/Kg raw	6.6	Cs137	7.0	Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Akebi seed	Soeno, Iwaki	Nov-12	Cs137	3.4	Bq/Kg raw	± 2.6	Bq/Kg raw	8.3	Cs137	7.0	Bq/Kg raw
			Cs134	4.9	Bq/Kg raw	± 2.3	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Akebi pulp	Soeno, Iwaki	Nov-12	Cs137	4.7	Bq/Kg raw	± 2.4	Bq/Kg raw	4.7	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Akebi skin	Soeno, Iwaki	Nov-12	Cs137	7.4	Bq/Kg raw	± 2.9	Bq/Kg raw	11.5	Cs137	7.0	Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Blueberry	Kamiyoshima, Yoshima, Iwaki	Dec-12	Cs137	7.0	Bq/Kg raw	± 2.6	Bq/Kg raw	9.7	Cs137	7.0	Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Fig	Nagasaki, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Lemon	Izumigaoka, Iwaki	Dec-12	Cs137	4.6	Bq/Kg raw	± 2.5	Bq/Kg raw	6.9	Cs137	7.0	Bq/Kg raw
			Cs134	2.3	Bq/Kg raw	± 2.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pheasant meat	Kajiro, Iwaki	Nov-12	Cs137	17.3	Bq/Kg raw	± 4.5	Bq/Kg raw	25.3	Cs137	7.0	Bq/Kg raw
			Cs134	8.0	Bq/Kg raw	± 2.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Wild boa meat	Between Fukushima and Niigata	Oct-12	Cs137	361	Bq/Kg raw	± 72.0	Bq/Kg raw	594	Cs137	7.0	Bq/Kg raw
			Cs134	233	Bq/Kg raw	± 47.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Wild boa meat (female)	Tono, Iwaki	Nov-12	Cs137	92.5	Bq/Kg raw	± 19.1	Bq/Kg raw	161	Cs137	7.0	Bq/Kg raw
			Cs134	68.8	Bq/Kg raw	± 13.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Wild boa meat (male)	Uchigo, Iwaki	Nov-12	Cs137	147	Bq/Kg raw	± 29.0	Bq/Kg raw	242	Cs137	7.0	Bq/Kg raw
			Cs134	94.5	Bq/Kg raw	± 18.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Wild boa meat (young female)	Tono, Iwaki	Nov-12	Cs137	118	Bq/Kg raw	± 24.0	Bq/Kg raw	197	Cs137	7.0	Bq/Kg raw
			Cs134	78.9	Bq/Kg raw	± 15.8	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Copper pheasant	Furudono, Ishikawa	Dec-12	Cs137	24.5	Bq/Kg raw	± 8.3	Bq/Kg raw	32.3	Cs137	8.0	Bq/Kg raw
			Cs134	7.8	Bq/Kg raw	± 4.8	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Copper pheasant	Miwa, Iwaki	Dec-12	Cs137	125	Bq/Kg raw	± 19.4	Bq/Kg raw	192	Cs137	8.0	Bq/Kg raw
			Cs134	67.2	Bq/Kg raw	± 10.3	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Mixed minced meat	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sweetfish	Unknown	Unknown	Cs137	5.9	Bq/Kg raw	± 2.9	Bq/Kg raw	8.4	Cs137	8.0	Bq/Kg raw
			Cs134	2.5	Bq/Kg raw	± 1.2	Bq/Kg raw		Cs134	8.0	Bq/Kg raw
Abalone	Ohtsu, Kitaibaraki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	4.6	Cs137	7.0	Bq/Kg raw
			Cs134	4.6	Bq/Kg raw	± 2.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Vienna sausage	Unknown	Unknown	Cs137	1.3	Bq/Kg raw	± 1.8	Bq/Kg raw	1.3	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
School lunch	Unknown	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
School lunch	Unknown	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
School lunch	Unknown	Dec-12	Cs137	1.4	Bq/Kg raw	± 1.9	Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Miso paste	Unknown	Unknown	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Well water	Ogawa, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	5.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	3.0	Bq/Kg raw
Ground water	Furudono, Ishikawa	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	1.5	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	1.4	Bq/Kg raw
Spring water	Furudono, Ishikawa	Nov-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	5.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	3.0	Bq/Kg raw
Nikko maple	Watanabe, Iwaki	Dec-12	Cs137	181	Bq/Kg raw	± 38.0	Bq/Kg raw	316	Cs137	7.0	Bq/Kg raw
			Cs134	135	Bq/Kg raw	± 28.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Saw dust	Akai, Taira, Iwaki	Dec-12	Cs137	54.0 <small>Bq/Kg dry</small>	± 13.0 <small>Bq/Kg dry</small>	97.0	Cs137	7.0 <small>Bq/Kg dry</small>
			Cs134	43.0 <small>Bq/Kg dry</small>	± 10.0 <small>Bq/Kg dry</small>		Cs134	6.0 <small>Bq/Kg dry</small>
Charcoal	Hirata, Ishikawa	Jun-11	Cs137	30.0 <small>Bq/Kg dry</small>	± 10.0 <small>Bq/Kg dry</small>	50.0	Cs137	7.0 <small>Bq/Kg dry</small>
			Cs134	20.0 <small>Bq/Kg dry</small>	± 7.0 <small>Bq/Kg dry</small>		Cs134	6.0 <small>Bq/Kg dry</small>
Ash	Shimoogoe, Taira, Iwaki	Dec-12	Cs137	4900 <small>Bq/Kg dry</small>	± 980 <small>Bq/Kg dry</small>	8,230	Cs137	7.0 <small>Bq/Kg dry</small>
			Cs134	3330 <small>Bq/Kg dry</small>	± 670 <small>Bq/Kg dry</small>		Cs134	6.0 <small>Bq/Kg dry</small>
Leaf mold	Onahama, Iwaki	Dec-12	Cs137	39.0 <small>Bq/Kg dry</small>	± 23.0 <small>Bq/Kg dry</small>	50.0	Cs137	8.0 <small>Bq/Kg dry</small>
			Cs134	11.0 <small>Bq/Kg dry</small>	± 9.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>
Compost	Onahama, Iwaki	Dec-12	Cs137	39.1 <small>Bq/Kg dry</small>	± 23.0 <small>Bq/Kg dry</small>	51.0	Cs137	9.9 <small>Bq/Kg dry</small>
			Cs134	11.9 <small>Bq/Kg dry</small>	± 9.1 <small>Bq/Kg dry</small>		Cs134	11.8 <small>Bq/Kg dry</small>
Field soil	Meiji Danchi, Iwaki	Dec-12	Cs137	199 <small>Bq/Kg dry</small>	± 29.0 <small>Bq/Kg dry</small>	303	Cs137	8.0 <small>Bq/Kg dry</small>
			Cs134	104 <small>Bq/Kg dry</small>	± 16.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>
Field soil	Tono, Iwaki	Dec-12	Cs137	47.0 <small>Bq/Kg dry</small>	± 13.0 <small>Bq/Kg dry</small>	71.0	Cs137	8.0 <small>Bq/Kg dry</small>
			Cs134	24.0 <small>Bq/Kg dry</small>	± 8.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>