



Radiation Measurement Results of 74 items in January 2013



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Brown rice	Kurihara, Miyagi	Oct-10	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
Brown rice	Yotsukura, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.3	Cs137	7.0 Bq/Kg raw		
			Cs134	1.3 Bq/Kg raw	± 2.2 Bq/Kg raw			Cs134		
Brown rice	Manome, taira, Iwaki	Oct-12	Cs137	1.3 Bq/Kg raw	± 2.2 Bq/Kg raw	1.3	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
Brown rice	Watanabemachibora, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.9	Cs137	7.0 Bq/Kg raw		
			Cs134	1.9 Bq/Kg raw	± 2.3 Bq/Kg raw			Cs134		
Brown rice	Watanabemachibora, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.3	Cs137	7.0 Bq/Kg raw		
			Cs134	1.3 Bq/Kg raw	± 2.2 Bq/Kg raw			Cs134		
Brown rice	Inoue, Yamada, Iwaki	Oct-12	Cs137	1.7 Bq/Kg raw	± 2.2 Bq/Kg raw	1.7	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Minamiuonuma, Niigata	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Niigata	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.1	Cs137	7.0 Bq/Kg raw		
			Cs134	2.1 Bq/Kg raw	± 2.1 Bq/Kg raw			Cs134		
White rice	Niigata	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Oosaki, Miyazaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Mie	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.6	Cs137	7.0 Bq/Kg raw		
			Cs134	1.6 Bq/Kg raw	± 2.1 Bq/Kg raw			Cs134		
White rice	Nagano	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Kurihara, Miyagi	Oct-10	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	7.0 Bq/Kg raw		
			Cs134	2.0 Bq/Kg raw	± 2.2 Bq/Kg raw			Cs134		
White rice	Tsuchiura, Ibaraki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Nihonmatsu	Oct-12	Cs137	1.5 Bq/Kg raw	± 2.1 Bq/Kg raw	1.5	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Nihonmatsu	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.0 Bq/Kg raw		
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw			Cs134		
White rice	Namie, Futaba	Oct-12	Cs137	1.8 Bq/Kg raw	± 2.2 Bq/Kg raw	4.1	Cs137	7.0 Bq/Kg raw		
			Cs134	2.3 Bq/Kg raw	± 1.9 Bq/Kg raw			Cs134		

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
White rice	Kitashirado, Taira, Iwak	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.5	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Toyoma, Taira, Iwaki	Oct-12	Cs137	3.5	Bq/Kg raw	± 2.2	Bq/Kg raw	5.1	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 1.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Akai, Taira, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Hashirikuma, Kashima, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.0	Cs137	7.0	Bq/Kg raw
			Cs134	3.0	Bq/Kg raw	± 3.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Watanabe, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Tono, Iwaki	Oct-12	Cs137	11.7	Bq/Kg raw	± 3.6	Bq/Kg raw	15.8	Cs137	7.0	Bq/Kg raw
			Cs134	4.1	Bq/Kg raw	± 2.2	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Tbito, Iwaki	不明	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Inoue, Yamada, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.2	Cs137	7.0	Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 1.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
White rice	Nakoso, Iwaki	Oct-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice(Brown rice)	Minamisoma	Oct-10	Cs137	1.5	Bq/Kg raw	± 2.4	Bq/Kg raw	1.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Sticky rice	Komoda, Kashima, Iwaki	Sep-12	Cs137	1.8	Bq/Kg raw	± 2.0	Bq/Kg raw	1.8	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Yotsukura, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Kubo, Kashima, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Japanese radish	Watanabe, Iwaki	Jan-13	Cs137	2.5	Bq/Kg raw	± 2.7	Bq/Kg raw	2.5	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pickled Japanese radish	Shimookeuri, Kawamae, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.4	Cs137	7.0	Bq/Kg raw
			Cs134	1.4	Bq/Kg raw	± 1.7	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Carrot	Kubo, Kashima, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Carrot	Watanabe, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Soybean	Kitsunozuka, Yotsukura, Iwaki	Dec-12	Cs137	14.8	Bq/Kg raw	± 4.4	Bq/Kg raw	18.3	Cs137	7.0	Bq/Kg raw
			Cs134	3.5	Bq/Kg raw	± 2.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Cauliflower	Kubo, Kashima, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Yotsukura, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Sakuragaoka, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Kubo, Kashima, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Broccoli	Watanabe, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection			
Chinese cabbage	Kitaibaraki	Dec-12	Cs137	1.9	Bq/Kg raw	± 2.2	Bq/Kg raw	3.5	Cs137	7.0	Bq/Kg raw
			Cs134	1.6	Bq/Kg raw	± 2.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Akai, Taira, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Chinese cabbage (Mixed)	Yagyū, Yotsukura, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Mizuna green	Sakuragaoka, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	5.0	Cs137	7.0	Bq/Kg raw
			Cs134	5.0	Bq/Kg raw	± 5.9	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Leek	Kitsunozuka, Yotsukura, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	3.4	Cs137	7.0	Bq/Kg raw
			Cs134	3.4	Bq/Kg raw	± 3.5	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Spinach	Watanabe, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Amitake mushroom	Hanawa, Higashishirakawa	Oct-12	Cs137	249	Bq/Kg raw	± 50.0	Bq/Kg raw	409	Cs137	7.0	Bq/Kg raw
			Cs134	160	Bq/Kg raw	± 32.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Jiyugaoka, Iwaki	Jan-13	Cs137	19.4	Bq/Kg raw	± 5.0	Bq/Kg raw	29.5	Cs137	7.0	Bq/Kg raw
			Cs134	10.1	Bq/Kg raw	± 3.1	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Yuzu	Shimokajiro, Onahama, Iwaki	Jan-13	Cs137	22.4	Bq/Kg raw	± 7.0	Bq/Kg raw	36.9	Cs137	7.0	Bq/Kg raw
			Cs134	14.5	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Kiwi fruit	Naraha, Futaba	Jan-13	Cs137	47.6	Bq/Kg raw	± 10.8	Bq/Kg raw	80.2	Cs137	7.0	Bq/Kg raw
			Cs134	32.6	Bq/Kg raw	± 7.4	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Dried persimmon (dried)	Shimokajiro, Onahama, Iwaki	Oct-12	Cs137	15.5	Bq/Kg raw	± 5.1	Bq/Kg raw	23.6	Cs137	7.0	Bq/Kg raw
			Cs134	8.1	Bq/Kg raw	± 3.6	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
School lunch	Unknown	Jan-13	Cs137	1.2	Bq/Kg raw	± 1.9	Bq/Kg raw	1.2	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
School lunch	Unknown	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
School lunch	Unknown	Jan-13	Cs137	1.6	Bq/Kg raw	± 2.0	Bq/Kg raw	1.6	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Rigid hornwort	Onahama, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	6.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	4.0	Bq/Kg raw
Seaweed	Onahama, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Well water	Koya, Uchigo, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Well water	Akai, Taira, Iwaki	Jan-13	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.9	Cs137	7.0	Bq/Kg raw
			Cs134	0.9	Bq/Kg raw	± 1.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Pickled Japanese radish	Yamada, Iwaki	Dec-12	Cs137	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw	0.0	Cs137	7.0	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0	Bq/Kg raw		Cs134	6.0	Bq/Kg raw
Bamboo(sawdust)	Akai, Taira, Iwaki	Jan-13	Cs137	27.0	Bq/Kg raw	± 8.0	Bq/Kg raw	41.0	Cs131	7.0	Bq/Kg raw
			Cs134	14.0	Bq/Kg raw	± 5.0	Bq/Kg raw		Cs128	6.0	Bq/Kg raw
Bamboo(sawdust)	Akai, Taira, Iwaki	Jan-13	Cs137	236	Bq/Kg raw	± 50.0	Bq/Kg raw	384	Cs125	7.0	Bq/Kg raw
			Cs134	148	Bq/Kg raw	± 31.0	Bq/Kg raw		Cs122	6.0	Bq/Kg raw
Bamboo(sawdust)	Akai, Taira, Iwaki	Jan-13	Cs137	8.0	Bq/Kg raw	± 4.0	Bq/Kg raw	17.0	Cs119	7.0	Bq/Kg raw
			Cs134	9.0	Bq/Kg raw	± 3.0	Bq/Kg raw		Cs116	6.0	Bq/Kg raw

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Bamboo(sawdust)	Akai, Taira, Iwaki	Jan-13	Cs137	20.0 <small>Bq/Kg raw</small>	± 7.0 <small>Bq/Kg raw</small>	21.0	Cs113	7.0 <small>Bq/Kg raw</small>	
			Cs134	1.0 <small>Bq/Kg raw</small>	± 5.0 <small>Bq/Kg raw</small>		Cs110	5.0 <small>Bq/Kg raw</small>	
Yamazakura tree	Uwadaira, Ogawa, Iwaki	Jan-13	Cs137	155 <small>Bq/Kg raw</small>	± 37.0 <small>Bq/Kg raw</small>	265	Cs107	7.0 <small>Bq/Kg raw</small>	
			Cs134	110 <small>Bq/Kg raw</small>	± 26.0 <small>Bq/Kg raw</small>		Cs104	6.0 <small>Bq/Kg raw</small>	
Oak	Uwadaira, Ogawa, Iwaki	Jan-13	Cs137	205 <small>Bq/Kg raw</small>	± 44.0 <small>Bq/Kg raw</small>	343	Cs101	7.0 <small>Bq/Kg raw</small>	
			Cs134	138 <small>Bq/Kg raw</small>	± 29.0 <small>Bq/Kg raw</small>		Cs98	6.0 <small>Bq/Kg raw</small>	
Fallen leaves	Kawauchi, Futaba	Jan-13	Cs137	407 <small>Bq/Kg raw</small>	± 88.0 <small>Bq/Kg raw</small>	693	Cs95	7.0 <small>Bq/Kg raw</small>	
			Cs134	286 <small>Bq/Kg raw</small>	± 60.0 <small>Bq/Kg raw</small>		Cs92	6.0 <small>Bq/Kg raw</small>	
Leaf mold	Kamikatayose, Taira, Iwaki	Dec-12	Cs137	139 <small>Bq/Kg dry</small>	± 38.0 <small>Bq/Kg dry</small>	193	Cs137	8.0 <small>Bq/Kg dry</small>	
			Cs134	54.0 <small>Bq/Kg dry</small>	± 20.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>	
Leaf mold	Misawa, Joban, Iwaki	Jan-13	Cs137	674 <small>Bq/Kg dry</small>	± 84.0 <small>Bq/Kg dry</small>	1,008	Cs137	8.0 <small>Bq/Kg dry</small>	
			Cs134	334 <small>Bq/Kg dry</small>	± 44.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>	
Field soil	Ogawa, Iwaki	Jan-13	Cs137	155 <small>Bq/Kg dry</small>	± 5.0 <small>Bq/Kg dry</small>	215	Cs137	8.0 <small>Bq/Kg dry</small>	
			Cs134	60.0 <small>Bq/Kg dry</small>	± 5.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>	
Field soil	Ojiroi, Kawamae, Iwaki	Dec-12	Cs137	9963 <small>Bq/Kg dry</small>	± 863 <small>Bq/Kg dry</small>	14,526	Cs137	8.0 <small>Bq/Kg dry</small>	
			Cs134	4563 <small>Bq/Kg dry</small>	± 402 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>	
Field soil	Ojiroi, Kawamae, Iwaki	Dec-12	Cs137	7460 <small>Bq/Kg dry</small>	± 1490 <small>Bq/Kg dry</small>	12,280	Cs137	7.0 <small>Bq/Kg dry</small>	
			Cs134	4820 <small>Bq/Kg dry</small>	± 960 <small>Bq/Kg dry</small>		Cs134	6.0 <small>Bq/Kg dry</small>	
Field soil	Kamikatayose, Taira, Iwaki	Jan-13	Cs137	199 <small>Bq/Kg dry</small>	± 28.0 <small>Bq/Kg dry</small>	292	Cs137	8.0 <small>Bq/Kg dry</small>	
			Cs134	93.0 <small>Bq/Kg dry</small>	± 14.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>	
Field soil	Akai, Taira, Iwaki	Jan-13	Cs137	250 <small>Bq/Kg dry</small>	± 33.0 <small>Bq/Kg dry</small>	362	Cs137	8.0 <small>Bq/Kg dry</small>	
			Cs134	112 <small>Bq/Kg dry</small>	± 18.0 <small>Bq/Kg dry</small>		Cs134	8.0 <small>Bq/Kg dry</small>	