



Radiation Measurement Results of 90 items in April 2013



※The analytical software we used for radioactivity measurement between November 2011 and March 2013 was older than the one we use currently, therefore, the both values for measured results and minimum limit of detection during this period may differ comparing the present measured values. Please refer those values as an indication for radiation dose at the time of the Daiichi Fukushima nuclear accident.

When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Yotsukura, Iwaki	Oct-12	Cs137	1.5 Bq/Kg raw	± 2.5 Bq/Kg raw	1.5	Cs137	3.8 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.4 Bq/Kg raw	
Brown rice	Kawabe, Iwaki	Apr-13	Cs137	7.3 Bq/Kg raw	± 2.4 Bq/Kg raw	9.8	Cs137	2.8 Bq/Kg raw	
			Cs134	2.5 Bq/Kg raw	± 1.7 Bq/Kg raw		Cs134	2.6 Bq/Kg raw	
White rice	Yamada, Iwaki	Oct-12	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	3.8 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.5 Bq/Kg raw	
White rice	Tabito, Iwaki	Oct-11	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	3.6 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.3 Bq/Kg raw	
White rice	Unknown	Unknown	Cs137	7.9 Bq/Kg raw	± 2.6 Bq/Kg raw	9.0	Cs137	3.0 Bq/Kg raw	
			Cs134	1.1 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	2.8 Bq/Kg raw	
Chinese chive	Yotsukura, Iwaki	Apr-13	Cs137	4.3 Bq/Kg raw	± 4.9 Bq/Kg raw	4.3	Cs137	7.9 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	7.3 Bq/Kg raw	
Chinese chive	Kamikajiro, Onahama, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.1	Cs137	6.0 Bq/Kg raw	
			Cs134	3.1 Bq/Kg raw	± 3.6 Bq/Kg raw		Cs134	5.4 Bq/Kg raw	
Turnip	Kamikajiro, Onahama, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	5.4 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	4.7 Bq/Kg raw	
Bok choy	Shimoyada, Kashima, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	3.7	Cs137	3.9 Bq/Kg raw	
			Cs134	3.7 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	3.4 Bq/Kg raw	
Leek	Nishiki, Iwaki	Apr-13	Cs137	1.3 Bq/Kg raw	± 2.1 Bq/Kg raw	1.3	Cs137	3.4 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.2 Bq/Kg raw	
Japanese radish leaf	Nishiki, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	3.9 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.5 Bq/Kg raw	
Chinese colza	Nishigo, Joban, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	7.1 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	6.2 Bq/Kg raw	
Cabbage	Kamioyoe, Taira, Iwaki	Mar-13	Cs137	1.5 Bq/Kg raw	± 1.7 Bq/Kg raw	1.5	Cs137	2.6 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	2.5 Bq/Kg raw	
Yam	Kashima, Iwaki	Mar-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.0	Cs137	4.1 Bq/Kg raw	
			Cs134	2.0 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	3.8 Bq/Kg raw	
Fatsia sprout	Shimoyoshima, Yoshima, Iwaki	Apr-13	Cs137	4.3 Bq/Kg raw	± 4.4 Bq/Kg raw	7.0	Cs137	6.6 Bq/Kg raw	
			Cs134	2.7 Bq/Kg raw	± 4.0 Bq/Kg raw		Cs134	6.1 Bq/Kg raw	
Fatsia sprout	Okaona, Onahama, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	2.4	Cs137	4.5 Bq/Kg raw	
			Cs134	2.4 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	3.9 Bq/Kg raw	
Fatsia sprout	Shimokawa, Izumi, Iwaki	Apr-13	Cs137	16.7 Bq/Kg raw	± 6.1 Bq/Kg raw	27.5	Cs137	6.6 Bq/Kg raw	
			Cs134	10.8 Bq/Kg raw	± 4.7 Bq/Kg raw		Cs134	6.2 Bq/Kg raw	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Fatsia sprout	Shimokawa, Izumi, Iwaki	Apr-13	Cs137	30.2	Bq/Kg raw	± 8.4	52.1	Cs137	6.4
			Cs134	21.9	Bq/Kg raw	± 6.2		Cs134	5.8
Fatsia sprout	Tabito, Iwaki	Apr-13	Cs137	6.7	Bq/Kg raw	± 2.7	12.7	Cs137	3.1
			Cs134	6.0	Bq/Kg raw	± 2.3		Cs134	2.8
Bamboo shoot	Ohisa, Iwaki	Apr-13	Cs137	23.3	Bq/Kg raw	± 5.7	35.7	Cs137	3.3
			Cs134	12.4	Bq/Kg raw	± 3.5		Cs134	3.1
Bamboo shoot	Ohisa, Iwaki	Apr-13	Cs137	15.0	Bq/Kg raw	± 4.1	23.2	Cs137	3.3
			Cs134	8.2	Bq/Kg raw	± 2.8		Cs134	3.1
Bamboo shoot	Ohisa, Iwaki	Apr-13	Cs137	11.6	Bq/Kg raw	± 3.4	14.8	Cs137	3.3
			Cs134	3.2	Bq/Kg raw	± 2.2		Cs134	3.1
Bamboo shoot	Ohisa, Iwaki	Apr-13	Cs137	20.6	Bq/Kg raw	± 5.2	30.1	Cs137	3.4
			Cs134	9.5	Bq/Kg raw	± 3.1		Cs134	3.1
Bamboo shoot	Yoshima, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	10.2
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	9.0
Bamboo shoot	Sakura, Taira, Iwaki	Apr-13	Cs137	10.3	Bq/Kg raw	± 3.7	19.2	Cs137	3.8
			Cs134	8.9	Bq/Kg raw	± 3.0		Cs134	3.3
Bamboo shoot	Akai, Taira, Iwaki	Apr-13	Cs137	18.7	Bq/Kg raw	± 5.0	29.9	Cs137	3.7
			Cs134	11.2	Bq/Kg raw	± 3.4		Cs134	3.3
Bamboo shoot	Akai, Taira, Iwaki	Apr-13	Cs137	11.9	Bq/Kg raw	± 3.8	19.2	Cs137	3.8
			Cs134	7.3	Bq/Kg raw	± 2.8		Cs134	3.4
Bamboo shoot	Akai, Taira, Iwaki	Apr-13	Cs137	7.2	Bq/Kg raw	± 5.2	7.2	Cs137	8.0
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	7.3
Bamboo shoot	Watanabe, Iwaki	Apr-13	Cs137	19.6	Bq/Kg raw	± 5.3	31.3	Cs137	4.1
			Cs134	11.7	Bq/Kg raw	± 3.6		Cs134	3.9
Bamboo shoot	Hashirikuma, Kashima, Iwaki	Apr-13	Cs137	104	Bq/Kg raw	± 21.0	163	Cs137	3.5
			Cs134	58.5	Bq/Kg raw	± 12.0		Cs134	3.1
Bamboo shoot	Yamadama, Iwaki	Apr-13	Cs137	11.9	Bq/Kg raw	± 3.9	20.5	Cs137	3.7
			Cs134	8.6	Bq/Kg raw	± 3.0		Cs134	3.5
Bamboo shoot	Miyo, Kashima, Iwaki	Apr-13	Cs137	8.7	Bq/Kg raw	± 3.5	15.6	Cs137	3.9
			Cs134	6.9	Bq/Kg raw	± 2.8		Cs134	3.4
Bamboo shoot	Onahama, Iwaki	Apr-13	Cs137	21.5	Bq/Kg raw	± 7.3	28.5	Cs137	4.2
			Cs134	7.0	Bq/Kg raw	± 5.0		Cs134	5.0
Bamboo shoot (boiled)	Shibahara, Ogawa, Iwaki	Apr-13	Cs137	2.9	Bq/Kg raw	± 2.2	2.9	Cs137	3.3
			Cs134	0.0	Bq/Kg raw	± 0.0		Cs134	2.9
Bamboo shoot (boiled)	Nakashio, Taira, Iwaki	Apr-13	Cs137	14.9	Bq/Kg raw	± 4.3	25.6	Cs137	3.5
			Cs134	10.7	Bq/Kg raw	± 3.2		Cs134	3.3
Bamboo shoot (boiled)	Shimotakaku, Taira, Iwaki	Apr-13	Cs137	100	Bq/Kg raw	± 21.0	160	Cs137	3.5
			Cs134	60.1	Bq/Kg raw	± 12.3		Cs134	3.1
Bamboo shoot (boiled)	Miyo, Kashima, Iwaki	Apr-13	Cs137	7.0	Bq/Kg raw	± 2.4	9.9	Cs137	2.6
			Cs134	2.9	Bq/Kg raw	± 1.7		Cs134	2.5
Bamboo shoot (boiled)	Shimokawa, Onahama, Iwaki	Apr-13	Cs137	12.9	Bq/Kg raw	± 4.1	17.6	Cs137	4.3
			Cs134	4.7	Bq/Kg raw	± 2.8		Cs134	4.0
Bamboo shoot (boiled)	Kanari, Onahama, Iwaki	Apr-13	Cs137	9.5	Bq/Kg raw	± 3.3	16.6	Cs137	3.4
			Cs134	7.1	Bq/Kg raw	± 2.6		Cs134	3.1
Bracken	Komagome, Yotsukura, Iwaki	Apr-13	Cs137	35.5	Bq/Kg raw	± 8.2	59.5	Cs137	3.2
			Cs134	24.0	Bq/Kg raw	± 5.5		Cs134	2.9

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Bracken	Kashima, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	2.4	Cs137	4.8	Bq/Kg raw
			Cs134	2.4	Bq/Kg raw	± 2.8		Bq/Kg raw	Cs134	4.2
Bracken(boiled)	Uchigo, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	3.8	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	3.5
Bracken(boiled)	Izumi, Iwaki	Apr-13	Cs137	1.5	Bq/Kg raw	± 2.3	1.5	Cs137	3.6	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	3.4
Royal fern	Ohisa, Iwaki	Apr-13	Cs137	58.8	Bq/Kg raw	± 12.7	92.9	Cs137	3.4	Bq/Kg raw
			Cs134	34.1	Bq/Kg raw	± 7.5		Bq/Kg raw	Cs134	3.1
Ostrich fern	Kanari, Onahama, Iwaki	Apr-13	Cs137	9.5	Bq/Kg raw	± 6.3	9.5	Cs137	10.3	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	9.2
Udo	Shimokawa, Izumi, Iwaki	Apr-13	Cs137	25.2	Bq/Kg raw	± 6.5	34.0	Cs137	4.9	Bq/Kg raw
			Cs134	8.8	Bq/Kg raw	± 3.6		Bq/Kg raw	Cs134	3.5
Koshiabura tree	Kadouno, Tono, Iwaki	Apr-13	Cs137	639	Bq/Kg raw	± 128	1,015	Cs137	6.2	Bq/Kg raw
			Cs134	376	Bq/Kg raw	± 75.0		Bq/Kg raw	Cs134	5.4
Fuki	Yoshima, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	1.8	Cs137	4.6	Bq/Kg raw
			Cs134	1.8	Bq/Kg raw	± 2.7		Bq/Kg raw	Cs134	4.1
Fuki	Makunouchi, Taira, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	4.9	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	4.6
Fuki	Shimotakaku, Taira, Iwaki	Apr-13	Cs137	6.6	Bq/Kg raw	± 3.9	9.4	Cs137	5.8	Bq/Kg raw
			Cs134	2.8	Bq/Kg raw	± 3.3		Bq/Kg raw	Cs134	5.1
Fuki	Suginami, Taira, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	0.0	Cs137	10.5	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	9.3
Fuki	Watanabe, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	6.5	Cs137	4.5	Bq/Kg raw
			Cs134	6.5	Bq/Kg raw	± 3.0		Bq/Kg raw	Cs134	4.1
Boiled Fuki	Onahama, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	2.7	Cs137	3.7	Bq/Kg raw
			Cs134	2.7	Bq/Kg raw	± 2.2		Bq/Kg raw	Cs134	3.4
Fuki (without skin)	Kamikajiro, Onahama, Iwaki	Apr-13	Cs137	0.0	Bq/Kg raw	± 0.0	1.2	Cs137	3.2	Bq/Kg raw
			Cs134	1.2	Bq/Kg raw	± 1.9		Bq/Kg raw	Cs134	2.9
Fuki (without skin)	Akai, Taira, Iwaki	Apr-13	Cs137	2.1	Bq/Kg raw	± 3.6	2.1	Cs137	5.6	Bq/Kg raw
			Cs134	0.0	Bq/Kg raw	± 0.0		Bq/Kg raw	Cs134	5.2
Amigasatake mushroom	Onahama, Iwaki	Apr-13	Cs137	13.8	Bq/Kg raw	± 9.3	18.5	Cs137	5.4	Bq/Kg raw
			Cs134	4.7	Bq/Kg raw	± 4.4		Bq/Kg raw	Cs134	5.4
Shiitake mushroom	Shibahara, Ogawa, Iwaki	Apr-13	Cs137	52.0	Bq/Kg raw	± 11.4	81.7	Cs137	3.4	Bq/Kg raw
			Cs134	29.7	Bq/Kg raw	± 6.7		Bq/Kg raw	Cs134	3.1
Shiitake mushroom	Suginami, Taira, Iwaki	Apr-13	Cs137	385	Bq/Kg raw	± 77.0	599	Cs137	3.9	Bq/Kg raw
			Cs134	214	Bq/Kg raw	± 43.0		Bq/Kg raw	Cs134	3.6
Shiitake mushroom	Numanouchi, Taira, Iwaki	Apr-13	Cs137	1014	Bq/Kg raw	± 108	1,444	Cs137	10.2	Bq/Kg raw
			Cs134	430	Bq/Kg raw	± 50.5		Bq/Kg raw	Cs134	9.0
Shiitake mushroom	Izumi, Iwaki	Apr-13	Cs137	81.5	Bq/Kg raw	± 18.3	114	Cs137	7.5	Bq/Kg raw
			Cs134	32.6	Bq/Kg raw	± 8.4		Bq/Kg raw	Cs134	7.0
Shiitake mushroom	Onahama, Iwaki	Apr-13	Cs137	122	Bq/Kg raw	± 21.2	162	Cs137	8.1	Bq/Kg raw
			Cs134	39.6	Bq/Kg raw	± 10.2		Bq/Kg raw	Cs134	7.2
Shiitake mushroom	Obanma, Iwaki	Apr-13	Cs137	10.5	Bq/Kg raw	± 5.4	15.3	Cs137	4.6	Bq/Kg raw
			Cs134	4.8	Bq/Kg raw	± 5.0		Bq/Kg raw	Cs134	6.3
Shiitake mushroom	Obanma, Iwaki	Apr-13	Cs137	143	Bq/Kg raw	± 29.0	215	Cs137	4.7	Bq/Kg raw
			Cs134	72.1	Bq/Kg raw	± 15.0		Bq/Kg raw	Cs134	4.3

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Shiitake mushroom	Negishi, Tono, Iwaki	Apr-13	Cs137	93.3 Bq/Kg raw	± 19.2 Bq/Kg raw	148	Cs137	3.5 Bq/Kg raw	
			Cs134	54.6 Bq/Kg raw	± 11.3 Bq/Kg raw		Cs134	3.3 Bq/Kg raw	
Shiitake mushroom	Tabito, Iwaki	Apr-13	Cs137	130 Bq/Kg raw	± 26.0 Bq/Kg raw	201	Cs137	4.5 Bq/Kg raw	
			Cs134	70.7 Bq/Kg raw	± 14.6 Bq/Kg raw		Cs134	4.0 Bq/Kg raw	
Shiitake mushroom	Shimoyamaguchi, Taira, Iwaki	Apr-13	Cs137	297 Bq/Kg raw	± 35.4 Bq/Kg raw	416	Cs137	4.9 Bq/Kg raw	
			Cs134	119 Bq/Kg raw	± 16.5 Bq/Kg raw		Cs134	5.6 Bq/Kg raw	
Shiitake mushroom	Mimaya, Uchigo, Iwaki	Apr-13	Cs137	356 Bq/Kg raw	± 71.0 Bq/Kg raw	569	Cs137	4.9 Bq/Kg raw	
			Cs134	213 Bq/Kg raw	± 43.0 Bq/Kg raw		Cs134	4.6 Bq/Kg raw	
Shiitake mushroom	Kawabe, Iwaki	Apr-13	Cs137	259 Bq/Kg raw	± 52.0 Bq/Kg raw	405	Cs137	4.8 Bq/Kg raw	
			Cs134	146 Bq/Kg raw	± 29.0 Bq/Kg raw		Cs134	4.5 Bq/Kg raw	
Amanatsu orange	Izumi, Iwaki	Apr-13	Cs137	12.2 Bq/Kg raw	± 4.2 Bq/Kg raw	20.5	Cs137	4.4 Bq/Kg raw	
			Cs134	8.3 Bq/Kg raw	± 3.2 Bq/Kg raw		Cs134	3.9 Bq/Kg raw	
Amanatsu orange marmalade	Uchigo, Iwaki	Feb-13	Cs137	11.2 Bq/Kg raw	± 3.1 Bq/Kg raw	16.0	Cs137	2.7 Bq/Kg raw	
			Cs134	4.8 Bq/Kg raw	± 2.0 Bq/Kg raw		Cs134	2.5 Bq/Kg raw	
Oyster (boiled)	Onahama, Iwaki	Mar-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	24.3	Cs137	3.6 Bq/Kg raw	
			Cs134	24.3 Bq/Kg raw	± 5.7 Bq/Kg raw		Cs134	3.4 Bq/Kg raw	
Char	Shitoki River, Iwaki	Apr-13	Cs137	151 Bq/Kg raw	± 21.9 Bq/Kg raw	222	Cs137	4.6 Bq/Kg raw	
			Cs134	71.4 Bq/Kg raw	± 11.3 Bq/Kg raw		Cs134	5.2 Bq/Kg raw	
Char	Shitoki River, Iwaki	Apr-13	Cs137	142 Bq/Kg raw	± 20.6 Bq/Kg raw	200	Cs137	4.9 Bq/Kg raw	
			Cs134	57.7 Bq/Kg raw	± 10.0 Bq/Kg raw		Cs134	5.3 Bq/Kg raw	
Seaweed (boiled)	Usuiso, Taira, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	3.8 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.5 Bq/Kg raw	
School lunch	Unknown	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	3.3 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.1 Bq/Kg raw	
School lunch	Unknown	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.0	Cs137	3.4 Bq/Kg raw	
			Cs134	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw		Cs134	3.2 Bq/Kg raw	
School lunch	Unknown	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	1.3	Cs137	3.3 Bq/Kg raw	
			Cs134	1.3 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	2.9 Bq/Kg raw	
Fermented rice bran	Unknown	Unknown	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	4.3	Cs137	3.5 Bq/Kg raw	
			Cs134	4.3 Bq/Kg raw	± 2.3 Bq/Kg raw		Cs134	3.2 Bq/Kg raw	
Fermented rice bran	Unknown	Apr-13	Cs137	3.6 Bq/Kg raw	± 2.1 Bq/Kg raw	5.8	Cs137	2.9 Bq/Kg raw	
			Cs134	2.2 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	2.6 Bq/Kg raw	
Well water	Akai, Taira, Iwaki	Apr-13	Cs137	0.0 Bq/Kg raw	± 0.0 Bq/Kg raw	0.8	Cs137	1.9 Bq/Kg raw	
			Cs134	0.8 Bq/Kg raw	± 1.1 Bq/Kg raw		Cs134	1.7 Bq/Kg raw	
Wood chip	Kashima, Iwaki	Apr-13	Cs137	2118 Bq/Kg raw	± 224 Bq/Kg raw	2,995	Cs107	8.0 Bq/Kg raw	
			Cs134	877 Bq/Kg raw	± 102 Bq/Kg raw		Cs104	8.0 Bq/Kg raw	
Charcoal	Tabito, Iwaki	Nov-11	Cs137	6030 Bq/Kg raw	± 1240 Bq/Kg raw	9,590	Cs89	7.0 Bq/Kg raw	
			Cs134	3560 Bq/Kg raw	± 710 Bq/Kg raw		Cs86	6.0 Bq/Kg raw	
Sand(Surface)	Unknown	Unknown	Cs137	9.0 Bq/Kg dry	± 4.0 Bq/Kg dry	9.0	Cs137	8.0 Bq/Kg dry	
			Cs134	0.0 Bq/Kg dry	± 0.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Sand (from 15cm deep)	Unknown	Unknown	Cs137	4.0 Bq/Kg dry	± 5.0 Bq/Kg dry	7.0	Cs137	8.0 Bq/Kg dry	
			Cs134	3.0 Bq/Kg dry	± 4.0 Bq/Kg dry		Cs134	8.0 Bq/Kg dry	
Sand (from 30cm deep)	Unknown	Unknown	Cs137	11.0 Bq/Kg dry	± 5.0 Bq/Kg dry	11.0	Cs131	8.0 Bq/Kg dry	
			Cs134	0.0 Bq/Kg dry	± 2.0 Bq/Kg dry		Cs128	8.0 Bq/Kg dry	
Sand (from 50cm deep)	Unknown	Unknown	Cs137	11.0 Bq/Kg dry	± 5.0 Bq/Kg dry	11.0	Cs125	8.0 Bq/Kg dry	
			Cs134	0.0 Bq/Kg dry	± 2.0 Bq/Kg dry		Cs122	8.0 Bq/Kg dry	

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Field soil	Akai, Taira, Iwaki	Apr-13	Cs137	253 <small>Bq/Kg dry</small>	± 33.0 <small>Bq/Kg dry</small>	351	Cs101	8.0 <small>Bq/Kg dry</small>
			Cs134	98.0 <small>Bq/Kg dry</small>	± 16.0 <small>Bq/Kg dry</small>		Cs98	8.0 <small>Bq/Kg dry</small>
Field soil	Iino, Chuodai, Iwaki	Apr-13	Cs137	78.0 <small>Bq/Kg dry</small>	± 16.0 <small>Bq/Kg dry</small>	114	Cs119	8.0 <small>Bq/Kg dry</small>
			Cs134	36.0 <small>Bq/Kg dry</small>	± 9.0 <small>Bq/Kg dry</small>		Cs116	8.0 <small>Bq/Kg dry</small>
Field soil	Nishiki, Iwaki	Apr-13	Cs137	170 <small>Bq/Kg dry</small>	± 24.0 <small>Bq/Kg dry</small>	232	Cs113	8.0 <small>Bq/Kg dry</small>
			Cs134	62.0 <small>Bq/Kg dry</small>	± 12.0 <small>Bq/Kg dry</small>		Cs110	8.0 <small>Bq/Kg dry</small>
Public garden soil	Takaku, Chuodai, Iwaki	Apr-13	Cs137	240 <small>Bq/Kg dry</small>	± 31.0 <small>Bq/Kg dry</small>	333	Cs95	8.0 <small>Bq/Kg dry</small>
			Cs134	93.0 <small>Bq/Kg dry</small>	± 15.0 <small>Bq/Kg dry</small>		Cs92	8.0 <small>Bq/Kg dry</small>

