




Radiation Measurement Results of 98 Items in April-May



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Bamboo shoots (raw)	Iwaki	Apr-15	Cs137	62.7 Bq/Kg raw	± 12.8 Bq/Kg raw	84.6	Cs137	5.4 Bq/Kg raw	
			Cs134	21.9 Bq/Kg raw	± 5.0 Bq/Kg raw		Cs134	2.8 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	Apr-15	Cs137	15.0 Bq/Kg raw	± 4.6 Bq/Kg raw	15.0	Cs137	5.1 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	4.7 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	May-15	Cs137	7.3 Bq/Kg raw	± 1.7 Bq/Kg raw	9.6	Cs137	1.1 Bq/Kg raw	
			Cs134	2.2 Bq/Kg raw	± 0.8 Bq/Kg raw		Cs134	1.0 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	May-15	Cs137	16.5 Bq/Kg raw	± 4.0 Bq/Kg raw	23.6	Cs137	2.5 Bq/Kg raw	
			Cs134	7.1 Bq/Kg raw	± 2.2 Bq/Kg raw		Cs134	2.4 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	May-15	Cs137	19.6 Bq/Kg raw	± 4.7 Bq/Kg raw	28.4	Cs137	2.9 Bq/Kg raw	
			Cs134	8.8 Bq/Kg raw	± 2.6 Bq/Kg raw		Cs134	2.6 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	May-15	Cs137	8.5 Bq/Kg raw	± 2.3 Bq/Kg raw	10.0	Cs137	3.2 Bq/Kg raw	
			Cs134	1.5 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	2.9 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	May-15	Cs137	26.4 Bq/Kg raw	± 6.4 Bq/Kg raw	35.7	Cs137	3.7 Bq/Kg raw	
			Cs134	9.3 Bq/Kg raw	± 3.1 Bq/Kg raw		Cs134	3.4 Bq/Kg raw	
Bamboo shoots (raw)	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.9 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.6 Bq/Kg raw	
Bamboo shoots (boiled)	Iwaki	Apr-15	Cs137	134 Bq/Kg raw	± 27.0 Bq/Kg raw	175	Cs137	5.2 Bq/Kg raw	
			Cs134	41.1 Bq/Kg raw	± 9.4 Bq/Kg raw		Cs134	4.8 Bq/Kg raw	
Bamboo shoots (boiled 3 times)	Iwaki	Apr-15	Cs137	6.7 Bq/Kg raw	± 2.6 Bq/Kg raw	9.9	Cs137	3.0 Bq/Kg raw	
			Cs134	3.2 Bq/Kg raw	± 1.8 Bq/Kg raw		Cs134	2.8 Bq/Kg raw	
Warabi (Edible wild plant)	Iwaki	Apr-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.2 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.9 Bq/Kg raw	
Warabi (Edible wild plant)	Iwaki	Apr-15	Cs137	14.6 Bq/Kg raw	± 3.4 Bq/Kg raw	21.2	Cs137	2.3 Bq/Kg raw	
			Cs134	6.6 Bq/Kg raw	± 1.9 Bq/Kg raw		Cs134	2.1 Bq/Kg raw	
Warabi (Edible wild plant)	Tomioka	Apr-15	Cs137	124 Bq/Kg raw	± 26.0 Bq/Kg raw	161	Cs137	6.3 Bq/Kg raw	
			Cs134	36.8 Bq/Kg raw	± 8.8 Bq/Kg raw		Cs134	5.8 Bq/Kg raw	
Warabi (Edible wild plant)	Iwaki	Apr-15	Cs137	5.0 Bq/Kg raw	± 2.1 Bq/Kg raw	5.0	Cs137	2.8 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.6 Bq/Kg raw	
Warabi (Edible wild plant)	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.4 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.7 Bq/Kg raw	
Warabi (Edible wild plant)	Furudono	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.0 Bq/Kg raw	
Warabi (Edible wild plant)	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.9 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.6 Bq/Kg raw	
Warabi (Edible wild plant)	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.6 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.4 Bq/Kg raw	

※“—” used in Measurement Result and Uncertainty shows that the value is below the detection limit. But it does not necessary mean 0(zero)Bq/Kg.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Warabi leaf (Edible wild plant)	Tomioka	Apr-15	Cs137	526 Bq/Kg raw	± 107 Bq/Kg raw	682	Cs137	22.5 Bq/Kg raw	
			Cs134	156 Bq/Kg raw	± 36.0 Bq/Kg raw		Cs134	21.2 Bq/Kg raw	
Zenmai (raw) (Edible wild plant)	Tomioka	Apr-15	Cs137	798 Bq/Kg raw	± 160 Bq/Kg raw	1058	Cs137	8.0 Bq/Kg raw	
			Cs134	260 Bq/Kg raw	± 52.0 Bq/Kg raw		Cs134	7.2 Bq/Kg raw	
Zenmai (boiled) (Edible wild plant)	Tomioka	Apr-15	Cs137	480 Bq/Kg raw	± 96.0 Bq/Kg raw	629	Cs137	4.7 Bq/Kg raw	
			Cs134	149 Bq/Kg raw	± 30.0 Bq/Kg raw		Cs134	4.3 Bq/Kg raw	
Zenmai leaf (Edible wild plant)	Tomioka	Apr-15	Cs137	3380 Bq/Kg raw	± 680 Bq/Kg raw	4410	Cs137	8.0 Bq/Kg raw	
			Cs134	1030 Bq/Kg raw	± 210 Bq/Kg raw		Cs134	7.5 Bq/Kg raw	
Zenmai Cotton (Edible wild plant)	Tomioka	Apr-15	Cs137	2250 Bq/Kg raw	± 450 Bq/Kg raw	3008	Cs137	46.7 Bq/Kg raw	
			Cs134	758 Bq/Kg raw	± 156 Bq/Kg raw		Cs134	42.6 Bq/Kg raw	
Udo (Edible wild plant)	Iwaki	Apr-15	Cs137	5.3 Bq/Kg raw	± 1.5 Bq/Kg raw	7.4	Cs137	1.5 Bq/Kg raw	
			Cs134	2.1 Bq/Kg raw	± 1.0 Bq/Kg raw		Cs134	1.4 Bq/Kg raw	
Udo (Edible wild plant)	Iwaki	Mar-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	1.7 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	1.6 Bq/Kg raw	
Shidoki (Edible wild plant)	Iwaki	Apr-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	5.7 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	5.3 Bq/Kg raw	
Shidoki (Edible wild plant)	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	16.4 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	13.1 Bq/Kg raw	
Water Dropwort	Iwaki	Apr-15	Cs137	9.4 Bq/Kg raw	± 2.7 Bq/Kg raw	9.4	Cs137	2.9 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.7 Bq/Kg raw	
Fuki (Edible wild plant)	Iwaki	Apr-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.2 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	3.0 Bq/Kg raw	
Fuki leaf (Edible wild plant)	Iwaki	Apr-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	6.5 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	6.1 Bq/Kg raw	
Rice	Hokkaido	Oct-14	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	2.4 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	2.2 Bq/Kg dry	
Rice	Akita Pref	Oct-14	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	2.4 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	2.2 Bq/Kg dry	
Rice	Iwaki	Oct-14	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	2.4 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	2.2 Bq/Kg dry	
Brown rice	Iwaki	Unknown	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	2.6 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	2.3 Bq/Kg dry	
Brown rice	Iwaki	Oct-14	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	2.1 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	2.0 Bq/Kg dry	
Santona	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.0 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.7 Bq/Kg raw	
Kashiwa leaf	Iwaki	May-15	Cs137	4.4 Bq/Kg raw	± 1.5 Bq/Kg raw	6.7	Cs137	1.9 Bq/Kg raw	
			Cs134	2.3 Bq/Kg raw	± 1.2 Bq/Kg raw		Cs134	1.7 Bq/Kg raw	
Daikon radish	Chiba Pref	Apr-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.6 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.4 Bq/Kg raw	
Daikon radish	Iwaki	May-15	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	2.0 Bq/Kg raw	

※“—” used in Measurement Result and Uncertainty shows that the value is below the detection limit. But it does not necessary mean 0(zero)Bq/Kg.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Bok Choy	Ibaraki Pref	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.0 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.7 Bq/Kg raw
New onion	Iwaki	Mar-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.7 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.4 Bq/Kg raw
Potato	Koriyama	Jun-14	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.9 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.7 Bq/Kg raw
Potato	Iwaki	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.3 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.1 Bq/Kg raw
Corbicula	Hinuma in Ibaraki Pref	Unknown	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.4 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.8 Bq/Kg raw
Dried shrimp	Iwate Pref	Unknown	Cs137	— Bq/Kg dry	±	— Bq/Kg dry	Under Minimum Limit of Detection	Cs137	3.4 Bq/Kg dry
			Cs134	— Bq/Kg dry	±	— Bq/Kg dry		Cs134	3.1 Bq/Kg dry
Shirasu (Fish)	Hiroshima Pref	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.6 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.4 Bq/Kg raw
Milk	Aizu in Fukushima Pref	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	1.7 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	1.6 Bq/Kg raw
Hen's egg	Hanawa	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	1.8 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	1.6 Bq/Kg raw
Soybean	Iwaki	Unknown	Cs137	— Bq/Kg dry	±	— Bq/Kg dry	Under Minimum Limit of Detection	Cs137	3.5 Bq/Kg dry
			Cs134	— Bq/Kg dry	±	— Bq/Kg dry		Cs134	3.1 Bq/Kg dry
Miso (Soybean paste)	Iwaki	Unknown	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.5 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.6 Bq/Kg raw
Okara (Soybean curd refuse)	Iwaki	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.9 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.6 Bq/Kg raw
Spirulina	Hawaii	Unknown	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.6 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.4 Bq/Kg raw
Welsh onion	Iwaki	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.8 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.6 Bq/Kg raw
Welsh onion	Iwaki	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.9 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	3.6 Bq/Kg raw
Yam	Gunma Pref	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	4.0 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	3.8 Bq/Kg raw
Snow Pea	Iwaki	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.0 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.7 Bq/Kg raw
Honey	Iwaki	Unknown	Cs137	12.7 Bq/Kg raw	±	3.0 Bq/Kg raw	16.1	Cs137	3.0 Bq/Kg raw
			Cs134	3.4 Bq/Kg raw	±	1.4 Bq/Kg raw		Cs134	2.7 Bq/Kg raw
Spring water	Iwaki	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	1.8 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	1.7 Bq/Kg raw
School lunch	Iwaki	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.6 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.4 Bq/Kg raw
School lunch	Iwaki	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.1 Bq/Kg raw

※“—” used in Measurement Result and Uncertainty shows that the value is below the detection limit. But it does not necessary mean 0(zero)Bq/Kg.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
			Cs137	Cs134	±	±		Cs137	Cs134
School lunch	Iwaki	Apr-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.0 Bq/Kg raw
School lunch	Iwaki	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.5 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	3.2 Bq/Kg raw
School lunch	Iwaki	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	2.9 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.7 Bq/Kg raw
School lunch	Iwaki	May-15	Cs137	— Bq/Kg raw	±	— Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.2 Bq/Kg raw
			Cs134	— Bq/Kg raw	±	— Bq/Kg raw		Cs134	2.9 Bq/Kg raw
Fireplace ash	Yamagata Pref	Unknown	Cs137	584 Bq/Kg dry	±	117 Bq/Kg dry	785	Cs137	5.5 Bq/Kg dry
			Cs134	201 Bq/Kg dry	±	41.0 Bq/Kg dry		Cs134	5.0 Bq/Kg dry
Fallen leaves	Iwaki	Apr-15	Cs137	362 Bq/Kg dry	±	79.0 Bq/Kg dry	493	Cs137	31.5 Bq/Kg dry
			Cs134	131 Bq/Kg dry	±	34.0 Bq/Kg dry		Cs134	29.2 Bq/Kg dry
Mulberry leaf	Kawamata	Oct-14	Cs137	11.0 Bq/Kg dry	±	5.1 Bq/Kg dry	18.5	Cs137	7.1 Bq/Kg dry
			Cs134	7.5 Bq/Kg dry	±	4.3 Bq/Kg dry		Cs134	6.6 Bq/Kg dry
Mulberry leaf	Miharu	Oct-14	Cs137	— Bq/Kg dry	±	— Bq/Kg dry	7.6	Cs137	8.1 Bq/Kg dry
			Cs134	7.6 Bq/Kg dry	±	4.8 Bq/Kg dry		Cs134	7.4 Bq/Kg dry
Mulberry leaf	Koriyama	Oct-14	Cs137	— Bq/Kg dry	±	— Bq/Kg dry	Under Minimum Limit of Detection	Cs137	4.4 Bq/Kg dry
			Cs134	— Bq/Kg dry	±	— Bq/Kg dry		Cs134	4.0 Bq/Kg dry
Mulberry leaf	Koriyama	Oct-14	Cs137	— Bq/Kg dry	±	— Bq/Kg dry	Under Minimum Limit of Detection	Cs137	6.2 Bq/Kg dry
			Cs134	— Bq/Kg dry	±	— Bq/Kg dry		Cs134	5.8 Bq/Kg dry
Soil	Miharu	Oct-14	Cs137	1100 Bq/Kg dry	±	220 Bq/Kg dry	1477	Cs137	4.5 Bq/Kg dry
			Cs134	377 Bq/Kg dry	±	75.0 Bq/Kg dry		Cs134	4.0 Bq/Kg dry
Soil	Miharu	Oct-14	Cs137	3570 Bq/Kg dry	±	710 Bq/Kg dry	4720	Cs137	4.7 Bq/Kg dry
			Cs134	1150 Bq/Kg dry	±	230 Bq/Kg dry		Cs134	4.2 Bq/Kg dry
Soil	Koriyama	Oct-14	Cs137	494 Bq/Kg dry	±	53.4 Bq/Kg dry	602	Cs137	1.0 Bq/Kg dry
			Cs134	108 Bq/Kg dry	±	16.7 Bq/Kg dry		Cs134	1.0 Bq/Kg dry
Soil	Koriyama	Oct-14	Cs137	1390 Bq/Kg dry	±	280 Bq/Kg dry	1842	Cs137	4.7 Bq/Kg dry
			Cs134	452 Bq/Kg dry	±	90.0 Bq/Kg dry		Cs134	4.4 Bq/Kg dry
Soil	Koriyama	Oct-14	Cs137	642 Bq/Kg dry	±	128 Bq/Kg dry	870	Cs137	4.3 Bq/Kg dry
			Cs134	228 Bq/Kg dry	±	46.0 Bq/Kg dry		Cs134	4.0 Bq/Kg dry
Soil	Koriyama	Oct-14	Cs137	1670 Bq/Kg dry	±	330 Bq/Kg dry	2199	Cs137	4.2 Bq/Kg dry
			Cs134	529 Bq/Kg dry	±	106 Bq/Kg dry		Cs134	3.8 Bq/Kg dry
Soil	Kawamata	Oct-14	Cs137	5041 Bq/Kg dry	±	441 Bq/Kg dry	6174	Cs137	1.0 Bq/Kg dry
			Cs134	1133 Bq/Kg dry	±	110 Bq/Kg dry		Cs134	1.0 Bq/Kg dry
Soil	Kawamata	Oct-14	Cs137	6260 Bq/Kg dry	±	1250 Bq/Kg dry	8290	Cs137	3.2 Bq/Kg dry
			Cs134	2030 Bq/Kg dry	±	410 Bq/Kg dry		Cs134	2.9 Bq/Kg dry
Soil	Shirakawa	Oct-14	Cs137	4340 Bq/Kg dry	±	870 Bq/Kg dry	5750	Cs137	5.6 Bq/Kg dry
			Cs134	1410 Bq/Kg dry	±	280 Bq/Kg dry		Cs134	5.0 Bq/Kg dry
Soil	Shirakawa	Nov-14	Cs137	100 Bq/Kg dry	±	21.0 Bq/Kg dry	145	Cs137	4.0 Bq/Kg dry
			Cs134	44.9 Bq/Kg dry	±	9.8 Bq/Kg dry		Cs134	3.6 Bq/Kg dry
Soil	Shirakawa	Nov-14	Cs137	1890 Bq/Kg dry	±	380 Bq/Kg dry	2493	Cs137	4.6 Bq/Kg dry
			Cs134	603 Bq/Kg dry	±	121 Bq/Kg dry		Cs134	4.2 Bq/Kg dry

※“—” used in Measurement Result and Uncertainty shows that the value is below the detection limit. But it does not necessary mean 0(zero)Bq/Kg.

★Gamma-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Soil	Shirakawa	Nov-14	Cs137	3790 Bq/Kg dry	± 760 Bq/Kg dry	4950	Cs137	4.8 Bq/Kg dry
			Cs134	1160 Bq/Kg dry	± 230 Bq/Kg dry		Cs134	4.4 Bq/Kg dry
Soil	Shirakawa	Nov-14	Cs137	162 Bq/Kg dry	± 32.0 Bq/Kg dry	223	Cs137	4.2 Bq/Kg dry
			Cs134	61.0 Bq/Kg dry	± 12.9 Bq/Kg dry		Cs134	4.0 Bq/Kg dry
Soil	Ishikawa	Nov-14	Cs137	360 Bq/Kg dry	± 72.0 Bq/Kg dry	496	Cs137	4.2 Bq/Kg dry
			Cs134	136 Bq/Kg dry	± 27.0 Bq/Kg dry		Cs134	3.9 Bq/Kg dry
Soil	Ishikawa	Nov-14	Cs137	407 Bq/Kg dry	± 81.0 Bq/Kg dry	555	Cs137	4.7 Bq/Kg dry
			Cs134	148 Bq/Kg dry	± 30.0 Bq/Kg dry		Cs134	4.2 Bq/Kg dry
Soil	Hirata	Nov-14	Cs137	316 Bq/Kg dry	± 37.9 Bq/Kg dry	387	Cs137	1.0 Bq/Kg dry
			Cs134	71.1 Bq/Kg dry	± 12.7 Bq/Kg dry		Cs134	1.0 Bq/Kg dry
Soil	Iwaki	Apr-15	Cs137	3.4 Bq/Kg dry	± 1.1 Bq/Kg dry	18.1	Cs137	1.0 Bq/Kg dry
			Cs134	14.7 Bq/Kg dry	± 3.0 Bq/Kg dry		Cs134	1.0 Bq/Kg dry
Soil	Iwaki	Apr-15	Cs137	343 Bq/Kg dry	± 69.0 Bq/Kg dry	467	Cs137	2.5 Bq/Kg dry
			Cs134	124 Bq/Kg dry	± 25.0 Bq/Kg dry		Cs134	2.2 Bq/Kg dry
Soil	Iwaki	Apr-15	Cs137	8.3 Bq/Kg dry	± 1.9 Bq/Kg dry	18.8	Cs137	0.9 Bq/Kg dry
			Cs134	10.5 Bq/Kg dry	± 2.2 Bq/Kg dry		Cs134	0.8 Bq/Kg dry
Soil	Koriyama	Apr-15	Cs137	71.6 Bq/Kg dry	± 13.5 Bq/Kg dry	81.6	Cs137	1.0 Bq/Kg dry
			Cs134	10.0 Bq/Kg dry	± 6.8 Bq/Kg dry		Cs134	1.0 Bq/Kg dry
Soil	Koriyama	Apr-15	Cs137	202 Bq/Kg dry	± 25.5 Bq/Kg dry	244	Cs137	1.0 Bq/Kg dry
			Cs134	41.2 Bq/Kg dry	± 9.3 Bq/Kg dry		Cs134	1.0 Bq/Kg dry

※“_” used in Measurement Result and Uncertainty shows that the value is below the detection limit. But it does not necessary mean 0(zero)Bq/Kg.



★Beta-ray

(Bq/Kg raw:raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty		Minimum Limit of Detection	
Bamboo shoots (boiled 3 times)	Iwaki	Apr-15	T(Organization)	Under Minimum Limit of Detection	Bq/Kg dry	±	—	Bq/Kg dry	2.10 Bq/Kg dry
			Sr90	Under Minimum Limit of Detection	Bq/Kg raw	±	—	Bq/Kg raw	0.06 Bq/Kg raw
Rapeseed oil	Minami-soma	Unknown	T(Organization)	Under Minimum Limit of Detection	Bq/Kg raw	±	—	Bq/Kg raw	3.50 Bq/Kg raw
			Sr90	Under Minimum Limit of Detection	Bq/Kg raw	±	—	Bq/Kg raw	0.05 Bq/Kg raw
Spirulina	Hawaii	Unknown	T(Organization)	Under Minimum Limit of Detection	Bq/Kg dry	±	—	Bq/Kg dry	4.10 Bq/Kg dry
			Sr90	Under Minimum Limit of Detection	Bq/Kg raw	±	—	Bq/Kg raw	0.22 Bq/Kg raw
Fuki (Edible wild plant)	Iwaki	Apr-15	T(Free)	Under Minimum Limit of Detection	Bq/L	±	—	Bq/L	4.40 Bq/L
			T(Organization)	5.70	Bq/Kg dry	±	2.70	Bq/Kg dry	2.50 Bq/Kg dry
			Sr90	1.34	Bq/Kg raw	±	0.47	Bq/Kg raw	0.45 Bq/Kg raw
Fuki leaf (Edible wild plant)	Iwaki	Apr-15	Sr90	1.73	Bq/Kg raw	±	0.49	Bq/Kg raw	0.47 Bq/Kg raw
Mulberry leaf	Kawamata	Oct-14	Sr90	7.99	Bq/Kg raw	±	0.14	Bq/Kg raw	0.04 Bq/Kg raw
Fallen leaves	Iwaki	Apr-15	T(Organization)	3.80	Bq/Kg dry	±	1.80	Bq/Kg dry	1.90 Bq/Kg dry
			Sr90	0.16	Bq/Kg raw	±	0.08	Bq/Kg raw	0.08 Bq/Kg raw

T(Free) : Tritium(Free water) T(Organization) : Tritium(Organization bound water) Sr90 : Strontium90

※The value below Minimum Limit of Detection does not necessary
mean 0(zero)Bq/Kg.

