



Radiation Measurement Results of 127 Items in August



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Potato	Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Potato	Ouse, Koriyama	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Potato	Tairasimotakaku, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Potato	Ueda, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Burdock	Ibaraki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Burdock	Gunma	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Japanese white radish	Aomori	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Eggplant	Tono, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Green pepper	Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.7 Bq/kg raw
Green pepper	Nishiki, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Japanese mustard spinach	Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Malabar spinach	Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Moroccan green bean	Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.1 Bq/kg raw
Asparagus	Aizu, Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.1 Bq/kg raw
Cucumber	Tono, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Cucumber	Nishiki, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.9 Bq/kg raw
Cucumber	Yonezawa, Yamagata	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Bitter gourd	Nihonmatsu, Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Bitter gourd	unknown	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Zucchini	Yonezawa, Yamagata	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw

*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessarily mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Zucchini	Nagano	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.0 Bq/kg raw
Pumpkin	Nihonmatsu, Fukushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Pumpkin	Joban, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Spaghetti squash	Yonezawa, Yamagata	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Green pepper	Tono, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Perilla	Hitachiota, Ibaraki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 5.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 4.6 Bq/kg raw
Green chili	Hiratamura, Ishikawa	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Corn(peel)	Ibaraki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 8.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 6.6 Bq/kg raw
Corn(core)	Ibaraki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.7 Bq/kg raw
Tomato	Fukushima, Fukushima	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Tomato	Tono, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Citron	Izumigaoka, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Water melon	Tono, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Pear	Yamagata	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Blackberry	Tairashimokabeya, Iwaki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Blueberry	Tairashimokabeya, Iwaki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.0 Bq/kg raw
Blueberry	Tono, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Blueberry	Joban, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.2 Bq/kg raw
Nameko mushroom	Fukushima	Aug-18	Cs137	13.2 Bq/kg raw	± 3.0 Bq/kg raw	13.2	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Elingi mushroom	Ogawa, Iwaki	Aug-18	Cs137	2.2 Bq/kg raw	± 1.4 Bq/kg raw	2.2	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Shimeji mushroom	Ibaraki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 7.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 5.9 Bq/kg raw
Dried bracken	Yonezawa, Yamagata	Aug-18	Cs137	5.6 Bq/kg raw	± 2.0 Bq/kg raw	5.6	Cs137 2.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.5 Bq/kg raw
Egg	Hiratamura, Ishikawa	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.0 Bq/kg raw
Tofu	Naruto, Tokushima	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw

*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Soy milk	Fukuoka	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Roundnose flounder (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 4.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 3.3 Bq/kg raw
Roundnose flounder (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Roundnose flounder (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Greenling (flesh)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Greenling (head, bone, guts)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Young yellowtail (flesh)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Young yellowtail (head, bone, guts)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 4.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 3.0 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.7 Bq/kg raw
Red rockfish (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Littlemouth flounder (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Roundnose flounder (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Roundnose flounder (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Smelt	Ibaraki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Carp (whole)	Yamagata	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Smelt	Ibaraki	Jul-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Basket clam	Taira, Iwaki	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Ascidian	Ishinomaki, Miyagi	unknown	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.0 Bq/kg raw
Hijiki seaweed	Chiba	Aug-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw

*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result	Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Sea water A (surface)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137 — Bq/L	± — Bq/L	Under Minimum Limit of Detection	Cs137 0.016 Bq/L
			Cs134 — Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water A (lower)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137 0.076 Bq/L	± 0.011 Bq/L	0.076	Cs137 0.018 Bq/L
			Cs134 — Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water B (surface)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137 — Bq/L	± — Bq/L	Under Minimum Limit of Detection	Cs137 0.017 Bq/L
			Cs134 — Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water B (lower)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137 0.047 Bq/L	± 0.011 Bq/L	0.047	Cs137 0.017 Bq/L
			Cs134 — Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137 — Bq/L	± — Bq/L	Under Minimum Limit of Detection	Cs137 0.016 Bq/L
			Cs134 — Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water C (lower)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Cs137 — Bq/L	± — Bq/L	Under Minimum Limit of Detection	Cs137 0.017 Bq/L
			Cs134 — Bq/L	± — Bq/L		Cs134 — Bq/L
School lunch	Uchigotakasaka, Iwaki	Aug-18	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.8 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
School lunch	Jobanmatsugadai, Iwaki	Aug-18	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Canned coffee	unknown	unknown	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Milk	Ibaraki	Aug-18	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.8 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.7 Bq/kg raw
Yogurt	Gunma	Aug-18	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Somen (dried noodles)	Nagasaki	2018	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Pine leaves	Tokaimura, Naka , Ibaraki	Jul-18	Cs137 227.0 Bq/kg raw	± 45.0 Bq/kg raw	255.2	Cs137 9.0 Bq/kg raw
			Cs134 28.2 Bq/kg raw	± 7.6 Bq/kg raw		Cs134 8.1 Bq/kg raw
Mulberry leaves	Mayumi, Hitachiota, Ibaraki	Jul-18	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.4 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 1.8 Bq/kg raw
Fallen leaves	Okuma, Futaba	Aug-18	Cs137 28700.0 Bq/kg raw	± 5700.0 Bq/kg raw	31640.0	Cs137 61.8 Bq/kg raw
			Cs134 2940.0 Bq/kg raw	± 590.0 Bq/kg raw		Cs134 48.6 Bq/kg raw
Weed①	Okuma, Futaba	Aug-18	Cs137 1230.0 Bq/kg raw	± 250.0 Bq/kg raw	1387.0	Cs137 41.1 Bq/kg raw
			Cs134 157.0 Bq/kg raw	± 39.0 Bq/kg raw		Cs134 33.5 Bq/kg raw
Weed · Branch	Okuma, Futaba	Aug-18	Cs137 515.0 Bq/kg raw	± 106.0 Bq/kg raw	571.5	Cs137 47.1 Bq/kg raw
			Cs134 56.5 Bq/kg raw	± 28.4 Bq/kg raw		Cs134 46.4 Bq/kg raw
Weed	Onahamahanabatake, Iwaki	Aug-18	Cs137 145.4 Bq/kg raw	± 14.7 Bq/kg raw	157.5	Cs137 3.8 Bq/kg raw
			Cs134 12.1 Bq/kg raw	± 3.5 Bq/kg raw		Cs134 3.4 Bq/kg raw
Swallow's nest	Onahama, Iwaki	2012	Cs137 405.0 Bq/kg raw	± 81.0 Bq/kg raw	466.7	Cs137 2.9 Bq/kg raw
			Cs134 61.7 Bq/kg raw	± 12.3 Bq/kg raw		Cs134 2.5 Bq/kg raw
Moss	Joban, Iwaki	Aug-18	Cs137 1170.0 Bq/kg raw	± 230.0 Bq/kg raw	1329.0	Cs137 17.8 Bq/kg raw
			Cs134 159.0 Bq/kg raw	± 34.0 Bq/kg raw		Cs134 16.6 Bq/kg raw
Leaf mold	Okuma, Futaba	Aug-18	Cs137 64700.0 Bq/kg dry	± 12900.0 Bq/kg dry	72770.0	Cs137 52.6 Bq/kg dry
			Cs134 8070.0 Bq/kg dry	± 1610.0 Bq/kg dry		Cs134 50.8 Bq/kg dry
Soil	Okuma, Futaba	Aug-18	Cs137 495000.0 Bq/kg dry	± 99000.0 Bq/kg dry	556500.0	Cs137 155.0 Bq/kg dry
			Cs134 61500.0 Bq/kg dry	± 12300.0 Bq/kg dry		Cs134 151.0 Bq/kg dry
Soil	Okuma, Futaba	Aug-18	Cs137 31600.0 Bq/kg dry	± 6300.0 Bq/kg dry	35320.0	Cs137 68.4 Bq/kg dry
			Cs134 3720.0 Bq/kg dry	± 740.0 Bq/kg dry		Cs134 67.3 Bq/kg dry
Soil	Okuma, Futaba	Aug-18	Cs137 25900.0 Bq/kg dry	± 5200.0 Bq/kg dry	28500.0	Cs137 17.6 Bq/kg dry
			Cs134 2600.0 Bq/kg dry	± 520.0 Bq/kg dry		Cs134 14.0 Bq/kg dry

*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Soil	Izumi,Fukushima	Aug-18	Cs137	3180.0	Bq/kg dry ± 344.0	3535.0	Cs137	12.2 Bq/kg dry
			Cs134	355.0	Bq/kg dry ± 45.5		Cs134	12.3 Bq/kg dry
Soil	Hisanohama, Iwaki	Aug-18	Cs137	1610.0	Bq/kg dry ± 174.0	1794.0	Cs137	11.0 Bq/kg dry
			Cs134	184.0	Bq/kg dry ± 23.7		Cs134	13.2 Bq/kg dry
Soil	Ohara, Iwaki	Aug-18	Cs137	383.0	Bq/kg dry ± 42.5	419.1	Cs137	5.9 Bq/kg dry
			Cs134	36.1	Bq/kg dry ± 6.0		Cs134	7.8 Bq/kg dry
Soil	Ohara, Iwaki	Aug-18	Cs137	51.6	Bq/kg dry ± 6.7	59.6	Cs137	4.9 Bq/kg dry
			Cs134	8.0	Bq/kg dry ± 2.1		Cs134	7.7 Bq/kg dry
Soil	Ohara, Iwaki	Aug-18	Cs137	45.8	Bq/kg dry ± 5.7	45.8	Cs137	3.5 Bq/kg dry
			Cs134	—	Bq/kg dry ± —		Cs134	4.1 Bq/kg dry
Soil	Joban, Iwaki	Aug-18	Cs137	123.0	Bq/kg dry ± 14.5	135.5	Cs137	4.3 Bq/kg dry
			Cs134	12.5	Bq/kg dry ± 2.4		Cs134	5.4 Bq/kg dry
Soil	Shimogawa, Izumi, Iwaki	Aug-18	Cs137	301.0	Bq/kg dry ± 34.1	337.8	Cs137	4.1 Bq/kg dry
			Cs134	36.8	Bq/kg dry ± 6.0		Cs134	6.1 Bq/kg dry
Soil	Joban, Iwaki	Aug-18	Cs137	123.0	Bq/kg dry ± 14.5	135.5	Cs137	4.3 Bq/kg dry
			Cs134	12.5	Bq/kg dry ± 2.4		Cs134	5.4 Bq/kg dry
Soil	Ogawa, Iwaki	Aug-18	Cs137	44700.0	Bq/kg dry ± 4860.0	49620.0	Cs137	55.8 Bq/kg dry
			Cs134	4920.0	Bq/kg dry ± 636.0		Cs134	50.2 Bq/kg dry
Soil	Ogawa, Iwaki	Aug-18	Cs137	32900.0	Bq/kg dry ± 3580.0	36490.0	Cs137	51.0 Bq/kg dry
			Cs134	3590.0	Bq/kg dry ± 464.0		Cs134	50.6 Bq/kg dry
Soil	Ogawa, Iwaki	Aug-18	Cs137	5310.0	Bq/kg dry ± 573.0	5879.0	Cs137	19.0 Bq/kg dry
			Cs134	569.0	Bq/kg dry ± 73.1		Cs134	19.1 Bq/kg dry
Soil	Ogawa, Iwaki	Aug-18	Cs137	5200.0	Bq/kg dry ± 576.0	5707.0	Cs137	16.8 Bq/kg dry
			Cs134	507.0	Bq/kg dry ± 72.9		Cs134	17.2 Bq/kg dry
Soil	Ogawa, Iwaki	Aug-18	Cs137	3150.0	Bq/kg dry ± 338.0	3482.0	Cs137	17.4 Bq/kg dry
			Cs134	332.0	Bq/kg dry ± 42.5		Cs134	18.6 Bq/kg dry
Soil	Sakura, Tochigi	Aug-18	Cs137	203.0	Bq/kg dry ± 23.1	223.2	Cs137	3.9 Bq/kg dry
			Cs134	20.2	Bq/kg dry ± 3.9		Cs134	6.0 Bq/kg dry
Soil	Tokaimura, Naka , Ibaraki	Aug-18	Cs137	852.0	Bq/kg dry ± 97.8	949.0	Cs137	6.7 Bq/kg dry
			Cs134	97.0	Bq/kg dry ± 15.1		Cs134	7.6 Bq/kg dry
Soil	Hitachiota, Ibaraki	Aug-18	Cs137	420.0	Bq/kg dry ± 47.7	463.4	Cs137	8.0 Bq/kg dry
			Cs134	43.4	Bq/kg dry ± 7.1		Cs134	11.1 Bq/kg dry
Soil	Tokaimura, Naka , Ibaraki	Aug-18	Cs137	179.0	Bq/kg dry ± 20.7	197.6	Cs137	5.1 Bq/kg dry
			Cs134	18.6	Bq/kg dry ± 3.3		Cs134	7.4 Bq/kg dry
Soil	Tokaimura, Naka , Ibaraki	Aug-18	Cs137	81.4	Bq/kg dry ± 10.1	89.8	Cs137	5.7 Bq/kg dry
			Cs134	8.4	Bq/kg dry ± 2.0		Cs134	5.7 Bq/kg dry
Soil	Tokaimura, Naka , Ibaraki	Aug-18	Cs137	76.4	Bq/kg dry ± 9.4	84.7	Cs137	5.1 Bq/kg dry
			Cs134	8.3	Bq/kg dry ± 2.1		Cs134	7.6 Bq/kg dry
Vacuum cleaner dust	Onahamahanabatake, Iwaki	Aug-18	Cs137	612.0	Bq/kg raw ± 122.0	687.4	Cs137	3.4 Bq/kg raw
			Cs134	75.4	Bq/kg raw ± 15.1		Cs134	3.1 Bq/kg raw

*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



★Beta-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Minimum Limit of Detection	
Flounder(flesh)	Off the coast of Fukushima Nuclear Power Plant1	Jul-17	T(Organization)	Under Minimum Limit of Detection Bq/Kg dry	± —	Bq/Kg dry	1.37 Bq/Kg dry
Sea water A (surface)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Free)	Under Minimum Limit of Detection Bq/L	± —	Bq/L	1.98 Bq/L
Sea water A (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Free)	Under Minimum Limit of Detection Bq/L	± —	Bq/L	1.98 Bq/L
Sea water B (surface)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Free)	Under Minimum Limit of Detection Bq/L	± —	Bq/L	1.98 Bq/L
Sea water B (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Free)	Under Minimum Limit of Detection Bq/L	± —	Bq/L	1.98 Bq/L
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Free)	Under Minimum Limit of Detection Bq/L	± —	Bq/L	1.98 Bq/L
Sea water C (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Free)	Under Minimum Limit of Detection Bq/L	± —	Bq/L	1.98 Bq/L
Honey	Naraha,Futaba	May-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± —	Bq/Kg dry	0.33 Bq/Kg dry
Greenling (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jun-16	Sr90	1.02 Bq/Kg dry	± 0.10	Bq/Kg dry	0.14 Bq/Kg dry
Greenling (whole)	Off the coast of Fukushima Nuclear Power Plant1	Jun-16	Sr90	0.53 Bq/Kg dry	± 0.15	Bq/Kg dry	0.22 Bq/Kg dry
Tuna(bone)	Fukushima	Jul-18	Sr90	0.27 Bq/Kg dry	± 0.14	Bq/Kg dry	0.21 Bq/Kg dry
Sea water A (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	Sr90	Under Minimum Limit of Detection Bq/L	± —	Bq/L	0.0014 Bq/L
Sea water B (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	Sr90	0.0013 Bq/L	± 0.0008	Bq/L	0.0013 Bq/L
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	Sr90	Under Minimum Limit of Detection Bq/L	± —	Bq/L	0.0011 Bq/L
Sea water C (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	Sr90	Under Minimum Limit of Detection Bq/L	± —	Bq/L	0.0006 Bq/L
Sea water A (surface)	Off the coast of Fukushima Nuclear Power Plant1	Jul-18	Sr90	Under Minimum Limit of Detection Bq/L	± —	Bq/L	0.0010 Bq/L

T(Free) : Tritium(Free water) T(Organization) : Tritium(Organization bound water) Sr90 : Strontium90

※The value below Minimum Limit of Detection does not necessary mean 0(zero)Bq/Kg.

