



# Radiation Measurement Results of 171 Items in November



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Brown rice	Obama, Iwaki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Brown rice	Tairakoizumi, Iwaki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Brown rice	Yamagata	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Rice	Tairanakakabeya, Iwaki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Rice	Joban, Iwaki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.7 Bq/kg raw
Rice	Ishikawa, Ishikawa	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Potato	Kashima, Minamisoma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Potato (with peel)	Joban, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Taro(peel)	Ibaraki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.8 Bq/kg raw
Yam	Hirata, Ishikawa	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Yam(with peel)	Izumigaoka, Iwaki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Sweet potato	Takiziri, Izumi, Iwaki	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Sweet potato	Nakoso, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Japanese white radish(pulp)	Kashima, Minamisoma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Japanese white radish(leaf)	Kashima, Minamisoma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Japanese white radish(pulp)	Soma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Japanese white radish(leaf)	Soma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Japanese white radish	Hirata, Ishikawa	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Turnip (with leaf)	Okinouti, Soma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Turnip(pulp)	Joban, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.7 Bq/kg raw

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessarily mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Turnip (leaf)	Joban, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.7 Bq/kg raw
Turnip (calyx, peel))	Joban, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 3.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.4 Bq/kg raw
Red turnip	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Cabbage	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Chinese cabbage	Fukushima	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Chinese cabbage	Kashima, Minamisoma	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Chinese cabbage	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Japanese mustard spinach	Kashima, Minamisoma	Nov-18	Cs137	5.4 Bq/kg raw	± 2.7 Bq/kg raw	5.4	Cs137 3.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.5 Bq/kg raw
Garland chrysanthemum	Otama, Adachi	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Garland chrysanthemum	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Mustard greens	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Broccoli	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Yacon	Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Pumpkin	Hirono, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Chayote	Otama, Adachi	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Wax gourd (seed, cotton)	Shimogo, Minamiaizu	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Wax gourd(peel)	Shimogo, Minamiaizu	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.2 Bq/kg raw
Citron	Kashima, Minamisoma	Nov-18	Cs137	5.4 Bq/kg raw	± 1.3 Bq/kg raw	5.4	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Citron	Izumigaoka, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Citron	Kashima, Iwaki	Nov-18	Cs137	6.5 Bq/kg raw	± 1.6 Bq/kg raw	6.5	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Citron	Onahamakamikaziro, Iwaki	Nov-18	Cs137	2.1 Bq/kg raw	± 0.9 Bq/kg raw	2.1	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Citron	Nishiki, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Astringent persimmon(pulp)	Fukushima	Nov-18	Cs137	2.5 Bq/kg raw	± 0.9 Bq/kg raw	2.5	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Persimmon(pulp)	Idate, Soma	Nov-18	Cs137	1.9 Bq/kg raw	± 0.9 Bq/kg raw	1.9	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Persimmon (peel, calyx)	Idate, Soma	Nov-18	Cs137	3.5	Bq/kg raw ± 1.8 Bq/kg raw	3.5	Cs137 2.3 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.8 Bq/kg raw
Persimmon(pulp)	Kashima, Minamisoma	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Persimmon (peel, calyx)	Kashima, Minamisoma	Nov-18	Cs137	2.6	Bq/kg raw ± 1.0 Bq/kg raw	2.6	Cs137 1.2 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Persimmon(pulp)	Nakamura, Soma	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.0 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 0.9 Bq/kg raw
Persimmon(pulp)	Otama, Adachi	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Persimmon (peel, calyx)	Otama, Adachi	Oct-18	Cs137	4.2	Bq/kg raw ± 1.9 Bq/kg raw	4.2	Cs137 2.5 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Persimmon (with peel)	Izumigaoka, Iwa ki	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Persimmon(pulp)	Onahama- hanabatake, Iwaki	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Persimmon (peel, seed)	Onahama- hanabatake, Iwaki	Oct-18	Cs137	4.1	Bq/kg raw ± 2.3 Bq/kg raw	4.1	Cs137 2.2 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.7 Bq/kg raw
Astringent persimmon (peel, seed)	Ena, Iwaki	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.4 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.8 Bq/kg raw
Apple(with peel)	Fukushima	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Apple(pulp)	Fukushima	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Apple(peel, calyx )	Fukushima	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.3 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.8 Bq/kg raw
Apple(pulp)	Fukushima	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Mandarin orange(pulp)	Onahama- okaona, Iwaki	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Mandarin orange(peel)	Onahama- okaona, Iwaki	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.5 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Mandarin orange(pulp)	Onahamakamikaziro, Iwaki	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Mandarin orange(peel)	Onahamakamikaziro, Iwaki	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.8 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 2.2 Bq/kg raw
Mandarin orange(pulp)	Atami, Shizuoka	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Mandarin orange(peel)	Atami, Shizuoka	Oct-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.6 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 2.0 Bq/kg raw
Chinese quince	Idate, Soma	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Chinese quince	Yokodai, Iwaki	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Kiwi fruit	Nakamura, Soma	Nov-18	Cs137	2.0	Bq/kg raw ± 0.7 Bq/kg raw	2.0	Cs137 1.0 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 0.9 Bq/kg raw
Kiwi fruit	Kashima, Iwaki	Nov-18	Cs137	—	Bq/kg raw ± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 1.4 Bq/kg raw

※"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Pear(pulp)	Yamagata	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Pear(peel)	Yamagata	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.9 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	13.1 Bq/kg raw	± 3.7 Bq/kg raw	13.1	Cs137 4.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 3.3 Bq/kg raw
Fox jacopever (flesh)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	4.1 Bq/kg raw	± 1.4 Bq/kg raw	4.1	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Fox jacopever (head)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Fox jacopever (bone)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.8 Bq/kg raw
Fox jacopever (guts)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	5.5 Bq/kg raw	± 3.1 Bq/kg raw	5.5	Cs137 4.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 3.1 Bq/kg raw
Fox jacopever (whole)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Goldeye rockfish (whole)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	2.1 Bq/kg raw	± 1.9 Bq/kg raw	2.1	Cs137 2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.7 Bq/kg raw
Goldeye rockfish (whole)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Yellowtail(whole )	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	2.5 Bq/kg raw	± 1.2 Bq/kg raw	2.5	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Shark(flesh)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	8.6 Bq/kg raw	± 1.9 Bq/kg raw	8.6	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Shark(head)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	3.2 Bq/kg raw	± 1.1 Bq/kg raw	3.2	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Shark(bone, skin)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	8.2 Bq/kg raw	± 2.3 Bq/kg raw	8.2	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Shark(guts)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	3.7 Bq/kg raw	± 1.3 Bq/kg raw	3.7	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Salmon(whole)	Hirono, Futaba	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Mackerel (small fish)	Off the coast of Onahama, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Cod(whole)	Otuchi, Kamihei , Iwate	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.9 Bq/kg raw
Groundwater	Futaba, Futaba	Nov-18	Cs137	— Bq/L	± — Bq/L	Under Minimum Limit of Detection	Cs137 0.9 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 0.8 Bq/L
River water	Kashima, Minamisoma	Nov-18	Cs137	0.022 Bq/L	± 0.010 Bq/L	0.022	Cs137 0.016 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water A (surface)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	0.029 Bq/L	± 0.011 Bq/L	0.029	Cs137 0.017 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water A (lower)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	0.035 Bq/L	± 0.010 Bq/L	0.035	Cs137 0.017 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water B (surface)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	0.028 Bq/L	± 0.010 Bq/L	0.028	Cs137 0.017 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water B (lower)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	0.020 Bq/L	± 0.010 Bq/L	0.020	Cs137 0.017 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L

※"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	0.023 Bq/L	± 0.010 Bq/L	<b>0.023</b>	Cs137 0.017 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L
Sea water C (lower)	Off the coast of Fukushima Nuclear Power Plant2	Oct-18	Cs137	0.023 Bq/L	± 0.010 Bq/L	<b>0.023</b>	Cs137 0.017 Bq/L
			Cs134	— Bq/L	± — Bq/L		Cs134 — Bq/L
Dried sweet potapo	Ibaraki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Salted plum	Otama, Adachi	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Soybeans	Fushiguro, Date	Oct-18	Cs137	1.0 Bq/kg raw	± 0.6 Bq/kg raw	<b>1.0</b>	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Green soybean	Yamagata	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.6 Bq/kg raw
Barley	Japan (production)	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.5 Bq/kg raw
Bran	unknown	2018	Cs137	1.6 Bq/kg raw	± 0.9 Bq/kg raw	<b>1.6</b>	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Amazake (fermented rice drink)	Aizuwakamatu	unknown	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Oyster mushroom	Otama, Adachi	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Shitake mushroom grown in log	Tairashimoyamaguti, Iwaki	Nov-18	Cs137	137.0 Bq/kg raw	± 27.0 Bq/kg raw	<b>151.6</b>	Cs137 1.9 Bq/kg raw
			Cs134	14.6 Bq/kg raw	± 3.2 Bq/kg raw		Cs134 1.7 Bq/kg raw
Shitake mushroom grown in log	Kawasaki, Kanagawa	Nov-18	Cs137	3.6 Bq/kg raw	± 1.0 Bq/kg raw	<b>3.6</b>	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.2 Bq/kg raw
Shitake mushroom grown in log	Kumamoto	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Honey	Fushiguro, Date	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Soy sauce	Ogawa, Iwaki	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Rice miso	Ogawa, Iwaki	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw
Salt	China (production)	unknown	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Beet sugar	Hokkaido	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.0 Bq/kg raw
Bran	unknown	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Green tea	Shizuoka	2018	Cs137	5.1 Bq/kg raw	± 2.5 Bq/kg raw	<b>5.1</b>	Cs137 3.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.3 Bq/kg raw
Green tea	Shizuoka	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 2.0 Bq/kg raw
Yogurt	Koriyama	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
Green tea (petbottle)	Japan (production)	2018	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Orange juice	Japan (production)	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.1 Bq/kg raw

※"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Grape juice	Japan (production)	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.4 Bq/kg raw
School lunch	Uchigotakasaka, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.7 Bq/kg raw
School lunch	Uchigotakasaka, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.7 Bq/kg raw
School lunch	Jobanmatsugadai, Iwaki	Nov-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 0.8 Bq/kg raw
Beeswax	Fushiguro, Date	Oct-18	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 1.3 Bq/kg raw
Fallen leaves	Hirono, Iwaki	Nov-18	Cs137	79.2 Bq/kg raw	± 16.8 Bq/kg raw	79.2	Cs137 8.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 7.9 Bq/kg raw
Pine leaves	Okuma, Futaba	Oct-18	Cs137	23000.0 Bq/kg raw	± 4600.0 Bq/kg raw	25590.0	Cs137 61.7 Bq/kg raw
			Cs134	2590.0 Bq/kg raw	± 520.0 Bq/kg raw		Cs134 60.8 Bq/kg raw
Fruits of heavenly bamboo	Okuma, Futaba	Oct-18	Cs137	2900.0 Bq/kg raw	± 580.0 Bq/kg raw	3218.0	Cs137 139.0 Bq/kg raw
			Cs134	318.0 Bq/kg raw	± 106.0 Bq/kg raw		Cs134 133.0 Bq/kg raw
Leaves	Okuma, Futaba	Oct-18	Cs137	1035.0 Bq/kg raw	± 139.1 Bq/kg raw	1161.0	Cs137 85.4 Bq/kg raw
			Cs134	126.0 Bq/kg raw	± 57.9 Bq/kg raw		Cs134 76.6 Bq/kg raw
Ash(wood)	Tairashimokabeya, Iwaki	Nov-18	Cs137	840.0 Bq/kg raw	± 168.0 Bq/kg raw	959.0	Cs137 3.1 Bq/kg raw
			Cs134	119.0 Bq/kg raw	± 24.0 Bq/kg raw		Cs134 2.8 Bq/kg raw
Moss	Kashima, Minamisoma	Nov-18	Cs137	16300.0 Bq/kg raw	± 3300.0 Bq/kg raw	18100.0	Cs137 27.5 Bq/kg raw
			Cs134	1800.0 Bq/kg raw	± 360.0 Bq/kg raw		Cs134 26.0 Bq/kg raw
Moss	Nishiki, Iwaki	Nov-18	Cs137	1240.0 Bq/kg raw	± 250.0 Bq/kg raw	1373.0	Cs137 6.4 Bq/kg raw
			Cs134	133.0 Bq/kg raw	± 27.0 Bq/kg raw		Cs134 6.0 Bq/kg raw
Moss	Iritono, Tono, Iwaki	Oct-18	Cs137	80.5 Bq/kg raw	± 16.1 Bq/kg raw	95.7	Cs137 3.1 Bq/kg raw
			Cs134	15.2 Bq/kg raw	± 3.5 Bq/kg raw		Cs134 2.4 Bq/kg raw
Soil	Idate, Soma	Nov-18	Cs137	7140.0 Bq/Kg dry	± 772.0 Bq/Kg dry	7879.0	Cs137 12.1 Bq/Kg dry
			Cs134	739.0 Bq/Kg dry	± 94.8 Bq/Kg dry		Cs134 11.0 Bq/Kg dry
Soil	Idate, Soma	Nov-18	Cs137	4750.0 Bq/Kg dry	± 513.0 Bq/Kg dry	5236.0	Cs137 10.5 Bq/Kg dry
			Cs134	486.0 Bq/Kg dry	± 62.5 Bq/Kg dry		Cs134 9.7 Bq/Kg dry
Soil	Kashima, Minamisoma	Nov-18	Cs137	1600.0 Bq/Kg dry	± 175.0 Bq/Kg dry	1771.0	Cs137 7.0 Bq/Kg dry
			Cs134	171.0 Bq/Kg dry	± 22.6 Bq/Kg dry		Cs134 7.8 Bq/Kg dry
Soil	Kashima, Minamisoma	Nov-18	Cs137	1350.0 Bq/Kg dry	± 147.0 Bq/Kg dry	1497.0	Cs137 7.6 Bq/Kg dry
			Cs134	147.0 Bq/Kg dry	± 19.4 Bq/Kg dry		Cs134 9.3 Bq/Kg dry
Soil	Kashima, Minamisoma	Nov-18	Cs137	1005.0 Bq/Kg dry	± 116.0 Bq/Kg dry	1120.0	Cs137 7.0 Bq/Kg dry
			Cs134	115.0 Bq/Kg dry	± 15.7 Bq/Kg dry		Cs134 8.6 Bq/Kg dry
Soil	Kashima, Minamisoma	Nov-18	Cs137	895.0 Bq/Kg dry	± 95.8 Bq/Kg dry	980.9	Cs137 7.3 Bq/Kg dry
			Cs134	85.9 Bq/Kg dry	± 11.4 Bq/Kg dry		Cs134 8.6 Bq/Kg dry
Soil	Soma	Nov-18	Cs137	191.0 Bq/Kg dry	± 21.7 Bq/Kg dry	207.6	Cs137 3.8 Bq/Kg dry
			Cs134	16.6 Bq/Kg dry	± 3.1 Bq/Kg dry		Cs134 5.2 Bq/Kg dry
Soil	Takiziri, Izumi , Iwaki	Oct-18	Cs137	68.1 Bq/Kg dry	± 8.1 Bq/Kg dry	75.9	Cs137 3.7 Bq/Kg dry
			Cs134	7.8 Bq/Kg dry	± 1.8 Bq/Kg dry		Cs134 4.7 Bq/Kg dry
Soil	Saitama, Saitama	Nov-18	Cs137	227.0 Bq/Kg dry	± 26.2 Bq/Kg dry	246.6	Cs137 5.6 Bq/Kg dry
			Cs134	19.6 Bq/Kg dry	± 3.7 Bq/Kg dry		Cs134 8.3 Bq/Kg dry
Soil	Saitama, Saitama	Nov-18	Cs137	134.0 Bq/Kg dry	± 15.7 Bq/Kg dry	144.2	Cs137 5.6 Bq/Kg dry
			Cs134	10.2 Bq/Kg dry	± 2.4 Bq/Kg dry		Cs134 8.1 Bq/Kg dry
Soil	Kawasaki, Kanagawa	Nov-18	Cs137	79.6 Bq/Kg dry	± 9.8 Bq/Kg dry	86.7	Cs137 1.6 Bq/Kg dry
			Cs134	7.1 Bq/Kg dry	± 2.2 Bq/Kg dry		Cs134 2.0 Bq/Kg dry

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Volcanic ash soil	Aso, Kumamoto (Mt.Aso)	Oct-18	Cs137	—	Bq/Kg dry ± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137 1.8 Bq/Kg dry
			Cs134	—	Bq/Kg dry ± — Bq/Kg dry		Cs134 1.7 Bq/Kg dry
Vacuum cleaner dust (Dyson)	Kashima, Minamisoma	Nov-18	Cs137	2730.0	Bq/kg raw ± 550.0 Bq/kg raw	2996.0	Cs137 58.1 Bq/kg raw
			Cs134	266.0	Bq/kg raw ± 62.0 Bq/kg raw		Cs134 47.6 Bq/kg raw
Vacuum cleaner dust	Joban, Iwaki	Nov-18	Cs137	450.3	Bq/kg raw ± 46.1 Bq/kg raw	482.2	Cs137 14.0 Bq/kg raw
			Cs134	31.9	Bq/kg raw ± 10.9 Bq/kg raw		Cs134 13.0 Bq/kg raw
Vacuum cleaner dust (HITACHI Cyclone)	Joban, Iwaki	Oct-18	Cs137	211.0	Bq/kg raw ± 44.0 Bq/kg raw	235.5	Cs137 20.2 Bq/kg raw
			Cs134	24.5	Bq/kg raw ± 13.0 Bq/kg raw		Cs134 19.8 Bq/kg raw
Vacuum cleaner dust (HITACHI Cyclone)	Onahama-ohara, Iwaki	Oct-18	Cs137	40.8	Bq/kg raw ± 10.1 Bq/kg raw	40.8	Cs137 10.1 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134 7.7 Bq/kg raw
Air cleaner filter	Onahamakamikaziro, Iwaki	Nov-18	Cs137	1640.0	Bq/kg raw ± 330.0 Bq/kg raw	1819.0	Cs137 9.7 Bq/kg raw
			Cs134	179.0	Bq/kg raw ± 36.0 Bq/kg raw		Cs134 9.1 Bq/kg raw
Air dust	Nakosodaiichi Junior High School (schoolyard)	Oct-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0042 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Nakosodaini Junior High School (schoolyard)	Oct-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0041 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Kawabe Junior High School (schoolyard)	Oct-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0045 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Katono Junior High School (schoolyard)	Oct-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0049 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Iritono Junior High School (schoolyard)	Nov-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0042 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Yotukura Junior High School (schoolyard)	Nov-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0039 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Wakabadai Nursery School (playground)	Nov-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0045 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³
Air dust	Funao Nursery School (playground)	Nov-18	Cs137	—	Bq/m³ ± — Bq/m³	Under Minimum Limit of Detection	Cs137 0.0044 Bq/m³
			Cs134	—	Bq/m³ ± — Bq/m³		Cs134 — Bq/m³

※"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



# ★Beta-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month		Measurement Result	Uncertainty	Minimum Limit of Detection
Striped marlin (guts)	Off the coast of Iwaki	Oct-18	T(Organization)	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	1.31 Bq/Kg dry
Shell ginger (leaf)	Kume Island, Okinawa	Sep-18	T(Organization)	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	1.67 Bq/Kg dry
Potato	Rokkasho, Aomori	Sep-18	T(Free)	Under Minimum Limit of Detection Bq/L	± — Bq/L	1.98 Bq/L
Striped marlin (flesh)	Off the coast of Iwaki	Oct-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.16 Bq/Kg dry
Striped marlin (bone)	Off the coast of Iwaki	Oct-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.14 Bq/Kg dry
Striped marlin (guts)	Off the coast of Iwaki	Oct-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.16 Bq/Kg dry
Perilla	Mayumi-Hitachiota, Ibaraki	Jul-18	Sr90	1.44 Bq/Kg dry	± 0.19 Bq/Kg dry	0.28 Bq/Kg dry
Mulberry leaves	Mayumi-Hitachiota, Ibaraki	Jul-18	Sr90	3.62 Bq/Kg dry	± 0.55 Bq/Kg dry	0.67 Bq/Kg dry
Pine leaves	Tokai-Naka, Ibaraki	Jul-18	Sr90	1.33 Bq/Kg dry	± 0.44 Bq/Kg dry	0.63 Bq/Kg dry
Soil	Mayumi-Hitachiota, Ibaraki	Jul-18	Sr90	4.68 Bq/Kg dry	± 1.25 Bq/Kg dry	1.86 Bq/Kg dry
Soil	Mayumi-Hitachiota, Ibaraki	Jul-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	2.02 Bq/Kg dry
Soil	Tokai-Naka, Ibaraki	Jul-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	2.02 Bq/Kg dry
Red soil	Kume Island, Okinawa	Sep-18	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	1.64 Bq/Kg dry
Sea water A (surface)	Off the coast of Fukushima Nuclear Power Plant 2	Oct-18	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0017 Bq/L
Sea water	Kume Island, Okinawa	Sep-18	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0011 Bq/L
Spring water	Kume Island, Okinawa	Sep-18	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0015 Bq/L
Deep ocean water	Kume Island, Okinawa	Sep-18	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0020 Bq/L

※The value below Minimum Limit of Detection does not necessary mean 0(zero)Bq/Kg.