



Radiation Measurement Results of 125 Items in February



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection				
Rice	Watari, Fukushima, Fukushima	Oct-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.8	Bq/kg raw
Rice	Joban, Iwaki	Oct-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.8	Bq/kg raw
Rice	Akita	Oct-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.8	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.7	Bq/kg raw
Potato	Iwaki	Dec-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Potato	Nagasaki	Dec-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Yam	Hirono, Futaba, Fukushima	Feb-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.4	Bq/kg raw
Yam	Aomori	Feb-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.7	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.4	Bq/kg raw
Sweet potato (with peel)	Kanazawa, Ishikawa	Nov-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.7	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.5	Bq/kg raw
Red radish	Iwaki	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Japanese white radish(leaves)	Iwaki	Feb-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.9	Bq/kg raw
Cabbage	Iwaki	Feb-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.4	Bq/kg raw
Chinese cabbage	Iritono, Tono, Iwaki	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Chinese cabbage	Ibaraki	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Chinese cabbage	Hyogo	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Carrot	Shirakawa, Fukushima	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.4	Bq/kg raw
Carrot	Tomisato, Chiba	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Green onion	Ryouzen, Date, Fukushima	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.5	Bq/kg raw
Green onion	Iwaki	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Green onion	Fukushima	Jan-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.5	Bq/kg raw
Broccoli	Iwaki	Feb-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.7	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.4	Bq/kg raw

※"_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Japanese mustard spinach	Ryouzen, Date, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Potherb mustard	Ryouzen, Date, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.6 Bq/kg raw
Potherb mustard	Ibaraki	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	2.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.3 Bq/kg raw
Canola flower	Iwaki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.8 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.6 Bq/kg raw
Canola flower	Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.6 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.4 Bq/kg raw
Turnip (leaf)	Iwaki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Yacon	Hirono, Futaba, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Yacon	Furudono, Ishikawa, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2 Bq/kg raw
Aralia sprout	Kawauchi, Futaba, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	9.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	7.3 Bq/kg raw
Aralia cordate	Samegawa, Higashishirakawa, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.4 Bq/kg raw
Japanese parsley	Ibaraki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	3.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.5 Bq/kg raw
Tomato	Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Soybeans	Tsukidate, Date, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.7 Bq/kg raw
Green soybeans (pulp)	Taiwan	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Green soybeans (pod)	Taiwan	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.5 Bq/kg raw
Ginkgo(pulp)	Fukushima	Nov-18	Cs137	2.0 Bq/kg raw	±	0.8 Bq/kg raw	2.0	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Ginkgo(peel)	Fukushima	Nov-18	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	3.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.4 Bq/kg raw
Ginkgo(pulp)	Chiba	Nov-18	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Ginkgo(peel)	Chiba	Nov-18	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	3.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.5 Bq/kg raw
Dried Japanese white radish	Samegawa, Higashishirakawa, Fukushima	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	3.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.4 Bq/kg raw
Dried Japanese white radish	Ohisa, Iwaki	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	1.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.5 Bq/kg raw
Dried Japanese white radish	Iwaki	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	4.8 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.7 Bq/kg raw
Dried Japanese white radish	Izumitamatsuyu, Iwaki	Jan-19	Cs137	3.5 Bq/kg raw	±	2.7 Bq/kg raw	3.5	Cs137	3.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.6 Bq/kg raw
Dried stems of taro	Samegawa, Higashishirakawa, Fukushima	Dec-18	Cs137	— Bq/kg raw	±	— Bq/kg raw	検出下限値以下	Cs137	4.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.2 Bq/kg raw

※"_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Boar · male (liver)	Nagasakijigiri, Iwaki	Feb-19	Cs137	23.7 Bq/kg raw	± 4.7 Bq/kg raw	23.7	Cs137	1.5 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.3 Bq/kg raw	
Boar · male (heart, liver)	Nagasakijigiri, Iwaki	Jan-19	Cs137	20.2 Bq/kg raw	± 4.2 Bq/kg raw	20.2	Cs137	1.6 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw	
Boar · male (heart, liver)	Orito, Iwaki	Feb-19	Cs137	9.5 Bq/kg raw	± 2.1 Bq/kg raw	9.5	Cs137	1.3 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.2 Bq/kg raw	
Boar · male (heart, liver)	Ena, Iwaki	Feb-19	Cs137	19.9 Bq/kg raw	± 4.1 Bq/kg raw	19.9	Cs137	1.6 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw	
Flounder (head and bony parts)	Tairanumanouchi, Iwaki	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.1 Bq/kg raw	
Boiled salmon bone (canned)	Kesenuma, Miyagi	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.9 Bq/kg raw	
Green laver	Soma, Fukushima	Feb-19	Cs137	2.1 Bq/kg raw	± 1.0 Bq/kg raw	2.1	Cs137	1.5 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.2 Bq/kg raw	
Seaweed stem	Miyagi	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.0 Bq/kg raw	
Mekabu seaweed	Kesenuma, Miyagi	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.7 Bq/kg raw	
Shitake mushroom grown in bacteria-bed	Hokkaido	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.4 Bq/kg raw	
Nameko mushroom	Tamura, koriyama, Fukushima	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.2 Bq/kg raw	
Nameko mushroom	Yamatama, Iwaki	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.3 Bq/kg raw	
Iyokan(pulp)	Shizuoka	Jan-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.9 Bq/kg raw	
Iyokan(peel)	Shizuoka	Jan-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.8 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.4 Bq/kg raw	
Salted plum	China (production)	unknown	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.1 Bq/kg raw	
Dried persimmon	Ryouzen, Date, Fukushima	Dec-18	Cs137	3.2 Bq/kg raw	± 1.4 Bq/kg raw	3.2	Cs137	1.2 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.0 Bq/kg raw	
Egg	Hobara, Date, Fukushima	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.3 Bq/kg raw	
Egg(shell)	Hobara, Date, Fukushima	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	3.8 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.9 Bq/kg raw	
Egg	Date, Fukushima	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.9 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.7 Bq/kg raw	
Egg	Yazu, Yazu, Tottori	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.1 Bq/kg raw	
Soy pulp	Naka, Ibaraki	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.4 Bq/kg raw	
Soy pulp	Nikko, Tochigi	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.1 Bq/kg raw	
Tofu	Maebashi, Gunma	Jan-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.2 Bq/kg raw	
Konjac	Tabito, Iwaki	Feb-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.0 Bq/kg raw	

※"_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Konjac	Utsunomiya, Tochigi	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Konjac	Utsunomiya, Tochigi	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.4 Bq/kg raw
Sake lees	Ena, Iwaki	2018	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.8 Bq/kg raw
Boiled udon	Otagawa, Tochigi	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Boiled soba	Otagawa, Tochigi	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2 Bq/kg raw
Corn(canned)	America	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Yellow pickled radish	Hirono, Futaba, Fukushima	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Yogurt	Tubame, Niigata	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Yogurt	Takasaki, Gunma	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Milk beverage	Arai, Motomiya, Fukushima	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Milk beverage	Iwase, Odate, Akita	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2 Bq/kg raw
Soft drink	Hakushu, Hokuto, Yamanashi	Jan-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Soy sauce	Taira, Iwaki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.6 Bq/kg raw
Soy sauce	Choshi, Chiba	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.6 Bq/kg raw
Sugar	Hekinan, Aichi	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Source	Ryuou, Gamou, Siga	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Potato starch	Hokkaido	2018	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Okonomiyaki powder	Minamisenju, Arakawa, Tokyo	2018	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.2 Bq/kg raw
Rice flour	Japan (production)	2018	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.1 Bq/kg raw
School lunch	Uchigotakasaka, Iwaki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.7 Bq/kg raw
School lunch	Uchigotakasaka, Iwaki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.8 Bq/kg raw
School lunch	Jobanmatsugadai, Iwaki	Feb-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Water of a dam	Murohara, Namie, Futaba	Feb-19	Cs137	— Bq/L	±	— Bq/L	Under Minimum Limit of Detection	Cs137	0.9 Bq/L
			Cs134	— Bq/L	±	— Bq/L		Cs134	0.8 Bq/L
Meltwater	Taira, Iwaki	Jan-19	Cs137	— Bq/L	±	— Bq/L	Under Minimum Limit of Detection	Cs137	0.7 Bq/L
			Cs134	— Bq/L	±	— Bq/L		Cs134	0.7 Bq/L

※"_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Soil	Tomioka, Futaba	Feb-19	Cs137	16900.0 Bq/Kg dry	± 3400.0 Bq/Kg dry	18280.0	Cs137	11.5 Bq/Kg dry	
			Cs134	1380.0 Bq/Kg dry	± 280.0 Bq/Kg dry		Cs134	9.0 Bq/Kg dry	
Soil	Iritono, Tono, Iwaki	Feb-19	Cs137	2060.0 Bq/Kg dry	± 209.0 Bq/Kg dry	2220.0	Cs137	7.7 Bq/Kg dry	
			Cs134	160.0 Bq/Kg dry	± 18.3 Bq/Kg dry		Cs134	8.5 Bq/Kg dry	
Soil	Iritono, Tono, Iwaki	Feb-19	Cs137	124.0 Bq/Kg dry	± 14.2 Bq/Kg dry	133.3	Cs137	4.0 Bq/Kg dry	
			Cs134	9.3 Bq/Kg dry	± 2.0 Bq/Kg dry		Cs134	5.8 Bq/Kg dry	
Soil	Iritono, Tono, Iwaki	Feb-19	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	2.8 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	3.4 Bq/Kg dry	
Soil	Iritono, Tono, Iwaki	Feb-19	Cs137	250.0 Bq/Kg dry	± 28.1 Bq/Kg dry	272.1	Cs137	4.6 Bq/Kg dry	
			Cs134	22.1 Bq/Kg dry	± 4.1 Bq/Kg dry		Cs134	6.4 Bq/Kg dry	
Soil	Iritono, Tono, Iwaki	Feb-19	Cs137	32.1 Bq/Kg dry	± 4.3 Bq/Kg dry	32.1	Cs137	4.4 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	6.8 Bq/Kg dry	
Soil	Uchigotakasaka, Iwaki	Feb-19	Cs137	1450.0 Bq/Kg dry	± 159.0 Bq/Kg dry	1577.0	Cs137	9.2 Bq/Kg dry	
			Cs134	127.0 Bq/Kg dry	± 18.8 Bq/Kg dry		Cs134	11.7 Bq/Kg dry	
Soil	Uchigotakasaka, Iwaki	Feb-19	Cs137	18.3 Bq/Kg dry	± 2.8 Bq/Kg dry	18.3	Cs137	3.6 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	4.5 Bq/Kg dry	
Soil	Uchigotakasaka, Iwaki	Feb-19	Cs137	22.4 Bq/Kg dry	± 3.4 Bq/Kg dry	22.4	Cs137	4.6 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	5.4 Bq/Kg dry	
Soil	Watanabe, Iwaki	Jan-19	Cs137	151.0 Bq/Kg dry	± 18.0 Bq/Kg dry	160.6	Cs137	4.9 Bq/Kg dry	
			Cs134	9.6 Bq/Kg dry	± 3.0 Bq/Kg dry		Cs134	7.5 Bq/Kg dry	
Soil	Hanamigawa, Chiba, Chiba	Jan-19	Cs137	211.0 Bq/Kg dry	± 18.0 Bq/Kg dry	230.0	Cs137	4.5 Bq/Kg dry	
			Cs134	19.0 Bq/Kg dry	± 3.4 Bq/Kg dry		Cs134	5.5 Bq/Kg dry	
Soil	Hanamigawa, Chiba, Chiba	Jan-19	Cs137	156.0 Bq/Kg dry	± 18.9 Bq/Kg dry	168.3	Cs137	4.8 Bq/Kg dry	
			Cs134	12.3 Bq/Kg dry	± 3.5 Bq/Kg dry		Cs134	5.9 Bq/Kg dry	
Soil	Hanamigawa, Chiba, Chiba	Jan-19	Cs137	95.1 Bq/Kg dry	± 11.2 Bq/Kg dry	103.4	Cs137	4.6 Bq/Kg dry	
			Cs134	8.3 Bq/Kg dry	± 1.8 Bq/Kg dry		Cs134	5.9 Bq/Kg dry	
Soil	Hanamigawa, Chiba, Chiba	Jan-19	Cs137	95.2 Bq/Kg dry	± 12.6 Bq/Kg dry	95.2	Cs137	4.0 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	4.7 Bq/Kg dry	
Soil	Hanamigawa, Chiba, Chiba	Jan-19	Cs137	38.3 Bq/Kg dry	± 5.4 Bq/Kg dry	38.3	Cs137	4.5 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	7.2 Bq/Kg dry	
Soil	Wakakodama, Gyoda, Saitama	Feb-19	Cs137	77.0 Bq/Kg dry	± 8.9 Bq/Kg dry	85.5	Cs137	4.0 Bq/Kg dry	
			Cs134	8.5 Bq/Kg dry	± 1.8 Bq/Kg dry		Cs134	6.2 Bq/Kg dry	
Soil	Nakatsugawa, Gifu	Feb-19	Cs137	— Bq/Kg dry	± — Bq/Kg dry	Under Minimum Limit of Detection	Cs137	3.5 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	3.5 Bq/Kg dry	
Gravel	Higashimizumoto, Katushika, Tokyo	Feb-19	Cs137	11.8 Bq/Kg dry	± 2.5 Bq/Kg dry	11.8	Cs137	2.8 Bq/Kg dry	
			Cs134	— Bq/Kg dry	± — Bq/Kg dry		Cs134	3.1 Bq/Kg dry	
Moss	Nishiki, Iwaki	Feb-19	Cs137	1770.0 Bq/Kg raw	± 350.0 Bq/Kg raw	1955.0	Cs137	10.7 Bq/Kg raw	
			Cs134	185.0 Bq/Kg raw	± 37.0 Bq/Kg raw		Cs134	9.8 Bq/Kg raw	
Vacuum cleaner dust (paper pack)	Joban, Iwaki	Feb-19	Cs137	303.0 Bq/Kg raw	± 40.6 Bq/Kg raw	303.0	Cs137	24.7 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	18.3 Bq/Kg raw	
Vacuum cleaner dust	Uchigoumidaisakai, Iwaki	Feb-19	Cs137	91.9 Bq/Kg raw	± 20.6 Bq/Kg raw	91.9	Cs137	12.5 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	9.9 Bq/Kg raw	
Vacuum cleaner dust(Cyclone)	Onahama, Iwaki	Feb-19	Cs137	68.8 Bq/Kg raw	± 17.2 Bq/Kg raw	68.8	Cs137	16.1 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	12.7 Bq/Kg raw	
Vacuum cleaner dust	Kume Island, Okinawa	Feb-19	Cs137	— Bq/Kg raw	± — Bq/Kg raw	Under Minimum Limit of Detection	Cs137	3.9 Bq/Kg raw	
			Cs134	— Bq/Kg raw	± — Bq/Kg raw		Cs134	3.0 Bq/Kg raw	

※"_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



★Beta-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty			Minimum Limit of Detection	
Sea water B (surface)	Off the coast of Fukushima Nuclear Power Plant 2	Oct-18	T(Free)	Under Minimum Limit of Detection	Bq/L	±	—	Bq/L	1.98	Bq/L
Sea water B (lower)	Off the coast of Fukushima Nuclear Power Plant 2	Oct-18	T(Free)	Under Minimum Limit of Detection	Bq/L	±	—	Bq/L	1.98	Bq/L
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant 2	Oct-18	T(Free)	Under Minimum Limit of Detection	Bq/L	±	—	Bq/L	1.98	Bq/L
Sea water C (lower)	Off the coast of Fukushima Nuclear Power Plant 2	Oct-18	T(Free)	Under Minimum Limit of Detection	Bq/L	±	—	Bq/L	1.98	Bq/L
Oyster	Matushima, Miyagi	Nov-18	Sr90	Under Minimum Limit of Detection	Bq/Kg dry	±	—	Bq/Kg dry	0.15	Bq/Kg dry
Soil	along River Esk, England	Oct-14	Sr90	6.98	Bq/Kg dry	±	0.90	Bq/Kg dry	1.28	Bq/Kg dry
Soil	golf course of Seascale village, England	Oct-14	Sr90	Under Minimum Limit of Detection	Bq/Kg dry	±	—	Bq/Kg dry	1.82	Bq/Kg dry
Soil	Russel park, England	Oct-14	Sr90	Under Minimum Limit of Detection	Bq/Kg dry	±	—	Bq/Kg dry	1.80	Bq/Kg dry
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant 1	Jul-18	Sr90	0.0012	Bq/L	±	0.0007	Bq/L	0.0011	Bq/L