



# Radiation Measurement Results of 157 Items in July



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection				
			Cs137	Cs134	±	±		Cs137	Cs134			
Rice	Sasaya, Fukushima, Fukushima	Oct-18	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.8	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.8	Bq/kg raw
Potato	Naraha, Futaba, Fukushima	Jul-19	Cs137	5.1	Bq/kg raw	±	1.8	Bq/kg raw	5.1	Cs137	1.5	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Potato	Hirono, Futaba, Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Potato	Izumi, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Potato (peel)	Izumi, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.8	Bq/kg raw
Potato	Iritono, Tono, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Potato	Nakoso, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Potato	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Sweet potato	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.6	Bq/kg raw
Sweet potato	Chiba	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Carrot	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.5	Bq/kg raw
Carrot	Chiba	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Onion	Kunimi, Date, Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Onion	Japan (production)	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
White eggplant	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Eggplant	Gunma	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Cucumber①	Naraha, Futaba, Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Cucumber②	Naraha, Futaba, Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Cucumber	Onahama-tamagawa, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Cucumber	Onahama, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.8	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.5	Bq/kg raw

※"\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection		
Cucumber	Nishiki, Iwaki	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1	Bq/kg raw
Cucumber	Nakoso, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2	Bq/kg raw
Cabbage	Iritono, Tono, Iwaki	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0	Bq/kg raw
Cabbage	Chiba	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1	Bq/kg raw
Qing-geng-cai	Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.8	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.7	Bq/kg raw
Qing-geng-cai	Ibaraki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.7	Bq/kg raw
Green bean	Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0	Bq/kg raw
Green bean (frozen)	Thai land	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0	Bq/kg raw
String beans	Iritono, Tono, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2	Bq/kg raw
Broccoli	Fukushima	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2	Bq/kg raw
Cauliflower (frozen)	Belgium	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2	Bq/kg raw
Bitter gourd	Miyazaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0	Bq/kg raw
Zucchini①	Naraha, Futaba, Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.7	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.5	Bq/kg raw
Zucchini②	Naraha, Futaba, Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0	Bq/kg raw
Zucchini	Tairashimotakaku, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1	Bq/kg raw
Red perilla	Iwaki	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.8	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.1	Bq/kg raw
Ginger	Ibaraki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9	Bq/kg raw
Scallion	Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3	Bq/kg raw
Scallion	Ibaraki	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.4	Bq/kg raw
Bean sprout	Arai, Motomiya, Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1	Bq/kg raw
Pea sprout (leaves)	Yamanashi	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.9	Bq/kg raw
Pea sprout (roots)	Yamanashi	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2	Bq/kg raw
Tomato	Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2	Bq/kg raw
Pumpkin	Okuma, Futaba, Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0	Bq/kg raw

※"\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection				
Pumpkin	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.8	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.7	Bq/kg raw
Japanese white radish(leaves)	Nishiki, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.8	Bq/kg raw
Corn	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Corn(peel, core)	Ibaraki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.8	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.8	Bq/kg raw
Butterbur	Iritono, Tono, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.5	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	2.3	Bq/kg raw
Shitake mushroom grown in bacteria-bed	Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Wood ear mushroom(raw)	Minamiyamashiro, Kyoto	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	10.5	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	8.0	Bq/kg raw
Nameko mushroom	Koriyama, Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Nameko mushroom	Mogami, Mogami, Yamagata	2018	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Fig	Izumigaoka, Iwaki	Jul-19	Cs137	5.2	Bq/kg raw	±	1.3	Bq/kg raw	5.2	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Loquat	Izumigaoka, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	6.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	6.2	Bq/kg raw
Blueberry	Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Blackberry	Taira, Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.2	Bq/kg raw
Plum	Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Water melon (peel)	Ibaraki	Jun-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.8	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.5	Bq/kg raw
Mandarin orange(frozen)	China (production)	unknown	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Cherry(frozen)	Hungary	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Salted mackerel(flesh)	Fukushima	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Greater amberjack (flesh)	Off the coast of Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.1	Bq/kg raw
Yellowtail (flesh)	Off the coast of Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Pacific cod (flesh)	Off the coast of Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	0.9	Bq/kg raw
Shark(flesh)	Off the coast of Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Skipjack tuna (bony parts)	Off the coast of Iwaki	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw
Japanese sardine (whole)	Chiba	Jul-19	Cs137	—	Bq/kg raw	±	—	Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2	Bq/kg raw
			Cs134	—	Bq/kg raw	±	—	Bq/kg raw		Cs134	1.0	Bq/kg raw

※"\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Goby(whole)	Same river estuary	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	9.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	7.7 Bq/kg raw
Dried whitebait	Ibaraki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Dried small sardines	Japan (production)	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Krill	Miyagi	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.5 Bq/kg raw
Silver mouthed Monodont	Onahama, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Oyster(frozen)	Korea (production)	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Boiled seaweed	Off the coast of Sanriku	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.4 Bq/kg raw
Saltwort	Chiba	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.7 Bq/kg raw
Sea lettuce	Onahama, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	4.6 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.5 Bq/kg raw
Dried green seaweed	Soma, Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.7 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.6 Bq/kg raw
Agar weed	Onahama, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.1 Bq/kg raw
Sea hare	Onahama, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Roasted laver	Fukuoka	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	3.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.8 Bq/kg raw
Boiled yellowfin tuna (canned)	Japan (production)	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Grasshoppers cooked in soy sauce	Furudono, Ishikawa, Fukushima	Oct-18	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.8 Bq/kg raw
Pork (minced)	Japan (production)	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Chicken breast	Japan (production)	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Wiener sausage	Kakegawa, Shizuoka	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2 Bq/kg raw
Salted plum	Yoshima, Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Rice miso	Taira, Iwaki	2019	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.2 Bq/kg raw
Natto	Mito, Ibaraki	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.6 Bq/kg raw
Natto(straw)	Mito, Ibaraki	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Tofu	Maebashi, Gunma	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Bread	Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw

※"\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Bread	Chiyoda-ku, Tokyo	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.6 Bq/kg raw
Rice flour (rice50%·glutinous rice50%)	Iwaki	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Frozen rice cake	Katsurao, Futaba, Fukushima	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Pocessed flour	Japan (production)	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Soy pulp	Japan (production)	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.4 Bq/kg raw
Sweet bean paste	Hokkaido	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
French fries (frozen)	Belgium	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Chicken hamburg	Ishidoriya, Hanamaki, Iwate	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Prepared liquid milk	Isesaki, Gunma	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.8 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.7 Bq/kg raw
Yogurt(drink)	Chuou-ku, Tokyo	Jul-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.0 Bq/kg raw
Tomato juice	Nagano, Nagano	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Amazake (fermented rice drink)	Tamura, Koriyama, Fukushima	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Soy milk	Japan (production)	Jun-19	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.0 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.9 Bq/kg raw
Sports drink (powder)	Japan (production)	May-18	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.1 Bq/kg raw
Lemon juice	India	unknown	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.9 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.8 Bq/kg raw
River water	Naraha, Futaba, Fukushima	2019	Cs137	— Bq/L	±	— Bq/L	Under Minimum Limit of Detection	Cs137	0.7 Bq/L
			Cs134	— Bq/L	±	— Bq/L		Cs134	0.6 Bq/L
Japanese cleyera	Iwaki	Jul-19	Cs137	26.7 Bq/kg raw	±	6.3 Bq/kg raw	26.7	Cs137	5.8 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	4.2 Bq/kg raw
Rotten wood(for growing shitake mushroom)	Japan (production)	unknown	Cs137	2.8 Bq/kg raw	±	1.1 Bq/kg raw	2.8	Cs137	1.5 Bq/kg raw
			Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	1.3 Bq/kg raw
Pine cone	Tomioaka, Futaba, Fukushima	May-19	Cs137	161.0 Bq/kg raw	±	32.0 Bq/kg raw	182.2	Cs137	8.7 Bq/kg raw
			Cs134	21.2 Bq/kg raw	±	6.6 Bq/kg raw		Cs134	7.8 Bq/kg raw
Pine cone	Naraha, Futaba, Fukushima	Apr-19	Cs137	308.0 Bq/kg raw	±	62.0 Bq/kg raw	340.7	Cs137	5.1 Bq/kg raw
			Cs134	32.7 Bq/kg raw	±	7.3 Bq/kg raw		Cs134	4.6 Bq/kg raw
Pine cone	Hirono, Futaba, Fukushima	May-19	Cs137	737.0 Bq/kg raw	±	147.0 Bq/kg raw	808.3	Cs137	9.6 Bq/kg raw
			Cs134	71.3 Bq/kg raw	±	15.9 Bq/kg raw		Cs134	8.6 Bq/kg raw
Pine cone①	Iidate, Soma, Fukushima	May-19	Cs137	5950.0 Bq/kg raw	±	1190.0 Bq/kg raw	6606.0	Cs137	14.1 Bq/kg raw
			Cs134	656.0 Bq/kg raw	±	131.0 Bq/kg raw		Cs134	13.8 Bq/kg raw
Pine cone②	Iidate, Soma, Fukushima	May-19	Cs137	1050.0 Bq/kg raw	±	210.0 Bq/kg raw	1164.0	Cs137	13.9 Bq/kg raw
			Cs134	114.0 Bq/kg raw	±	26.0 Bq/kg raw		Cs134	13.7 Bq/kg raw
Pine cone	Moriai, Fukushima, Fukushima	Apr-19	Cs137	2160.0 Bq/kg raw	±	430.0 Bq/kg raw	2365.0	Cs137	11.6 Bq/kg raw
			Cs134	205.0 Bq/kg raw	±	41.0 Bq/kg raw		Cs134	10.4 Bq/kg raw

※"\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Pine cone	Matsukawa, Fukushima, Fukushima	May-19	Cs137	547.0 Bq/kg raw	± 109.0 Bq/kg raw	597.9	Cs137	5.7 Bq/kg raw	
			Cs134	50.9 Bq/kg raw	± 11.0 Bq/kg raw		Cs134	5.2 Bq/kg raw	
Pine cone	Sukagawa, Fukushima	May-19	Cs137	433.0 Bq/kg raw	± 87.0 Bq/kg raw	473.6	Cs137	11.6 Bq/kg raw	
			Cs134	40.6 Bq/kg raw	± 11.1 Bq/kg raw		Cs134	11.4 Bq/kg raw	
Pine cone	Shimookuri, Kawamae, Iwaki	Apr-19	Cs137	32.3 Bq/kg raw	± 7.9 Bq/kg raw	32.3	Cs137	6.9 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	6.8 Bq/kg raw	
Pine cone	Ogawa, Iwaki	Apr-19	Cs137	352.0 Bq/kg raw	± 70.0 Bq/kg raw	387.8	Cs137	5.5 Bq/kg raw	
			Cs134	35.8 Bq/kg raw	± 8.1 Bq/kg raw		Cs134	4.9 Bq/kg raw	
Pine cone	Tairasimotakaku, Iwaki	Apr-19	Cs137	200.0 Bq/kg raw	± 40.0 Bq/kg raw	222.3	Cs137	6.5 Bq/kg raw	
			Cs134	22.3 Bq/kg raw	± 6.1 Bq/kg raw		Cs134	6.4 Bq/kg raw	
Pine cone	tairayotsunami, Iwaki	Apr-19	Cs137	119.0 Bq/kg raw	± 24.0 Bq/kg raw	130.5	Cs137	4.4 Bq/kg raw	
			Cs134	11.5 Bq/kg raw	± 3.5 Bq/kg raw		Cs134	4.3 Bq/kg raw	
Pine cone	Chuoudaikashima, Iwaki	Apr-19	Cs137	965.0 Bq/kg raw	± 193.0 Bq/kg raw	1064.8	Cs137	6.5 Bq/kg raw	
			Cs134	99.8 Bq/kg raw	± 20.2 Bq/kg raw		Cs134	6.3 Bq/kg raw	
Pine cone	Onahama- shimokaziro, Iwaki	Apr-19	Cs137	275.0 Bq/kg raw	± 55.0 Bq/kg raw	301.0	Cs137	6.5 Bq/kg raw	
			Cs134	26.0 Bq/kg raw	± 6.7 Bq/kg raw		Cs134	6.3 Bq/kg raw	
Pine cone	Shimogawa, Izumi, Iwaki	Apr-19	Cs137	444.0 Bq/kg raw	± 89.0 Bq/kg raw	481.3	Cs137	12.4 Bq/kg raw	
			Cs134	37.3 Bq/kg raw	± 10.8 Bq/kg raw		Cs134	11.3 Bq/kg raw	
Moss	Jyobanmizunoya, Iwaki	Jul-19	Cs137	556.0 Bq/kg raw	± 111.0 Bq/kg raw	618.4	Cs137	1.5 Bq/kg raw	
			Cs134	62.4 Bq/kg raw	± 12.5 Bq/kg raw		Cs134	1.3 Bq/kg raw	
Moss	Nishiki, Iwaki	Jul-19	Cs137	883.0 Bq/kg raw	± 177.0 Bq/kg raw	958.8	Cs137	3.5 Bq/kg raw	
			Cs134	75.8 Bq/kg raw	± 15.2 Bq/kg raw		Cs134	2.7 Bq/kg raw	
Soil (under the drain)	Onahama- sumiyoshi, Iwaki	Jul-19	Cs137	3940.0 Bq/kg dry	± 427.0 Bq/kg dry	4274.0	Cs137	9.4 Bq/kg dry	
			Cs134	334.0 Bq/kg dry	± 44.3 Bq/kg dry		Cs134	9.0 Bq/kg dry	
Organic culture soil	Japan (production)	unknown	Cs137	— Bq/kg dry	± — Bq/kg dry	Under Minimum Limit of Detection	Cs137	1.8 Bq/kg dry	
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.7 Bq/kg dry	
Curtain	Izumigaoka, Iwaki	Jul-19	Cs137	501.0 Bq/kg raw	± 100.0 Bq/kg raw	547.5	Cs137	3.2 Bq/kg raw	
			Cs134	46.5 Bq/kg raw	± 9.3 Bq/kg raw		Cs134	2.9 Bq/kg raw	
Cleaning sheet (ventilation fan dust)	Izumigaoka, Iwaki	Jul-19	Cs137	35500.0 Bq/kg raw	± 7100.0 Bq/kg raw	39120.0	Cs137	82.9 Bq/kg raw	
			Cs134	3620.0 Bq/kg raw	± 720.0 Bq/kg raw		Cs134	75.8 Bq/kg raw	
cleaning sheet (dust outside the window frame on the second floor)	Izumigaoka, Iwaki	Jul-19	Cs137	4290.0 Bq/kg raw	± 860.0 Bq/kg raw	4628.0	Cs137	4.9 Bq/kg raw	
			Cs134	338.0 Bq/kg raw	± 68.0 Bq/kg raw		Cs134	3.8 Bq/kg raw	
Cleaning sheet (dust inside the window frame)	Onahamasuwa, Iwaki	Jun-19	Cs137	462.3 Bq/kg raw	± 43.2 Bq/kg raw	493.3	Cs137	8.4 Bq/kg raw	
			Cs134	31.0 Bq/kg raw	± 7.5 Bq/kg raw		Cs134	7.8 Bq/kg raw	
Cleaning sheet (dust on the floor)	Onahamasuwa, Iwaki	Jul-19	Cs137	81.8 Bq/kg raw	± 11.7 Bq/kg raw	81.8	Cs137	8.2 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	7.1 Bq/kg raw	
Dust cloth (dust on the floor)	Ueda, Iwaki	Jul-19	Cs137	898.0 Bq/kg raw	± 180.0 Bq/kg raw	990.0	Cs137	13.3 Bq/kg raw	
			Cs134	92.0 Bq/kg raw	± 20.6 Bq/kg raw		Cs134	12.3 Bq/kg raw	
Empty can	Naraha, Futaba, Fukushima	Jul-19	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	33.7 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	32.7 Bq/kg raw	
Vacuum cleaner dust (TOSHIBA, paper pack)	Miharu, Tamura, Fukushima	Feb-19	Cs137	182.0 Bq/kg raw	± 36.0 Bq/kg raw	211.5	Cs137	10.5 Bq/kg raw	
			Cs134	29.5 Bq/kg raw	± 8.5 Bq/kg raw		Cs134	9.7 Bq/kg raw	
Vacuum cleaner dust	Keitoku, Kitakata, Fukushima	Jul-19	Cs137	46.4 Bq/kg raw	± 6.0 Bq/kg raw	46.4	Cs137	3.4 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.6 Bq/kg raw	
Vacuum cleaner dust (Dyson)	Onahama- hanabatake, Iwaki	Jun-19	Cs137	591.6 Bq/kg raw	± 54.3 Bq/kg raw	623.9	Cs137	5.4 Bq/kg raw	
			Cs134	32.3 Bq/kg raw	± 7.2 Bq/kg raw		Cs134	4.8 Bq/kg raw	
Vacuum cleaner dust (Dyson)	Onahama-ohara, Iwaki	Jun-19	Cs137	110.1 Bq/kg raw	± 17.1 Bq/kg raw	110.1	Cs137	12.2 Bq/kg raw	
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	9.2 Bq/kg raw	

※"\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result	Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Air dust	Onahamadaiichi Junior High School (schoolyard)	Jun-19	Cs137	— Bq/m <sup>3</sup> ± — Bq/m <sup>3</sup>	検出下限値以下	Cs137	0.0046 Bq/m <sup>3</sup>
			Cs134	— Bq/m <sup>3</sup> ± — Bq/m <sup>3</sup>		Cs134	— Bq/m <sup>3</sup>

※"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



## ★Beta-ray

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Minimum Limit of Detection
Surgassum (seaweed)	Genkainada	2019	T(Organization)	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.93 Bq/Kg dry
Slime flounder (flesh)	Off the coast of Fukushima Nuclear Power Plant1	Apr-18	T(Organization)	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	1.39 Bq/Kg dry
Mountain water①	Shimookeuri, Kawamae, Iwaki	May-19	T(Free)	Under Minimum Limit of Detection Bq/L	± — Bq/L	2.02 Bq/L
Mountain water②	Shimookeuri, Kawamae, Iwaki	May-19	T(Free)	Under Minimum Limit of Detection Bq/L	± — Bq/L	2.02 Bq/L
Citron	okawara, Okuma, Futaba, Fukushima	Nov-15	Sr90	4.38 Bq/Kg dry	± 0.54 Bq/Kg dry	0.77 Bq/Kg dry
Citron	Naraha, Futaba, Fukushima	Nov-15	Sr90	1.00 Bq/Kg dry	± 0.09 Bq/Kg dry	0.13 Bq/Kg dry
Persimmon	okawara, Okuma, Futaba, Fukushima	Nov-15	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.16 Bq/Kg dry
Persimmon	Minamisoma, Fukushima	Nov-15	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.17 Bq/Kg dry
Greenling (flesh, bone)	Off the coast of Iwaki	Jun-19	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.13 Bq/Kg dry
Yellowtail (bone)	Off the coast of Iwaki	Jun-19	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.12 Bq/Kg dry
Greater amberjack	Off the coast of Iwaki	Jul-19	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.16 Bq/Kg dry
Spirulina	Hawaii	2015	Sr90	Under Minimum Limit of Detection Bq/Kg dry	± — Bq/Kg dry	0.13 Bq/Kg dry
Tap water①	Kawauchi, Futaba, Fukushima	Dec-15	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0017 Bq/L
Tap water②	Kawauchi, Futaba, Fukushima	Dec-15	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0016 Bq/L
Sea water C (surface)	Off the coast of Fukushima Nuclear Power Plant1	Apr-19	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0014 Bq/L
Sea water C (lower)	Off the coast of Fukushima Nuclear Power Plant1	Apr-19	Sr90	Under Minimum Limit of Detection Bq/L	± — Bq/L	0.0014 Bq/L

※The value below Minimum Limit of Detection does not necessary mean 0(zero)Bq/Kg.