



# Radiation Measurement Results of 176 Items in May



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

## ★Gamma-ray

Measuring instrument		Feature	Guide to lower limit※
Na I Scintillation Spectrometer			
Product of ATOMTEX AT1320A	Product of BERTHOLD LB2045	· Gamma-ray spectrometer with Na I scintillation detector.	Food (Sample 1kg) Lower limit 1.0Bq/Kg Soil (Sample 1kg) Lower limit 2.5Bq/Kg Material (Sample 1kg) Lower limit 1.0Bq/Kg Water (Sample 20L) Lower limit 0.02Bq/L
Germanium Semiconductor detector			
ORTEC GEM30-70		· Radioactivity measurement series. Quantitative analysis based on "Gamma-ray spectrometry with germanium semiconductor detector." · Relative efficiency 35%	Food (Sample 2kg) Lower limit 0.04Bq/Kg Soil (Sample 1kg) Lower limit 0.06Bq/Kg Material (Sample 1kg) Lower limit 0.06Bq/Kg Water (Sample 20L) Lower limit 0.001Bq/L

※The lower limit varies depending on the sample weight and measurement time.

Measuring instrument:Na I Scintillation Spectrometer (Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result	Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Glutinous rice	Fukushima Pref.	Oct-20	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.6 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.5 Bq/kg raw
Onion	Watanabe, Iwaki	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.2 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 2.1 Bq/kg raw
Cabbage	Nihonmatsu, Fukushima	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.0 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.7 Bq/kg raw
Cabbage	Motomiya, Fukushima	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.9 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 2.7 Bq/kg raw
Chinese cabbage	Miharu, Tamura, Fukushima	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 3.0 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 2.8 Bq/kg raw
Japanese white radish(pulp)	Iwaki city	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 2.0 Bq/kg raw
Japanese white radish(leaf)	Iwaki city	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.7 Bq/kg raw
Turnip(pulp)	Iwaki city	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.3 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 2.1 Bq/kg raw
Turnip (leaf)	Iwaki city	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.1 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.7 Bq/kg raw
Radish	Iwaki city	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.5 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.2 Bq/kg raw
Welsh onion	Otama, Adachi, Fukushima	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 2.3 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 2.0 Bq/kg raw
Green onion	Fukushima Pref.	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.8 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.5 Bq/kg raw
Cucumber	Fukushima, Fukushima Pref.	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.8 Bq/kg raw
Cucumber	Iwaki city	May-21	Cs137	— Bq/kg raw	Under Minimum Limit of Detection	Cs137 1.9 Bq/kg raw
			Cs134	— Bq/kg raw		Cs134 1.8 Bq/kg raw

※"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessarily mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Zucchini	Fukushima, Fukushima Pref.	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.9 Bq/kg raw
Zucchini	Iwaki city	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.4 Bq/kg raw
Snow pea	Hobara, Date, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.2 Bq/kg raw
Spinach	Iwaki city	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.0 Bq/kg raw
Santouna	Nihonmatsu, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.8 Bq/kg raw
Kukitachina	Otama, Adachi, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.7 Bq/kg raw
Norabouna	Iwaki city	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.7 Bq/kg raw
Kale	Iwaki city	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.3 Bq/kg raw
Watercress	Miyakoji, Tamura, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	3.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	3.6 Bq/kg raw
Mizuna	Nihonmatsu, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.6 Bq/kg raw
Leek	Fukushima Pref.	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Garlic sprout	Date, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.2 Bq/kg raw
Bamboo shoot (Small)	Matsukawa, Fukushima Fukushima Pref.	May-21	Cs137	19.8 Bq/kg raw	± 4.4 Bq/kg raw	19.8	Cs137	2.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.5 Bq/kg raw
Bamboo shoot (Large)	Matsukawa, Fukushima Fukushima Pref.	May-21	Cs137	13.1 Bq/kg raw	± 3.1 Bq/kg raw	13.1	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.1 Bq/kg raw
Bamboo shoot	Matsukawa, Fukushima Fukushima Pref.	May-21	Cs137	10.4 Bq/kg raw	± 2.4 Bq/kg raw	10.4	Cs137	1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.7 Bq/kg raw
Bamboo shoot	Ryouzen, Date, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.9 Bq/kg raw
Bamboo shoot①	Miharu, Tamura, Fukushima	May-21	Cs137	8.7 Bq/kg raw	± 2.2 Bq/kg raw	8.7	Cs137	1.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Bamboo shoot②	Miharu, Tamura, Fukushima	May-21	Cs137	6.8 Bq/kg raw	± 2.0 Bq/kg raw	6.8	Cs137	2.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.2 Bq/kg raw
Bamboo shoot	Nishida, Koriyama Fukushima	May-21	Cs137	2.5 Bq/kg raw	± 1.4 Bq/kg raw	2.5	Cs137	2.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.1 Bq/kg raw
Bamboo shoot	Kamiogawa, Ogawa, Iwaki	May-21	Cs137	7.4 Bq/kg raw	± 1.4 Bq/kg raw	7.4	Cs137	1.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.0 Bq/kg raw
Bracken(Wild)	Fukushima, Fukushima Pref.	May-21	Cs137	22.4 Bq/kg raw	± 3.2 Bq/kg raw	22.4	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Bracken(Wild)	Motomiya, Fukushima	May-21	Cs137	9.7 Bq/kg raw	± 2.2 Bq/kg raw	9.7	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.6 Bq/kg raw
Bracken(Wild)	Otama, Adachi, Fukushima	May-21	Cs137	8.4 Bq/kg raw	± 2.0 Bq/kg raw	8.4	Cs137	1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Bracken(Wild)	Miyakoji, Tamura, Fukushima	May-21	Cs137	3.2 Bq/kg raw	± 1.7 Bq/kg raw	3.2	Cs137	2.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.3 Bq/kg raw

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Bracken	Minamiaizu, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	3.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.8 Bq/kg raw
Bracken(Wild)	Esashi, Osyu, Iwate	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	4.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	3.4 Bq/kg raw
Dried bracken	Yonezawa, Yamagata	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	4.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	3.7 Bq/kg raw
Ostrich fern	Watari, Watari-gun, Miyagi	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.1 Bq/kg raw
Butterbur	Soma, Fukushima	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	4.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	3.3 Bq/kg raw
Butterbur	Iwaki city	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.9 Bq/kg raw
Butterbur	Kashima, Iwaki	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.8 Bq/kg raw
Udo(Cultivation)	Soma, Fukushima	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.4 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.9 Bq/kg raw
Mountain udo	Kitashiobara, Yama, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Mountain udo	Minamiaizu, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	4.2 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	3.4 Bq/kg raw
Mountain udo(Leaf)	Iwaki city	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.2 Bq/kg raw
Japanese mugwort	Minamiaizuchi, Minamiaizu, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.7 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.2 Bq/kg raw
Hosta	Yamagata Pref.	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	3.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.8 Bq/kg raw
Walnut	Nishigo, Nishishirakawa, Fukushima	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	15.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	12.3 Bq/kg raw
Pressed barley	Japan (production)	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.8 Bq/kg raw
Milk	Fukushima Pref.	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Milk	Chiba Pref.	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.6 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.5 Bq/kg raw
Yogurt	Miyagi Pref.	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.0 Bq/kg raw
Soybeans	Noda, Chiba	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.3 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.9 Bq/kg raw
Soy pulp	Nihonmatsu, Fukushima	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.8 Bq/kg raw
Chicken liver	Japan (production)	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.1 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.8 Bq/kg raw
Dried udon	Tamura, Fukushima	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.9 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.7 Bq/kg raw
Glass noodles	China (production)	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.5 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	2.0 Bq/kg raw
Tuna flakes	Japan (production)	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.8 Bq/kg raw

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Pancake mix	Japan (production)	Apr-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	2.0 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	1.9 Bq/kg raw
Trifolium pratense	Onahama, Iwaki	May-21	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	5.8 Bq/kg raw
			Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	5.5 Bq/kg raw
Soil	Onami, Fukushima, Fukushima Pref.	May-21	Cs137	7110.0 Bq/kg dry	± 723.0 Bq/kg dry	7421.0	Cs137	10.9 Bq/kg dry
			Cs134	311.0 Bq/kg dry	± 33.9 Bq/kg dry		Cs134	9.0 Bq/kg dry
Soil	Onami, Fukushima, Fukushima Pref.	May-21	Cs137	2370.0 Bq/kg dry	± 244.0 Bq/kg dry	2474.0	Cs137	6.2 Bq/kg dry
			Cs134	104.0 Bq/kg dry	± 12.1 Bq/kg dry		Cs134	5.7 Bq/kg dry
Soil①	Okawara, Okuma, Futaba, Fukushima	May-21	Cs137	78.5 Bq/kg dry	± 9.1 Bq/kg dry	83.1	Cs137	1.8 Bq/kg dry
			Cs134	4.6 Bq/kg dry	± 1.1 Bq/kg dry		Cs134	2.5 Bq/kg dry
Soil②	Okawara, Okuma, Futaba, Fukushima	May-21	Cs137	68.8 Bq/kg dry	± 7.7 Bq/kg dry	68.8	Cs137	2.0 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	2.4 Bq/kg dry
Soil①	Nagasaki, Iwaki	Apr-21	Cs137	4020.0 Bq/kg dry	± 408.0 Bq/kg dry	4198.0	Cs137	5.0 Bq/kg dry
			Cs134	178.0 Bq/kg dry	± 19.1 Bq/kg dry		Cs134	4.9 Bq/kg dry
Soil②	Nagasaki, Iwaki	Apr-21	Cs137	109.0 Bq/kg dry	± 12.0 Bq/kg dry	113.0	Cs137	1.5 Bq/kg dry
			Cs134	4.0 Bq/kg dry	± 0.9 Bq/kg dry		Cs134	2.0 Bq/kg dry
Soil①	Onahama-tamagawa, Iwaki	May-21	Cs137	2540.0 Bq/kg dry	± 261.0 Bq/kg dry	2651.0	Cs137	6.3 Bq/kg dry
			Cs134	111.0 Bq/kg dry	± 12.7 Bq/kg dry		Cs134	5.9 Bq/kg dry
Soil②	Onahama-tamagawa, Iwaki	May-21	Cs137	7.5 Bq/kg dry	± 1.1 Bq/kg dry	7.5	Cs137	1.7 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.9 Bq/kg dry
Soil	Izumi-tamatsuyu, Iwaki	May-21	Cs137	— Bq/kg dry	± — Bq/kg dry	Under Minimum Limit of Detection	Cs137	1.1 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.3 Bq/kg dry
Soil (in the park)	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	3820.0 Bq/kg dry	± 392.0 Bq/kg dry	3986.0	Cs137	9.0 Bq/kg dry
			Cs134	166.0 Bq/kg dry	± 19.0 Bq/kg dry		Cs134	8.3 Bq/kg dry
Soil (in the park) under a maple tree	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	2550.0 Bq/kg dry	± 259.0 Bq/kg dry	2652.0	Cs137	4.2 Bq/kg dry
			Cs134	102.0 Bq/kg dry	± 11.4 Bq/kg dry		Cs134	3.9 Bq/kg dry
Soil (in the park)	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	907.0 Bq/kg dry	± 93.5 Bq/kg dry	952.4	Cs137	2.8 Bq/kg dry
			Cs134	45.4 Bq/kg dry	± 5.2 Bq/kg dry		Cs134	2.8 Bq/kg dry
Soil (in the park)	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	750.0 Bq/kg dry	± 77.5 Bq/kg dry	789.3	Cs137	3.2 Bq/kg dry
			Cs134	39.3 Bq/kg dry	± 4.7 Bq/kg dry		Cs134	3.5 Bq/kg dry
Soil (in the park)	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	577.0 Bq/kg dry	± 59.7 Bq/kg dry	599.7	Cs137	2.6 Bq/kg dry
			Cs134	22.7 Bq/kg dry	± 3.0 Bq/kg dry		Cs134	2.8 Bq/kg dry
Soil (in the park)	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	397.0 Bq/kg dry	± 40.7 Bq/kg dry	413.5	Cs137	1.4 Bq/kg dry
			Cs134	16.5 Bq/kg dry	± 2.0 Bq/kg dry		Cs134	1.6 Bq/kg dry
Soil (in the park) under a cherry-blossom tree	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	344.0 Bq/kg dry	± 35.3 Bq/kg dry	358.2	Cs137	1.3 Bq/kg dry
			Cs134	14.2 Bq/kg dry	± 1.8 Bq/kg dry		Cs134	1.5 Bq/kg dry
Soil (in the park) under the slide	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	183.0 Bq/kg dry	± 19.5 Bq/kg dry	189.9	Cs137	2.1 Bq/kg dry
			Cs134	6.9 Bq/kg dry	± 1.3 Bq/kg dry		Cs134	2.6 Bq/kg dry
Soil (in the park) under the Swing	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	96.7 Bq/kg dry	± 10.8 Bq/kg dry	101.1	Cs137	1.9 Bq/kg dry
			Cs134	4.4 Bq/kg dry	± 1.0 Bq/kg dry		Cs134	2.4 Bq/kg dry
Soil (in the park) In front of a bench	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	56.1 Bq/kg dry	± 6.0 Bq/kg dry	57.7	Cs137	0.9 Bq/kg dry
			Cs134	1.6 Bq/kg dry	± 0.4 Bq/kg dry		Cs134	1.2 Bq/kg dry
Soil (in the park) Sandbox	Noda-chuo Park Noda, Fukushima, Fukushima Pref.	Apr-21	Cs137	— Bq/kg dry	± — Bq/kg dry	Under Minimum Limit of Detection	Cs137	1.3 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.6 Bq/kg dry
Sea sand(surface)	Haragamaobama Beach ①, Fukushima Pef.	Apr-21	Cs137	6.4 Bq/kg dry	± 0.8 Bq/kg dry	6.4	Cs137	0.9 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.9 Bq/kg dry
Sea sand(15cm)	Haragamaobama Beach ①, Fukushima Pef.	Apr-21	Cs137	7.6 Bq/kg dry	± 0.9 Bq/kg dry	7.6	Cs137	0.7 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.8 Bq/kg dry

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Sea sand(30cm)	Haragamaobama Beach①, Fukushima Pref.	Apr-21	Cs137	— Bq/kg dry	± — Bq/kg dry	Under Minimum Limit of Detection	Cs137	0.9 Bq/kg dry
Sea sand(50cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.1 Bq/kg dry
Sea sand (surface)		Apr-21	Cs137	14.6 Bq/kg dry	± 1.9 Bq/kg dry	14.6	Cs137	1.5 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.1 Bq/kg dry
Sea sand(30cm)	Haragamaobama Beach②, Fukushima Pref.	Apr-21	Cs137	8.8 Bq/kg dry	± 1.1 Bq/kg dry	9.4	Cs137	0.5 Bq/kg dry
Sea sand(50cm)			Cs134	0.6 Bq/kg dry	± 0.2 Bq/kg dry		Cs134	0.6 Bq/kg dry
Sea sand (surface)		Apr-21	Cs137	17.0 Bq/kg dry	± 2.1 Bq/kg dry	17.0	Cs137	1.4 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.1 Bq/kg dry
Sea sand(30cm)	Apr-21	Cs137	34.3 Bq/kg dry	± 3.6 Bq/kg dry	35.2	Cs137	0.5 Bq/kg dry	
Sea sand(50cm)		Cs134	0.9 Bq/kg dry	± 0.2 Bq/kg dry		Cs134	0.5 Bq/kg dry	
Sea sand (surface)		Apr-21	Cs137	34.9 Bq/kg dry	± 3.9 Bq/kg dry	34.9	Cs137	1.3 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.0 Bq/kg dry
Sea sand(30cm)	Haragamaobama Beach③, Fukushima Pref.	Apr-21	Cs137	13.2 Bq/kg dry	± 1.5 Bq/kg dry	13.2	Cs137	0.5 Bq/kg dry
Sea sand(50cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.5 Bq/kg dry
Sea sand (surface)		Apr-21	Cs137	40.0 Bq/kg dry	± 4.3 Bq/kg dry	40.0	Cs137	1.0 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.7 Bq/kg dry
Sea sand(30cm)	Apr-21	Cs137	14.4 Bq/kg dry	± 1.8 Bq/kg dry	14.4	Cs137	1.3 Bq/kg dry	
Sea sand(50cm)		Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.0 Bq/kg dry	
Sea sand (surface)		Apr-21	Cs137	83.4 Bq/kg dry	± 9.0 Bq/kg dry	86.3	Cs137	1.1 Bq/kg dry
Sea sand(15cm)			Cs134	2.9 Bq/kg dry	± 0.5 Bq/kg dry		Cs134	1.3 Bq/kg dry
Sea sand(30cm)	Haragamaobama Beach④, Fukushima Pref.	Apr-21	Cs137	3.1 Bq/kg dry	± 0.4 Bq/kg dry	3.1	Cs137	0.5 Bq/kg dry
Sea sand(50cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.5 Bq/kg dry
Sea sand (surface)		Apr-21	Cs137	5.0 Bq/kg dry	± 0.6 Bq/kg dry	5.0	Cs137	0.7 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.7 Bq/kg dry
Sea sand(30cm)	Apr-21	Cs137	2.3 Bq/kg dry	± 0.5 Bq/kg dry	2.3	Cs137	1.3 Bq/kg dry	
Sea sand(50cm)		Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.3 Bq/kg dry	
Sea sand (surface)		Apr-21	Cs137	2.0 Bq/kg dry	± 0.4 Bq/kg dry	2.0	Cs137	1.1 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.1 Bq/kg dry
Sea sand(30cm)	Yotsukura Beach① Fukushima Pref.	Apr-21	Cs137	18.6 Bq/kg dry	± 2.1 Bq/kg dry	18.6	Cs137	0.6 Bq/kg dry
Sea sand(50cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.8 Bq/kg dry
Sea sand (surface)		Apr-21	Cs137	10.1 Bq/kg dry	± 1.2 Bq/kg dry	10.1	Cs137	0.6 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.6 Bq/kg dry
Sea sand(30cm)	Yotsukura Beach② Fukushima Pref.	Apr-21	Cs137	6.7 Bq/kg dry	± 0.8 Bq/kg dry	6.7	Cs137	0.6 Bq/kg dry
Sea sand(50cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.6 Bq/kg dry
Sea sand (surface)		Apr-21	Cs137	7.7 Bq/kg dry	± 1.0 Bq/kg dry	7.7	Cs137	1.1 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.1 Bq/kg dry
Sea sand(30cm)	Apr-21	Cs137	11.1 Bq/kg dry	± 1.5 Bq/kg dry	11.1	Cs137	1.4 Bq/kg dry	
Sea sand(50cm)		Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.4 Bq/kg dry	
Sea sand (surface)	Yotsukura Beach③ Fukushima Pref.	Apr-21	Cs137	21.6 Bq/kg dry	± 2.5 Bq/kg dry	22.6	Cs137	0.6 Bq/kg dry
Sea sand(15cm)			Cs134	1.0 Bq/kg dry	± 0.3 Bq/kg dry		Cs134	0.7 Bq/kg dry
Sea sand(30cm)	Apr-21	Cs137	11.2 Bq/kg dry	± 1.3 Bq/kg dry	11.2	Cs137	0.5 Bq/kg dry	
Sea sand(50cm)		Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.6 Bq/kg dry	
Sea sand (surface)	Yotsukura Beach④ Fukushima Pref.	Apr-21	Cs137	11.6 Bq/kg dry	± 1.3 Bq/kg dry	11.6	Cs137	0.6 Bq/kg dry
Sea sand(15cm)			Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	0.6 Bq/kg dry
Sea sand(30cm)	Apr-21	Cs137	7.0 Bq/kg dry	± 1.0 Bq/kg dry	7.0	Cs137	1.1 Bq/kg dry	
Sea sand(50cm)		Cs134	— Bq/kg dry	± — Bq/kg dry		Cs134	1.1 Bq/kg dry	
Sea sand (surface)	Apr-21	Cs137	32.4 Bq/kg dry	± 3.4 Bq/kg dry	33.3	Cs137	0.6 Bq/kg dry	
Sea sand(15cm)		Cs134	0.9 Bq/kg dry	± 0.2 Bq/kg dry		Cs134	0.6 Bq/kg dry	

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Sea sand(15cm)	Yotsukura Beach④ Fukushima Pref.	Apr-21	Cs137	19.2	Bq/kg dry ± 2.4 Bq/kg dry	19.2	Cs137	1.4 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	1.1 Bq/kg dry
		Apr-21	Cs137	26.9	Bq/kg dry ± 3.0 Bq/kg dry	26.9	Cs137	0.9 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	0.7 Bq/kg dry
		Apr-21	Cs137	19.0	Bq/kg dry ± 2.1 Bq/kg dry	19.0	Cs137	0.6 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	0.7 Bq/kg dry
		Apr-21	Cs137	33.4	Bq/kg dry ± 3.6 Bq/kg dry	35.6	Cs137	0.6 Bq/kg dry
			Cs134	2.2	Bq/kg dry ± 0.4 Bq/kg dry		Cs134	0.6 Bq/kg dry
Sea sand(15cm)	Yotsukura Beach⑤ Fukushima Pref.	Apr-21	Cs137	27.2	Bq/kg dry ± 2.9 Bq/kg dry	27.2	Cs137	0.6 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	0.6 Bq/kg dry
		Apr-21	Cs137	18.4	Bq/kg dry ± 2.0 Bq/kg dry	18.4	Cs137	0.6 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	0.6 Bq/kg dry
		Apr-21	Cs137	15.6	Bq/kg dry ± 1.7 Bq/kg dry	15.6	Cs137	0.6 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	0.6 Bq/kg dry
		Apr-21	Cs137	16.9	Bq/kg dry ± 2.0 Bq/kg dry	16.9	Cs137	1.2 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	1.1 Bq/kg dry
Sea sand(15cm)	Yotsukura Beach⑥ Fukushima Pref.	Apr-21	Cs137	34.2	Bq/kg dry ± 4.0 Bq/kg dry	34.2	Cs137	1.7 Bq/kg dry
			Cs134	—	Bq/kg dry ± — Bq/kg dry		Cs134	1.3 Bq/kg dry
		Apr-21	Cs137	73.7	Bq/kg dry ± 7.9 Bq/kg dry	75.8	Cs137	1.1 Bq/kg dry
			Cs134	2.1	Bq/kg dry ± 0.4 Bq/kg dry		Cs134	1.3 Bq/kg dry
		Apr-21	Cs137	129.0	Bq/kg dry ± 13.9 Bq/kg dry	134.3	Cs137	1.4 Bq/kg dry
			Cs134	5.3	Bq/kg dry ± 0.9 Bq/kg dry		Cs134	1.4 Bq/kg dry
Ash (Laurier tree)	Izumigaoka, Iwaki	May-21	Cs137	554.0	Bq/kg raw ± 48.6 Bq/kg raw	571.8	Cs137	3.5 Bq/kg raw
			Cs134	17.8	Bq/kg raw ± 4.3 Bq/kg raw		Cs134	2.9 Bq/kg raw
Vacuum cleaner dust (paper pack, Roomba)	Ichikawa, Chiba	Apr-21	Cs137	33.3	Bq/kg raw ± 9.0 Bq/kg raw	33.3	Cs137	7.0 Bq/kg raw
Dust cloth (After cleaning of sash)	Ichikawa, Chiba	Apr-21	Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134	6.0 Bq/kg raw
			Cs137	14.1	Bq/kg raw ± 4.7 Bq/kg raw	14.1	Cs137	5.3 Bq/kg raw
			Cs134	—	Bq/kg raw ± — Bq/kg raw		Cs134	4.2 Bq/kg raw

\*"—" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



## ★Gamma-ray

Measuring instrument		Feature	Guide to lower limit※
NaI Scintillation Spectrometer			
Product of ATOMTEX AT1320A	Product of BERTHOLD LB2045	• Gamma-ray spectrometer with NaI scintillation detector.	Food (Sample 1kg) Lower limit 1.0Bq/Kg Soil (Sample 1kg) Lower limit 2.5Bq/Kg Material (Sample 1kg) Lower limit 1.0Bq/Kg Water (Sample 20L) Lower limit 0.02Bq/L
Germanium Semiconductor detector			
ORTEC GEM30-70		• Radioactivity measurement series. Quantitative analysis based on "Gamma-ray spectrometry with germanium semiconductor detector." • Relative efficiency 35%	Food (Sample 2kg) Lower limit 0.04Bq/Kg Soil (Sample 1kg) Lower limit 0.06Bq/Kg Material (Sample 1kg) Lower limit 0.06Bq/Kg Water (Sample 20L) Lower limit 0.001Bq/L

※The lower limit varies depending on the sample weight and measurement time.

Measuring instrument: Germanium Semiconductor detector (Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result	Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Brown rice	Esashi, Osyu, Iwate	Oct-20	Cs137 0.68 Bq/kg raw	± 0.05 Bq/kg raw	<b>0.68</b>	Cs137 0.08 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.08 Bq/kg raw
Rice bran	Gotenba, Shizuoka	Apr-21	Cs137 11.6 Bq/kg raw	± 0.3 Bq/kg raw	<b>11.6</b>	Cs137 0.3 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.3 Bq/kg raw
Japanese honeywort	Izumi, Iwaki	May-21	Cs137 1.5 Bq/kg raw	± 0.09 Bq/kg raw	<b>1.5</b>	Cs137 0.1 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.1 Bq/kg raw
Plum(pulp)	Izumigaoka, Iwaki	May-21	Cs137 0.34 Bq/kg raw	± 0.08 Bq/kg raw	<b>0.34</b>	Cs137 0.1 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.1 Bq/kg raw
Dried zenmai (Wild)	Minamiaizuchō, Minamiaizu, Fukushima	May-11	Cs137 1.4 Bq/kg raw	± 0.3 Bq/kg raw	<b>1.4</b>	Cs137 0.6 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.6 Bq/kg raw
Dried zenmai (Wild)	Minamiaizuchō, Minamiaizu, Fukushima	May-18	Cs137 441.1 Bq/kg raw	± 2.3 Bq/kg raw	<b>453.7</b>	Cs137 0.8 Bq/kg raw
			Cs134 12.6 Bq/kg raw	± 0.5 Bq/kg raw		Cs134 0.8 Bq/kg raw
Dried zenmai (Wild)	Minamiaizuchō, Minamiaizu, Fukushima	May-19	Cs137 56.8 Bq/kg raw	± 0.8 Bq/kg raw	<b>59.4</b>	Cs137 0.7 Bq/kg raw
			Cs134 2.6 Bq/kg raw	± 0.3 Bq/kg raw		Cs134 0.6 Bq/kg raw
Bracken(Wild)	Miyakojī, Tamura, Fukushima (Old road)	May-21	Cs137 56.7 Bq/kg raw	± 1.2 Bq/kg raw	<b>59.0</b>	Cs137 0.6 Bq/kg raw
			Cs134 2.3 Bq/kg raw	± 0.4 Bq/kg raw		Cs134 0.7 Bq/kg raw
Dried bracken (Wild)	Minamiaizuchō, Minamiaizu, Fukushima	May-11	Cs137 15.2 Bq/kg raw	± 0.3 Bq/kg raw	<b>15.9</b>	Cs137 0.4 Bq/kg raw
			Cs134 0.7 Bq/kg raw	± 0.2 Bq/kg raw		Cs134 0.4 Bq/kg raw
Bracken(Wild)	Shimogo, Minamiaizu, Fukushima	May-21	Cs137 1.1 Bq/kg raw	± 0.1 Bq/kg raw	<b>1.1</b>	Cs137 0.2 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.2 Bq/kg raw
Bracken(Wild)	Izumi, Iwaki	May-21	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.1 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.2 Bq/kg raw
Bracken(Wild)	Esashi, Osyu, Iwate	May-21	Cs137 2.6 Bq/kg raw	± 0.1 Bq/kg raw	<b>2.6</b>	Cs137 0.2 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.2 Bq/kg raw
Bracken(Wild) After removing bitter taste	Esashi, Osyu, Iwate	May-21	Cs137 0.4 Bq/kg raw	± 0.1 Bq/kg raw	<b>0.4</b>	Cs137 0.2 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.2 Bq/kg raw
Bracken	Yamagata Pref.	Apr-21	Cs137 — Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.7 Bq/kg raw
			Cs134 — Bq/kg raw	± — Bq/kg raw		Cs134 0.7 Bq/kg raw
Koshiabura	Unknown (Purchase of Inawashiro)	May-21	Cs137 100.2 Bq/kg raw	± 1.9 Bq/kg raw	<b>104.3</b>	Cs137 1.2 Bq/kg raw
			Cs134 4.1 Bq/kg raw	± 0.6 Bq/kg raw		Cs134 1.2 Bq/kg raw
Koshiabura(Wild)	Osawa, Yonezawa, Yamagata	May-21	Cs137 75.4 Bq/kg raw	± 2.5 Bq/kg raw	<b>78.5</b>	Cs137 2.3 Bq/kg raw
			Cs134 3.1 Bq/kg raw	± 1.3 Bq/kg raw		Cs134 2.6 Bq/kg raw

\* "\_" used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessarily mean 0(zero)Bq/kg.

\* Dried royal fern and dried bracken in Minami-Aizu, Minami-Aizu, Fukushima, as of the measurement date May 2021.

## ★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result			Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Koshiabura(Wild)	Esashi,Osyu, Iwate	May-21	Cs137	16.7	Bq/kg raw	± 1.4	Bq/kg raw	16.7	Cs137 1.3 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 1.5 Bq/kg raw
New leaves of Koshiabura (Wild)	Onami,Fukushima, Fukushima Pref. (In the mountains)	May-21	Cs137	2598.1	Bq/kg raw	± 7.3	Bq/kg raw	2705.3	Cs137 1.6 Bq/kg raw
			Cs134	107.2	Bq/kg raw	± 1.6	Bq/kg raw		Cs134 1.7 Bq/kg raw
New leaves of Koshiabura (Wild)	Onami,Fukushima, Fukushima Pref. (In the garden)	May-21	Cs137	7.2	Bq/kg raw	± 0.2	Bq/kg raw	7.2	Cs137 0.3 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.2 Bq/kg raw
New leaves of Koshiabura (Wild)	Marumori,Igu, Miyagi	May-21	Cs137	317.1	Bq/kg raw	± 9.5	Bq/kg raw	326.4	Cs137 5.9 Bq/kg raw
			Cs134	9.3	Bq/kg raw	± 2.6	Bq/kg raw		Cs134 4.9 Bq/kg raw
Aralia sprout(Wild)	Sakuragi, Fukushima, Fukushima Pref.	May-21	Cs137	85.9	Bq/kg raw	± 3.4	Bq/kg raw	89.1	Cs137 2.7 Bq/kg raw
			Cs134	3.2	Bq/kg raw	± 1.3	Bq/kg raw		Cs134 2.6 Bq/kg raw
Aralia sprout(Wild)	Miyakoji,Tamura, Fukushima (Old road)	May-21	Cs137	148.7	Bq/kg raw	± 4.0	Bq/kg raw	154.3	Cs137 2.1 Bq/kg raw
			Cs134	5.6	Bq/kg raw	± 1.2	Bq/kg raw		Cs134 2.1 Bq/kg raw
Aralia sprout(Wild)	Ono,Tamura, Fukushima	May-21	Cs137	6.8	Bq/kg raw	± 0.3	Bq/kg raw	6.8	Cs137 0.5 Bq/kg raw
			—	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.6 Bq/kg raw
Aralia sprout(Wild)	Kamiogawa, Ogawa,Iwaki	Apr-21	Cs137	15.2	Bq/kg raw	± 1.2	Bq/kg raw	15.2	Cs137 2.0 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 2.5 Bq/kg raw
Aralia sprout(Wild)	Osawa,Yonezawa, Yamagata	May-21	Cs137	2.3	Bq/kg raw	± 0.5	Bq/kg raw	2.3	Cs137 1.1 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 1.1 Bq/kg raw
Ostrich fern (Wild)	Minamiaizu, Minamiaizu, Fukushima	May-21	Cs137	2.0	Bq/kg raw	± 0.2	Bq/kg raw	2.0	Cs137 0.3 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.4 Bq/kg raw
Shidoke(Wild)	Miyakoji,Tamura, Fukushima (Old road)	May-21	Cs137	178.2	Bq/kg raw	± 3.8	Bq/kg raw	184.9	Cs137 1.3 Bq/kg raw
			Cs134	6.7	Bq/kg raw	± 1.0	Bq/kg raw		Cs134 1.6 Bq/kg raw
Butterbur sprout (Wild)	Minamiaizuchi, Minamiaizu, Fukushima	Mar-21	Cs137	0.35	Bq/kg raw	± 0.08	Bq/kg raw	0.35	Cs137 0.1 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.1 Bq/kg raw
Butterbur	Tairashimokabeya, Iwaki	Apr-21	Cs137	—	Bq/kg raw	± —	Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.2 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.2 Bq/kg raw
Japanese mugwort	Minamiaizuchi, Minamiaizu, Fukushima	May-21	Cs137	—	Bq/kg raw	± —	Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.3 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.4 Bq/kg raw
Nobiru	Minamiaizuchi, Minamiaizu, Fukushima	Mar-21	Cs137	—	Bq/kg raw	± —	Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.2 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.3 Bq/kg raw
Shitake mushroom log grown	Ogawa,Iwaki	May-21	Cs137	6.5	Bq/kg raw	± 0.2	Bq/kg raw	6.5	Cs137 0.2 Bq/kg raw
			Cs134	—	Bq/kg raw	± —	Bq/kg raw		Cs134 0.2 Bq/kg raw
Black buss①	Iitate,Soma, Fukushima	Apr-21	Cs137	78.7	Bq/kg raw	± 0.8	Bq/kg raw	82.0	Cs137 0.3 Bq/kg raw
			Cs134	3.3	Bq/kg raw	± 0.2	Bq/kg raw		Cs134 0.4 Bq/kg raw
Black buss②	Iitate,Soma, Fukushima	Apr-21	Cs137	65.7	Bq/kg raw	± 0.9	Bq/kg raw	68.7	Cs137 0.4 Bq/kg raw
			Cs134	3.0	Bq/kg raw	± 0.2	Bq/kg raw		Cs134 0.4 Bq/kg raw

※"\_"used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



## ★Beta-ray

Measuring instrument		Feature
Liquid Scintillation Counter		
Product of Hidex HIDEX 300SSL	Product of PerkinElmer Japan Quantulus GCT 6220	<p>Equipment for measuring low-energy beta-ray emission nuclides</p> <p>Measuring nuclide</p> <p>Strontium90      Half-life 30 years</p> <p>Organic bound      Half-life 12.3 years</p> <p>Free-water tritium      Half-life 12.3 years</p> <p>All samples are measured in liquid condition after several days of pretreatment.</p>
		

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty	Minimum Limit of Detection	
Tap water	Onahama, hanabatake, Iwaki	Apr-21	T (Free)	0.29	Bq/L	± 0.16	Bq/L
Sea water (surface)	Futabacho,Futaba, Fukushima	Apr-21	T (Free)	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Sea water (surface)	Soma Port/ Fukushima Pref.	Apr-21	T (Free)	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Sea water A (surface)	Off the coast of Fukushima Nuclear Power Plant1	May-21	T (Free)	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Sea water B (surface)	Off the coast of Fukushima Nuclear Power Plant1	May-21	T (Free)	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Flounder(pulp)	Oshima,Fukui, Fukui Pref.	Mar-21	T (Organic)	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Wakame seaweed	Oi-cho, Oi,Fukui	Mar-21	T (Organic)	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Yellowtail	Off the coast of Fukushima Nuclear Power Plant1	Nov-19	Sr90	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Flounder(bone)	Oshima,Fukui, Fukui Pref.	Mar-21	Sr90	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Boar(bone)	Iwaki city	Sep-14	Sr90	25.34	Bq/kg dry	± 1.33	Bq/kg dry
Soil	Takada Park Onahamahanabatake, Iwaki	Apr-20	Sr90	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Soil	Onahama-chuo Park Onahama-minami	Mar-21	Sr90	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Soil	Tomigaura Park Onahamagotenba,Iwaki	Mar-20	Sr90	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Soil	Shonandai-Nishi Park 1Shonandai,Iwaki	Mar-20	Sr90	Under Minimum Limit of Detection	Bq/kg dry	± —	Bq/kg dry
Tap water	Hansaki,Odaka, Minamisoma, Fukushima	Apr-21	Sr90	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Groundwater	Futabacho, Futaba,Fukushima	Apr-21	Sr90	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Tap water	Onahama, hanabatake, Iwaki	Apr-21	Sr90	Under Minimum Limit of Detection	Bq/L	± —	Bq/L
Tap water	Minamidai, Iwaki	Apr-21	Sr90	Under Minimum Limit of Detection	Bq/L	± —	Bq/L

# Measurement results of 16 items by germanium semiconductor detector


Dr.Tetsuji Imanaka, Institute of Multiple Nuclear Science, Kyoto University

In order to convey more measurement results to everyone, we have asked Dr. Tetsuji Imanaka of the Institute of Advanced Nuclear Science, Kyoto University, to measure low-dose samples using germanium semiconductor detectors. Measurement samples are not only from Fukushima Prefecture but also come from other prefectures. Please compare data based on measurements from various regions and use them to protect your children from radiation exposure.

## ★Gamma-ray

Measuring instrument : Germanium Semiconductor detector

- Product of CANBERRA(CA),USA GX3018      Relative efficiency 30% or more
- Product of ORTEC(OR),USA GMX25-70      Relative efficiency 35%



(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measuring instrument type	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection
Rice	Iitate,Soma, Fukushima	Oct-20	OR	Cs137	0.2 Bq/kg raw	± 0.03 Bq/kg raw	0.2	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Rice	Shinti,Soma, Fukushima	Oct-20	OR	Cs137	0.3 Bq/kg raw	± 0.04 Bq/kg raw	0.3	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Rice	Miyagi Pref.	Oct-20	OR	Cs137	0.02 Bq/kg raw	± 0.01 Bq/kg raw	0.02	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Sweet potato	Miura,Inashiki, Ibaraki	Feb-21	OR	Cs137	0.5 Bq/kg raw	± 0.07 Bq/kg raw	0.5	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Green onion	Iwaki city	Mar-21	OR	Cs137	0.5 Bq/kg raw	± 0.09 Bq/kg raw	0.5	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Broccoli	Iitate,Soma, Fukushima	Feb-21	OR	Cs137	0.03 Bq/kg raw	± 0.02 Bq/kg raw	0.03	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Yam bulblet	Tomioka,Futaba, Fukushima	Feb-21	CA	Cs137	2.9 Bq/kg raw	± 0.1 Bq/kg raw	3.04	Cs137 — Bq/kg raw
				Cs134	0.14 Bq/kg raw	± 0.03 Bq/kg raw		Cs134 — Bq/kg raw
Japanese yam	Hirata,Ishikawa, Fukushima	Jan-21	OR	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137 0.1 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Butterbur sprout (cultivation)	Nishigo, Nishishirakawa, Fukushima	Mar-21	OR	Cs137	10.6 Bq/kg raw	± 0.3 Bq/kg raw	10.9	Cs137 — Bq/kg raw
				Cs134	0.3 Bq/kg raw	± 0.1 Bq/kg raw		Cs134 — Bq/kg raw
Butterbur sprout (cultivation)	Asakawa,Ishikawa, Fukushima	Mar-21	OR	Cs137	0.9 Bq/kg raw	± 0.13 Bq/kg raw	0.9	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Butterbur sprout (cultivation)	Funehiki, Tamura,Fukushima	Mar-21	CA	Cs137	0.5 Bq/kg raw	± 0.09 Bq/kg raw	0.5	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Butterbur sprout (cultivation)	Shimogo, Minamiaizu, Fukushima	Mar-21	CA	Cs137	0.3 Bq/kg raw	± 0.06 Bq/kg raw	0.3	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Shidoke (cultivation)	Furudono, Ishikawa,, Fukushima	Mar-21	CA	Cs137	0.45 Bq/kg raw	± 0.08 Bq/kg raw	0.45	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Shitake mushroom grown in bacteria-bed	Kawauchi,Futaba, Fukushima	Feb-21	OR	Cs137	0.2 Bq/kg raw	± 0.04 Bq/kg raw	0.2	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Shitake mushroom grown in bacteria-bed	Minamiuonuma, Miyagi	Feb-21	CA	Cs137	0.6 Bq/kg raw	± 0.05 Bq/kg raw	0.6	Cs137 — Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134 — Bq/kg raw
Kiwi fruit	Izumizaki, Nishishirakawa, Fukushima	Mar-21	OR	Cs137	1.9 Bq/kg raw	± 0.06 Bq/kg raw	1.95	Cs137 — Bq/kg raw
				Cs134	0.05 Bq/kg raw	± 0.02 Bq/kg raw		Cs134 — Bq/kg raw