



Radiation Measurement Results of 131 Items in July



When samples include natural radionuclides we can't deny the possibility of their radiation value counted together in our results.

The list below only shows the measurement results of the samples brought in.

Radioactive contamination level may differ according to sampling points even within the same address.

★Gamma-ray

Measuring instrument		Feature	Guide to lower limit※
Na I Scintillation Spectrometer			
Product of ATOMTEX AT1320A	Product of BERTHOLD LB2045	• Gamma-ray spectrometer with Na I scintillation detector.	Food (Sample 1kg) Lower limit 1.0Bq/Kg
			Soil (Sample 1kg) Lower limit 2.5Bq/Kg
			Material (Sample 1kg) Lower limit 1.0Bq/Kg
			Water (Sample 20L) Lower limit 0.02Bq/L

※The lower limit varies depending on the sample weight and measurement time.

Measuring instrument: Na I Scintillation Spectrometer (Bq/kg raw: Weight of raw sample Bq/kg dry: Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Sea sand (surface)	Iwasawa Coast① Hirono, Futaba, Fukushima perf.	Jun-25	Cs137	19.4 Bq/kg dry	± 2.1 Bq/kg dry		19.4	Cs137	0.7 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	0.8 Bq/kg dry
Sea sand (15cm deep)		Jun-25	Cs137	— Bq/kg dry	± — Bq/kg dry		Under Minimum Limit of Detection	Cs137	2.5 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	2.2 Bq/kg dry
Sea sand (30cm deep)		Jun-25	Cs137	14.2 Bq/kg dry	± 1.8 Bq/kg dry		14.2	Cs137	1.7 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	1.8 Bq/kg dry
Sea sand (50cm deep)		Jun-25	Cs137	13.7 Bq/kg dry	± 1.6 Bq/kg dry		13.7	Cs137	0.8 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	0.8 Bq/kg dry
Sea sand (surface)	Iwasawa Coast② Hirono, Futaba, Fukushima perf.	Jun-25	Cs137	20.3 Bq/kg dry	± 2.2 Bq/kg dry		20.3	Cs137	0.7 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	0.7 Bq/kg dry
Sea sand (15cm deep)		Jun-25	Cs137	14.9 Bq/kg dry	± 1.9 Bq/kg dry		14.9	Cs137	1.6 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	1.7 Bq/kg dry
Sea sand (30cm deep)		Jun-25	Cs137	23.8 Bq/kg dry	± 2.6 Bq/kg dry		23.8	Cs137	0.7 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	0.7 Bq/kg dry
Sea sand (50cm deep)		Jun-25	Cs137	24.7 Bq/kg dry	± 2.9 Bq/kg dry		24.7	Cs137	1.4 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	1.4 Bq/kg dry
Soil (under the slide)	Nagamachi Park, Mimaya, Uchigou, Iwaki	Mar-25	Cs137	216.0 Bq/kg dry	± 23.0 Bq/kg dry		216.0	Cs137	3.4 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	3.4 Bq/kg dry
Soil (spring play equipment)	Nagamachi Park, Mimaya, Uchigou, Iwaki	Mar-25	Cs137	149.0 Bq/kg dry	± 15.5 Bq/kg dry		149.0	Cs137	1.3 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	1.3 Bq/kg dry
Soil (under the swing)	Nagamachi Park, Mimaya, Uchigou, Iwaki	Mar-25	Cs137	103.0 Bq/kg dry	± 11.3 Bq/kg dry		103.0	Cs137	2.1 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	2.2 Bq/kg dry
Soil (in the park)	Nagamachi Park, Mimaya, Uchigou, Iwaki	Mar-25	Cs137	32.0 Bq/kg dry	± 3.8 Bq/kg dry		32.0	Cs137	2.0 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	2.0 Bq/kg dry
Soil (sand box)	Nagamachi Park, Mimaya, Uchigou, Iwaki	Mar-25	Cs137	— Bq/kg dry	± — Bq/kg dry		Under Minimum Limit of Detection	Cs137	0.8 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	0.7 Bq/kg dry
Soil (in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	574.0 Bq/kg dry	± 58.6 Bq/kg dry		579.2	Cs137	1.5 Bq/kg dry
			Cs134	5.2 Bq/kg dry	± 0.8 Bq/kg dry			Cs134	1.4 Bq/kg dry
Soil (in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	355.0 Bq/kg dry	± 37.0 Bq/kg dry		360.0	Cs137	2.1 Bq/kg dry
			Cs134	5.0 Bq/kg dry	± 1.0 Bq/kg dry			Cs134	2.1 Bq/kg dry
Soil (in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	331.0 Bq/kg dry	± 34.7 Bq/kg dry		331.0	Cs137	3.6 Bq/kg dry
			Cs134	— Bq/kg dry	± — Bq/kg dry			Cs134	2.8 Bq/kg dry

But it does not necessary mean 0(zero)Bq/kg.


★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Soil(in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	180.0 Bq/kg dry	±	19.1 Bq/kg dry	180.0	Cs137	2.7 Bq/kg dry
			Cs134	— Bq/kg dry	±	— Bq/kg dry		Cs134	2.0 Bq/kg dry
Soil(in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	156.0 Bq/kg dry	±	17.0 Bq/kg dry	156.0	Cs137	3.6 Bq/kg dry
			Cs134	— Bq/kg dry	±	— Bq/kg dry		Cs134	2.6 Bq/kg dry
Soil(in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	154.0 Bq/kg dry	±	15.9 Bq/kg dry	154.0	Cs137	1.5 Bq/kg dry
			Cs134	— Bq/kg dry	±	— Bq/kg dry		Cs134	1.1 Bq/kg dry
Soil(in the park)	Futabamachi Park, Tairayagawase, Iwaki	Mar-25	Cs137	— Bq/kg dry	±	— Bq/kg dry	Under Minimum Limit of Detection	Cs137	2.3 Bq/kg dry
			Cs134	— Bq/kg dry	±	— Bq/kg dry		Cs134	1.9 Bq/kg dry

※“—” used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



★Gamma-ray

Measuring instrument		Feature	Guide to lower limit※	
Germanium Semiconductor detector				
ORTEC GEM30-70	CANBERRA GC4020	• Radioactivity measurement series. - Quantitative analysis based on "Gamma-ray spectrometry with germanium semiconductor detector." • ORTEC GEM30-70 Relative efficiency 35% • CANBERRA GC4020 Relative efficiency 43%	Food (Sample 2kg)	Lower limit 0.04Bq/Kg
			Soil (Sample 1kg)	Lower limit 0.06Bq/Kg
			Material (Sample 1kg)	Lower limit 0.06Bq/Kg
			Water (Sample 20L)	Lower limit 0.001Bq/L

※The lower limit varies depending on the sample weight and measurement time.

Measuring instrument: Germanium Semiconductor detector

(Bq/kg raw: Weight of raw sample Bq/kg dry: Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measuring instrument type	Measurement Result		Uncertainty	Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Souma, Fukushima Pref.	Oct-24	CA	Cs137	0.9 Bq/kg raw	± 0.03 Bq/kg raw	0.9	Cs137	0.05 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.05 Bq/kg raw
Brown rice (half-milled rice)	Souma, Fukushima Pref.	Oct-24	CA	Cs137	0.2 Bq/kg raw	± 0.02 Bq/kg raw	0.2	Cs137	0.04 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.05 Bq/kg raw
Glutinous rice	Fukushima Pref.	Oct-24	CA	Cs137	0.4 Bq/kg raw	± 0.05 Bq/kg raw	0.4	Cs137	0.09 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.1 Bq/kg raw
Potato	Ryouzen, Date, Fukushima Pref.	Jun-25	OR	Cs137	0.2 Bq/kg raw	± 0.1 Bq/kg raw	0.2	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.2 Bq/kg raw
Eggplant	Nishida, Koriyama, Fukushima Pref.	Jul-25	CA	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.2 Bq/kg raw
Eggplant	Funehiki, Tamura, Fukushima Pref.	Jul-25	CA	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.2 Bq/kg raw
Pumpkin	Sizuoka Pref.	Jul-25	OR	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.2 Bq/kg raw
Cauliflower	Otsuki, Koriyama, Fukushima Pref.	Jul-25	CA	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.1 Bq/kg raw
common bean	Fukushima, Fukushima Pref.	Jun-25	OR	Cs137	1.3 Bq/kg raw	± 0.2 Bq/kg raw	1.3	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.4 Bq/kg raw
common bean	Tukidate, Date, Fukushima Pref.	Jun-25	OR	Cs137	0.6 Bq/kg raw	± 0.03 Bq/kg raw	0.6	Cs137	0.06 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.07 Bq/kg raw
Water spinach	Ryouzen, Date, Fukushima Pref.	Jun-25	CA	Cs137	0.8 Bq/kg raw	± 0.05 Bq/kg raw	0.8	Cs137	0.09 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.1 Bq/kg raw
Water spinach	Date, Fukushima Pref.	Jun-25	CA	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.1 Bq/kg raw
Water spinach	Date, Fukushima Pref.	Jun-25	OR	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.3 Bq/kg raw
Malabar spinach	Date, Fukushima Pref.	Jun-25	OR	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.3 Bq/kg raw
Moloheiya	Date, Fukushima Pref.	Jun-25	CA	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.1 Bq/kg raw
Rhubarb	Fukushima, Fukushima Pref.	Jun-25	OR	Cs137	1.1 Bq/kg raw	± 0.2 Bq/kg raw	1.1	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.3 Bq/kg raw
Anredera cordifolia	kunimi, Date, Fukushima Pref.	Jun-25	OR	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.4 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.5 Bq/kg raw
Saltwort	Kori, Date, Fukushima Pref.	Jun-25	OR	Cs137	0.1 Bq/kg raw	± 0.04 Bq/kg raw	0.1	Cs137	0.08 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.09 Bq/kg raw
Green soybeans	Date, Fukushima Pref.	Jun-25	OR	Cs137	— Bq/kg raw	± — Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	± — Bq/kg raw		Cs134	0.3 Bq/kg raw

※"—"used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measuring instrument type	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Zucchini	Asaka, Koriyama, Fukushima Pref.	Jul-25	CA	Cs137	2.2 Bq/kg raw	±	0.1 Bq/kg raw	2.2	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.1 Bq/kg raw
Zucchini	Kori, Date, Fukushima Pref.	Jun-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.2 Bq/kg raw
Cucumber	Ouse, Koriyama, Fukushima Pref.	Jul-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.1 Bq/kg raw
Dried stems of taro	Japan (production)	Jun-25	OR	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	5.4 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	5.6 Bq/kg raw
Ume	Funehiki, Tamura, Fukushima Pref.	Jul-25	CA	Cs137	1.2 Bq/kg raw	±	0.1 Bq/kg raw	1.2	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.1 Bq/kg raw
Ume	Mihota, Koriyama, Fukushima Pref.	Jul-25	OR	Cs137	0.6 Bq/kg raw	±	0.1 Bq/kg raw	0.6	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.2 Bq/kg raw
Ume	Fukushima, Fukushima Pref.	Jun-25	OR	Cs137	0.5 Bq/kg raw	±	0.1 Bq/kg raw	0.5	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.2 Bq/kg raw
Tomato	Ouse, Koriyama, Fukushima Pref.	Jul-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.1 Bq/kg raw
Artichoke	Fukushima, Fukushima Pref.	Jun-25	OR	Cs137	0.3 Bq/kg raw	±	0.1 Bq/kg raw	0.3	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.2 Bq/kg raw
Fig	Izumigaoka, Iwaki	Jul-25	OR	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.5 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.5 Bq/kg raw
Loquat	Kunimi, Date, Fukushima Pref.	Jun-25	CA	Cs137	0.9 Bq/kg raw	±	0.1 Bq/kg raw	0.9	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.2 Bq/kg raw
Loquat	Izumigaoka, Iwaki	Jul-25	OR	Cs137	0.4 Bq/kg raw	±	0.03 Bq/kg raw	0.4	Cs137	0.06 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.06 Bq/kg raw
Plum	Koriyama, Fukushima Pref.	Jul-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.1 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.1 Bq/kg raw
Apricot	Kori, Date, Fukushima Pref.	Jun-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.1 Bq/kg raw
Raspberry	Fukushima Pref.	Jun-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.4 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.4 Bq/kg raw
Butterbur (cultivation)	Cichigasyuku, Katta, Miyagi Pref.	Jun-25	CA	Cs137	0.4 Bq/kg raw	±	0.1 Bq/kg raw	0.4	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.3 Bq/kg raw
Shiitake mushroom (powder)	Japan (production)	Jun-25	OR	Cs137	66.6 Bq/kg raw	±	3.5 Bq/kg raw	66.6	Cs137	3.7 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.6 Bq/kg raw
Wood ear mushroom (dried)	Fukushima, Fukushima Pref.	Jun-25	CA	Cs137	14.7 Bq/kg raw	±	1.70 Bq/kg raw	14.7	Cs137	2.6 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	3.0 Bq/kg raw
Fox jacopever	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	1.2 Bq/kg raw	±	0.1 Bq/kg raw	1.2	Cs137	0.2 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.3 Bq/kg raw
Fox jacopever	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	0.7 Bq/kg raw	±	0.1 Bq/kg raw	0.7	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.3 Bq/kg raw
pufferfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	1.9 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	2.0 Bq/kg raw
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	CA	Cs137	0.5 Bq/kg raw	±	0.1 Bq/kg raw	0.5	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.2 Bq/kg raw
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	0.5 Bq/kg raw	±	0.1 Bq/kg raw	0.5	Cs137	0.3 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.3 Bq/kg raw
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	0.8 Bq/kg raw	±	0.3 Bq/kg raw	0.8	Cs137	0.6 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	0.6 Bq/kg raw

※“_”used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0 (zero) Bq/kg.

★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measuring instrument type	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
				Cs137	Bq/kg raw	±	Bq/kg raw		Cs137	Bq/kg raw
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	0.4	±	0.1	0.4	Cs137	0.2
				Cs134	—	±	—		Cs134	0.3
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	CA	Cs137	1.4	±	0.2	1.4	Cs137	0.4
				Cs134	—	±	—		Cs134	0.4
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	CA	Cs137	0.6	±	0.08	0.6	Cs137	0.1
				Cs134	—	±	—		Cs134	0.1
White rockfish	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.7
				Cs134	—	±	—		Cs134	0.8
White rockfish (four fish)	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	OR	Cs137	0.7	±	0.06	0.7	Cs137	0.1
				Cs134	—	±	—		Cs134	0.09
Konjac	Japan (production)	Jun-25	OR	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.1
				Cs134	—	±	—		Cs134	0.1
Rice flour	Fukushima Pref.	Jun-25	OR	Cs137	0.5	±	0.1	0.5	Cs137	0.2
				Cs134	—	±	—		Cs134	0.2
Buckwheat flour	Kori, Date, Fukushima Pref.	Jun-25	OR	Cs137	0.8	±	0.1	0.8	Cs137	0.3
				Cs134	—	±	—		Cs134	0.3
Sea water (surface)	Souma Port/ Fukushima Pref.	Jun-25	CA	Cs137	0.006	±	0.001	0.006	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Murakami Coast/ Fukushima Pref.	Jun-25	CA	Cs137	0.006	±	0.001	0.006	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Ukedo Port/ Fukushima Pref.	Jun-25	CA	Cs137	0.005	±	0.001	0.005	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Futaba Beach/ Fukushima Pref.	Jun-25	CA	Cs137	0.01	±	0.001	0.01	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Iwasawa Beach/ Fukushima Pref.	Jun-25	CA	Cs137	0.004	±	0.001	0.004	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Yotsukura Port/ Fukushima Pref.	May-25	CA	Cs137	0.015	±	0.001	0.015	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Ena Port/ Fukushima Pref.	May-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Onahama Port/ Fukushima Pref.	May-25	CA	Cs137	0.004	±	0.001	0.004	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface)	Obama Port/ Fukushima Pref.	May-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface) Suspended solid	Souma Port/ Fukushima Pref.	Jun-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface) Suspended solid	Murakami Coast/ Fukushima Pref.	Jun-25	CA	Cs137	0.003	±	0.001	0.003	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface) Suspended solid	Ukedo Port/ Fukushima Pref.	Jun-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.001
				Cs134	—	±	—		Cs134	0.001
Sea water (surface) Suspended solid	Futaba Beach/ Fukushima Pref.	Jun-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.001
				Cs134	—	±	—		Cs134	0.001
Sea water (surface) Suspended solid	Iwasawa Beach/ Fukushima Pref.	Jun-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002
Sea water (surface) Suspended solid	Yotsukura Port/ Fukushima Pref.	May-25	CA	Cs137	0.008	±	0.001	0.008	Cs137	0.002
				Cs134	—	±	—		Cs134	0.001
Sea water (surface) Suspended solid	Ena Port/ Fukushima Pref.	May-25	CA	Cs137	—	±	—	Under Minimum Limit of Detection	Cs137	0.002
				Cs134	—	±	—		Cs134	0.002

※“—”used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0 (zero) Bq/kg.



★Gamma-ray

(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measuring instrument type	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Sea water (surface) Suspended solid	Onahama Port/ Fukushima Pref.	May-25	CA	Cs137	— Bq/L	± —	Bq/L	Under Minimum Limit of Detection	Cs137	0.002 Bq/L
				Cs134	— Bq/L	± —	Bq/L		Cs134	0.002 Bq/L
Sea water (surface) Suspended solid	Obama Port/ Fukushima Pref.	May-25	CA	Cs137	— Bq/L	± —	Bq/L	Under Minimum Limit of Detection	Cs137	0.002 Bq/L
				Cs134	— Bq/L	± —	Bq/L		Cs134	0.002 Bq/L
Bamboo grass	Fukushima Pref.	Jun-25	OR	Cs137	38.3 Bq/kg raw	± 1.3	Bq/kg raw	38.3	Cs137	1.0 Bq/kg raw
				Cs134	— Bq/kg raw	± —	Bq/kg raw		Cs134	1.0 Bq/kg raw
Rose of sharon	Izumigaoka, Iwaki	Jul-25	OR	Cs137	1.7 Bq/kg raw	± 0.2	Bq/kg raw	1.7	Cs137	0.4 Bq/kg raw
				Cs134	— Bq/kg raw	± —	Bq/kg raw		Cs134	0.3 Bq/kg raw
Soil(in the park)	Soutoucho Park, Tairayagawase, Iwaki	Mar-25	OR	Cs137	671.0 Bq/kg 乾	± 2.2	Bq/kg 乾	679.2	Cs137	0.7 Bq/kg 乾
				Cs134	8.2 Bq/kg 乾	± 0.4	Bq/kg 乾		Cs134	0.7 Bq/kg 乾
Soil(in the park)	Soutoucho Park, Tairayagawase, Iwaki	Mar-25	CA	Cs137	666.6 Bq/kg 乾	± 3.0	Bq/kg 乾	675.7	Cs137	1.0 Bq/kg 乾
				Cs134	9.1 Bq/kg 乾	± 0.4	Bq/kg 乾		Cs134	1.2 Bq/kg 乾
Soil(in the park)	Soutoucho Park, Tairayagawase, Iwaki	Mar-25	CA	Cs137	447.5 Bq/kg 乾	± 1.8	Bq/kg 乾	454.0	Cs137	0.6 Bq/kg 乾
				Cs134	6.5 Bq/kg 乾	± 0.3	Bq/kg 乾		Cs134	0.8 Bq/kg 乾
Soil(in the park)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	OR	Cs137	1216.3 Bq/kg 乾	± 26.2	Bq/kg 乾	1231.8	Cs137	8.8 Bq/kg 乾
				Cs134	15.5 Bq/kg 乾	± 4.8	Bq/kg 乾		Cs134	8.8 Bq/kg 乾
Soil (park entrance)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	CA	Cs137	873.8 Bq/kg 乾	± 3.2	Bq/kg 乾	885.5	Cs137	0.9 Bq/kg 乾
				Cs134	11.7 Bq/kg 乾	± 0.4	Bq/kg 乾		Cs134	1.0 Bq/kg 乾
Soil (in the park)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	OR	Cs137	804.7 Bq/kg 乾	± 3.6	Bq/kg 乾	815.2	Cs137	1.1 Bq/kg 乾
				Cs134	10.5 Bq/kg 乾	± 0.7	Bq/kg 乾		Cs134	1.2 Bq/kg 乾
Soil (horizontal bar)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	CA	Cs137	766.8 Bq/kg 乾	± 3.0	Bq/kg 乾	778.5	Cs137	0.9 Bq/kg 乾
				Cs134	11.7 Bq/kg 乾	± 0.4	Bq/kg 乾		Cs134	1.1 Bq/kg 乾
Soil (bench)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	CA	Cs137	748.1 Bq/kg 乾	± 2.6	Bq/kg 乾	758.2	Cs137	0.7 Bq/kg 乾
				Cs134	10.1 Bq/kg 乾	± 0.3	Bq/kg 乾		Cs134	0.9 Bq/kg 乾
Soil(in the park)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	OR	Cs137	593.9 Bq/kg 乾	± 7.5	Bq/kg 乾	602.1	Cs137	2.9 Bq/kg 乾
				Cs134	8.2 Bq/kg 乾	± 1.5	Bq/kg 乾		Cs134	2.7 Bq/kg 乾
Soil(in the park)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	OR	Cs137	583.5 Bq/kg 乾	± 2.7	Bq/kg 乾	590.8	Cs137	0.9 Bq/kg 乾
				Cs134	7.3 Bq/kg 乾	± 0.5	Bq/kg 乾		Cs134	1.0 Bq/kg 乾
Soil (pavilion)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	OR	Cs137	323.4 Bq/kg 乾	± 1.6	Bq/kg 乾	326.7	Cs137	0.6 Bq/kg 乾
				Cs134	3.3 Bq/kg 乾	± 0.4	Bq/kg 乾		Cs134	0.7 Bq/kg 乾
Soil (park entrance)	Nagamachi Park, Uchigoumimaya, Iwaki	Mar-25	OR	Cs137	284.1 Bq/kg 乾	± 1.6	Bq/kg 乾	287.6	Cs137	0.7 Bq/kg 乾
				Cs134	3.5 Bq/kg 乾	± 0.4	Bq/kg 乾		Cs134	0.7 Bq/kg 乾
Fallout	Haramachi, Minamisouma, Fukushima Pref.	Jun-25	OR	Cs137	2199.2 Bq/kg 乾	± 8.3	Bq/kg 乾	2226.7	Cs137	3.2 Bq/kg 乾
				Cs134	27.5 Bq/kg 乾	± 1.8	Bq/kg 乾		Cs134	3.2 Bq/kg 乾
Lichen	Haramachi, Minamisouma, Fukushima Pref.	Jun-25	OR	Cs137	39856.0 Bq/kg 乾	± 67.1	Bq/kg 乾	40318.8	Cs137	13.8 Bq/kg 乾
				Cs134	462.8 Bq/kg 乾	± 10.0	Bq/kg 乾		Cs134	14.4 Bq/kg 乾

※“_”used in Measurement Result and Uncertainty shows that the value is below the detection limit.

But it does not necessary mean 0(zero)Bq/kg.



★Beta-ray

Measuring instrument		Feature
Liquid Scintillation Counter		
Product of Hidex HIDEX 300SLL	Product of PerkinElmer Japan Quantulus GCT 6220	Equipment for measuring low-energy beta-ray emission nuclides
		Measuring nuclide Strontium90 Half-life 30 years Organic bound tritium Half-life 12.3 years Free-water tritium Half-life 12.3 years All samples are measured in liquid condition after several days of pretreatment.

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Minimum Limit of Detection	
White rockfish	Souma Port/ Fukushima Pref.	Jun-24	T(Tissue free water)	Under Minimum Limit of Detection Bq/L	± - Bq/L	0.38 Bq/L		
Black seabastes	Sendai Bay/ Miyagi Pref.	Apr-25	T(Organic bound)	Under Minimum Limit of Detection Bq/Kg生	± - Bq/Kg生	0.10 Bq/Kg生		
Sea waterC (surface)	Off the coast of Fukushima Nuclear Power Plant1	Nov-24	T(free)	0.11 Bq/L	± 0.04 Bq/L	0.04 Bq/L		
Sea waterC (lower)	Off the coast of Fukushima Nuclear Power Plant1	Nov-25	T(free)	0.25 Bq/L	± 0.05 Bq/L	0.04 Bq/L		
Sea waterD (surface)	Off the coast of Fukushima Nuclear Power Plant1	Nov-24	T(free)	0.15 Bq/L	± 0.04 Bq/L	0.04 Bq/L		
Sea waterD (lower)	Off the coast of Fukushima Nuclear Power Plant1	Nov-24	T(free)	0.11 Bq/L	± 0.04 Bq/L	0.04 Bq/L		
Sea waterA (surface)	Off the coast of Fukushima Nuclear Power Plant1	Mar-25	T(free)	0.12 Bq/L	± 0.04 Bq/L	0.04 Bq/L		
Sea waterA (lower)	Off the coast of Fukushima Nuclear Power Plant1	Mar-25	T(free)	0.16 Bq/L	± 0.05 Bq/L	0.04 Bq/L		
Sea waterA (surface)	Kasumigaura/ Ibaraki Pref.	Oct-22	T(free)	0.14 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Sea waterA (lower)	Kasumigaura/ Ibaraki Pref.	Oct-22	T(free)	0.11 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Rain Water	Onahama, Iwaki	Jun-25	T(free)	0.58 Bq/L	± 0.06 Bq/L	0.04 Bq/L		
Tap water	Saku, Nagano Pref.	Jun-22	T(free)	0.09 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Tap water	Haramachi, Minamisouma, Fukushima Pref.	Jul-22	T(free)	0.10 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Tap water	Okuma, Futaba, Fukushima Pref.	Jul-22	T(free)	0.20 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Tap water	Rikuzentakata, Iwate Pref.	Jul-22	T(free)	0.15 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Tap water	Fujieda, Sizuoka Pref.	Jul-22	T(free)	0.14 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Tap water	Hidaka, Wakayama Pref.	Aug-22	T(free)	0.15 Bq/L	± 0.08 Bq/L	0.08 Bq/L		
Tap water	Tadami, Minamiaizu, Fukushima Pref.	May-23	T(free)	0.09 Bq/L	± 0.08 Bq/L	0.08 Bq/L		

(Bq/Kg raw:Weight of raw sample Bq/Kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measurement Result		Uncertainty		Minimum Limit of Detection	
Sea water (surface)	Iwasawa Beach/ Fukushima Pref.	Jun-25	Sr90	Under Minimum Limit of Detection Bq/L	± - Bq/L		0.0005 Bq/L	
Sea water (surface)	Ishinomakihamaich i/ Miyagi Pref.	Jun-25	Sr90	Under Minimum Limit of Detection Bq/L	± - Bq/L		0.0005 Bq/L	
Sea water (surface)	Sendai new port/ Miyagi Pref.	Jun-25	Sr90	Under Minimum Limit of Detection Bq/L	± - Bq/L		0.0006 Bq/L	
Sea waterC (surface)	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	Sr90	0.0013 Bq/L	± 0.0003 Bq/L		0.0004 Bq/L	
Sea waterC (lower)	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	Sr90	0.0009 Bq/L	± 0.0003 Bq/L		0.0004 Bq/L	
Sea waterD (surface)	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	Sr90	Under Minimum Limit of Detection Bq/L	± - Bq/L		0.0006 Bq/L	
Sea waterD (lower)	Off the coast of Fukushima Nuclear Power Plant1	Jul-25	Sr90	Under Minimum Limit of Detection Bq/L	± - Bq/L		0.0006 Bq/L	
Sea water (surface)	Tomiooka Port/ Fukushima Pref.	Jul-25	Sr90	Under Minimum Limit of Detection Bq/L	± - Bq/L		0.0005 Bq/L	

Measurement results of 16 items by germanium semiconductor detector


Dr.Tetsuji Imanaka, Institute of Multiple Nuclear Science, Kyoto University

In order to convey more measurement results to everyone, we have asked Dr. Tetsuji Imanaka of the Institute of Advanced Nuclear Science, Kyoto University, to measure low-dose samples using germanium semiconductor detectors. Measurement samples are not only from Fukushima Prefecture but also come from other prefectures. Please compare data based on measurements from various regions and use them to protect your children from radiation exposure.

★Gamma-ray

Measuring instrument: Germanium Semiconductor detector

- Product of CANBERRA(CA),USA GX3018 Relative efficiency 30% or more
- Product of ORTEC(OR),USA GMX25-70 Relative efficiency 35%




(Bq/kg raw:Weight of raw sample Bq/kg dry:Weight of dried sample)

Samples	Sampling Point	Sampling Month	Measuring instrument type	Measurement Result		Uncertainty		Total Amount of Cesium	Minimum Limit of Detection	
Brown rice	Namiie,Futaba, Fukushima Pref.	Oct-24	CA	Cs137	4.5 Bq/kg raw	±	0.09 Bq/kg raw	0.45	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Japanese mustard spinach	Iwaki City	Apr-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.13 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Japanese parsley	Hitachiomiya, Ibaraki Pref.	May-25	OR	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.17 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Koshiabara (wild)	Miwa,Iwaki	Apr-25	OR	Cs137	160 Bq/kg raw	±	0.7 Bq/kg raw	161.60	Cs137	Bq/kg raw
				Cs134	1.6 Bq/kg raw	±	0.08 Bq/kg raw		Cs134	Bq/kg raw
Wild bamboo shoot(boiled)	Ibaraki Pref.	Apr-25	CA	Cs137	26 Bq/kg raw	±	0.2 Bq/kg raw	26.22	Cs137	Bq/kg raw
				Cs134	0.22 Bq/kg raw	±	0.03 Bq/kg raw		Cs134	Bq/kg raw
Butterbur	Hitachiomiya, Ibaraki Pref.	Apr-25	OR	Cs137	0.06 Bq/kg raw	±	0.02 Bq/kg raw	0.06	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Mountain udo	Iwaki City	May-25	OR	Cs137	0.07 Bq/kg raw	±	0.03 Bq/kg raw	0.07	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Ostrich fern	Hitachiomiya, Ibaraki Pref.	May-25	CA	Cs137	0.42 Bq/kg raw	±	0.03 Bq/kg raw	0.42	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Aralia sprout	Yamagata Pref.	Apr-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.12 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
White rockfish	Sendai Bay/ Miyagi Pref.	Apr-25	OR	Cs137	0.26 Bq/kg raw	±	0.06 Bq/kg raw	0.26	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Shitake mushroom grown in bacteria-bed(raw)	Minamisanriku, Miyagi Pref.	Apr-25	CA	Cs137	0.97 Bq/kg raw	±	0.07 Bq/kg raw	0.97	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Shitake mushroom grown in bacteria-bed	Niigata Pref.	Apr-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.6 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Wood ear mushroom	Ibaragki Pref.	May-25	OR	Cs137	1.3 Bq/kg raw	±	0.06 Bq/kg raw	1.3	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
White oyster mushroom(dried)	Siroato, Higashiibaraki, Ibaraki Pref.	May-25	CA	Cs137	8.0 Bq/kg raw	±	0.9 Bq/kg raw	8.0	Cs137	Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
White mushroom	Ibaragki Pref.	May-25	CA	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.16 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw
Green pepper	Miyazaki Pref.	May-25	OR	Cs137	— Bq/kg raw	±	— Bq/kg raw	Under Minimum Limit of Detection	Cs137	0.14 Bq/kg raw
				Cs134	— Bq/kg raw	±	— Bq/kg raw		Cs134	Bq/kg raw

Air dose rate July 2025

Measuring Instrument		Measuring Place
CsI Scintillation survey meter	NaI Scintillation survey meter	Yokocho Park, Onahama, Iwaki, Fukushima
⑧HITACHI ALOKA TCS-1172	⑦HORIBA Radi PA-1100	
		
Feature : Measuring air (space) radiation dose and radioactive surface contamination of human body and other things.		



Measuring instrument		HORIBA Radi	
Measuring Date	Weather	Near the surface of the ground (μSv/h)	1m above the ground (μSv/h)
2025/7/1	☀	0.064	0.055
2025/7/2	☀	0.068	0.063
2025/7/3	☀	0.064	0.053
2025/7/4	☀	0.062	0.053
Measuring Date	Weather	Near the surface of the ground (μSv/h)	1m above the ground (μSv/h)
2025/7/7	☀	0.059	0.056
2025/7/8	☀	0.064	0.053
2025/7/9	☀	0.064	0.058
2025/7/10	☁	0.063	0.060
2025/7/11	☀	0.064	0.052
Measuring Date	Weather	Near the surface of the ground (μSv/h)	1m above the ground (μSv/h)
2025/7/14	☂	0.066	0.054
2025/7/15	☂	0.066	0.058
2025/7/16	☁/☂	0.061	0.052
2025/7/17	☀	0.062	0.053
2025/7/18	☀	0.061	0.052
Measuring Date	Weather	Near the surface of the ground (μSv/h)	1m above the ground (μSv/h)
2025/7/22	☀	0.071	0.060
2025/7/23	☀	0.069	0.058
2025/7/24	☀	0.065	0.058
2025/7/25	☀	0.062	0.052
Measuring Date	Weather	Near the surface of the ground (μSv/h)	1m above the ground (μSv/h)
2025/7/28	☀	0.063	0.061
2025/7/29	☀	0.066	0.055
2025/7/30	☀	0.064	0.053
2025/7/31	☀	0.061	0.053